

100-1178

02-16-2011

1/45



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Betty Brame
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, N.C. 27419

FEB 16 2011

Subject: Application for Pesticide Notification (PRN 98-10)
Submission date: 1/27/2011
Product Name: Quilt Fungicide
EPA Reg. No.: 100-1178
EPA Decision Number: 445138

Dear Betty Brame,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated April 28, 2008. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.


The Agency acknowledges your request for the alternate brand name "Azoxypop Fungicide".

The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have questions concerning this letter, please contact Shaunta Hill at 703-347-8961 or myself at 703-308-3194.

Sincerely,

A handwritten signature in cursive script that reads "Shaja Brothers-Joyner".

Shaja Brothers-Joyner
Product Manager 20
Fungicide Branch
Registration Division (7505P)

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-------------------------------

Application for Pesticide - Section I

1. Company/Product Number 100-1178	2. EPA Product Manager Shaja B. Joyner	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) AzoxyProp Fungicide	PM# 20	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II


<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.
 Per PR Notice 98-10, Syngenta Crop Protection, LLC is submitting Notification of an alternate brand name for Quilt® Fungicide, EPA Reg. No. 100-1178. The alternate brand name for Quilt Fungicide will be AzoxyProp Fungicide. Enclosed is one copy of the label with the alternate brand name highlighted, a clean label copy, a CD of the label with the alternate brand name for electronic comparison, a Certificate With Respect to Label Integrity form, and EPA Form 8570-1.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No			<input checked="" type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
*Certification must be submitted		If "Yes" No. per Unit Packaging wgt. Container	If "Yes" No. per Unit Packaging wgt. container		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 30, 264 gallons and Bulk		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other <u>Pressure Sensitive</u> <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Betty Brame	Title Regulatory Specialist II	Telephone No. (Include Area Code) 336-632-7466			
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory Specialist II			
4. Typed Name Betty Brame		5. Date January 27, 2011			

5
45

AzoxyProp

GROUP 3 11 FUNGICIDES

Fungicide

Broad spectrum fungicide for control of plant diseases

Active Ingredient:

Azoxystrobin 7.0%

Propiconazole 11.7%

Other Ingredients: 81.3%

Total: 100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1178

EPA Est.

SCP 1178A

_____ gallons
Net Contents

NOTIFICATION
FEB 16 2011

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- Protective eyewear

Remove and wash contaminated clothing before reuse.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Do not apply directly to water except as specified on this label. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify state and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- Protective eyewear

GENERAL INFORMATION

AzoxyProp Fungicide is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. AzoxyProp may also improve the yield and/or quality of the crop even under limited disease pressure due to plant performance characteristics. These additional benefits are due to positive effects on plant physiology. The effects may vary according to other factors such as the crop, crop hybrid, or environment. AzoxyProp may be applied as a foliar spray in alternating spray programs or in

tank mixes with other crop protection products. All applications should be made according to the use directions that follow.

GENERAL USE PRECAUTIONS

Do not use in nurseries, greenhouses or landscape plantings.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult: http://128.104.224.199/espp_front/view.jsp. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

GENERAL USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of AzoxyProp has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

Rotational Crops: Soybeans may be planted as a double crop following a cereal crop which has been treated with AzoxyProp. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding. Oats, rye and wild rice may be planted 45 days after AzoxyProp application. To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of AzoxyProp application to the preceding crop, unless the second crop appears on this label.

Crop Tolerance/Phytotoxicity: AzoxyProp Fungicide demonstrates some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone can contribute to phytotoxicity. Under certain environmental conditions, tank mixes of AzoxyProp plus herbicides and/or fertilizers may cause crop injury in barley, triticale and wheat.

ATTENTION

AzoxyProp is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray AzoxyProp where spray drift may reach apple trees.

C

C

$\frac{11}{45}$

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply AzoxyProp to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Spray Drift Management: To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. More information on managing spray drift can be found on the Syngenta Crop Protection website under Stewardship:

http://www.syngentacropprotectionus.com/enviro/driftmanagement/index.asp?nav=drift_management

Integrated Pest Management: AzoxyProp should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The SPECIFIC USE DIRECTIONS section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area. AzoxyProp may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

GROUP 3 11 FUNGICIDES

AzoxyProp Fungicide is a mixture of Group 3 (propiconazole) and Group 11 (azoxystrobin) fungicides. AzoxyProp has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. SYNGENTA encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the specific crop recommendations that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups. In situations requiring multiple sprays, develop season long spray programs for Group 11 (QoI) fungicides. The program should meet the goal of no more than 1/3 of the total sprays per season, when a Group 11 fungicide is used as a solo product, or 1/2 the total sprays when a Group 11 fungicide

is used in a mixture. Programs that include both solo Group 11 products and/or mixes containing Group 11 products should be no more than ½ the total sprays.

AzoxyProp should not be alternated or tank mixed with any fungicide to which resistance has already developed.

SPRAY EQUIPMENT

AzoxyProp may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be *16-mesh or coarser*.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - (1) Maintain 35-40 psi at nozzles
 - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

MIXING INSTRUCTIONS

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

AzoxyProp Alone (no tank mix):

- Add ½-¾ of the required amount of water to the spray or mixing tank.
- With the agitator running, add AzoxyProp to the tank.
- Continue agitation while adding the remainder of the water.

14
45

- Begin application of the spray solution after AzoxyProp has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

AzoxyProp + Tank Mixtures: AzoxyProp is usually compatible with all tank mix partners listed on this label. Do not combine AzoxyProp in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious to the crop under your conditions of use. To determine the physical compatibility of AzoxyProp with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables (suspoemulsions), and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank mix partner(s) into the tank in the same order as described above in the "AzoxyProp + Tank Mixtures" section.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the AzoxyProp to the spray tank.
- Allow AzoxyProp to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.
- No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product may not be mixed with any product which prohibits such mixing.

APPLICATION PROCEDURES

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner that will result in exposure to humans or animals.

Ground Application:

- For field crops (non-trees), apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- For field crops (non-trees), apply in a minimum spray volume of 2 gallons per acre unless specified otherwise.
- For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- Do not apply via ULV.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches per acre of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Note: Do not inject AzoxyProp at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part AzoxyProp. AzoxyProp is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

APPLICATION INSTRUCTIONS

Apply AzoxyProp Fungicide at rates and timings as described in this label.

Directions for Use Through Sprinkler and Drip Chemigation Systems (apply AzoxyProp only to crops for which chemigation is specified on this label):

Spray Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Use Precautions for Sprinkler and Drip Irrigation Applications

Sprinkler Irrigation: Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, drip (trickle), or hand move irrigation systems. Do not apply this product through any other type of irrigation system except as specified on this label.

Apply with center pivot or continuous-move equipment distributing ½ acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Do not apply when wind speed favors drift beyond the area intended for treatment. Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Directions for Use

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Almonds	Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	14 - 27.5	Apply AzoxyProp at early bloom stage. If disease pressure is low, a second application of 14 fl. oz./A may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50-75% bloom and petal fall. AzoxyProp may only be used on only two blossom blight applications. Orbit may be used alone for one of these applications.
	Anthracnose (<i>Collectotrichum acutatum</i>)	14 – 27.5	Apply AzoxyProp beginning at bud break on a 7-14 day interval. Make no more than two consecutive applications before switching to another fungicide with a different mode of action.

Application Instructions: Make no more than two sequential applications of a Group 11 fungicide prior to alternating with two applications of another product with a different mode of action than Group 11 fungicides. Almond diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. AzoxyProp may be applied by ground or by air (minimum of 15 gals./A). Aerial application may be used if necessary but disease control may be reduced. AzoxyProp may be applied by air only at growth stages prior to and including 5 weeks after petal fall.

CHANGE ON NEXT AMENDMENT – INSTRUCTIONS ON ALL OTHER CROPS.

Specific Use Restrictions:

- 1) Do not apply more than 110 fl. oz./A/season of AzoxyProp per crop.
- 2) Do not apply more than 0.9 lb. a.i. of propiconazole containing products.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 60 days of harvest.
- 5) Do not graze livestock in treated areas or cut treated cover crop for feed.

5/6

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Berries, bushberry sub-group Blueberry Currant Elderberry Gooseberry Huckleberry	Mummyberry <i>(Monilinia vaccinicorymbosi)</i>	14 - 20.5	Make first application of AzoxyProp beginning at green tip and repeat in 7-10 days. If conditions are favorable for disease development, additional application may need to be made at pink bud and repeating every 7 to 10 days through petal fall. Do not apply more than two consecutive applications before alternating to a non-Group 11 containing fungicide.
Juneberry Lingonberry Salal And cultivars and/or hybrids of these	Leaf Spot and Stem Canker <i>(Septoria albopunctata)</i> Powdery Mildew <i>(Microsphaera vaccinii)</i> Leaf Spot <i>(Septoria spp.)</i> Phomopsis Twig Blight, Fruit Rot, and Stem Canker <i>(P. vaccinii)</i> Botryosphaeria Canker <i>(Botryosphaeria spp.)</i>	14 - 20.5	AzoxyProp applications should begin prior to disease development and continue throughout the season on a 14 day interval. Make no more than two consecutive sprays before alternating to a non-Group 11 fungicide. Make no more than 3 applications per crop of AzoxyProp on other Group 11 fungicides.

Application Instructions: AzoxyProp may be applied by ground (minimum of 15 gals./A) or by air.

Specific Use Restrictions:

- 1) Do not apply more than 103 fl. oz./A/season of AzoxyProp per crop.
- 2) Do not apply more than 0.84 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 0.75 lb. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 30 days of harvest.

19/45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Berries, Caneberry sub-group Blackberry Bingleberry Boysenberry Dewberry Loganberry Lowberry Marionberry Olallieberry Raspberry, red and black Youngberry And cultivars and/or hybrids of these.	Rust <i>(Pucciniastrum vaccinii)</i> Leaf and Cane Spot <i>(Septoria rubi)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i> Leaf Spot <i>(Septoria spp.)</i>	14 - 20.5	AzoxyProp applications should begin prior to disease development and continue throughout the season on a 7-14 day interval. Make no more than two consecutive sprays before alternating to a non-Group 11 fungicide. Make no more than 3 applications per crop of AzoxyProp on other Group 11 fungicides.

Application Instructions: AzoxyProp may be applied by ground (minimum of 15 gals./A) or by air.

Specific Use Restrictions:

- 1) Do not apply more than 103 fl. oz./A/season of AzoxyProp per crop.
- 2) Do not apply more than 0.84 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 30 days of harvest.

2/12

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Carrots	Early Blight (<i>Cercospora carotae</i>) Powdery Mildew (<i>Erysiphe polygoni</i>) Suppression of Late Blight (<i>Alternaria dauci</i>)	14	Apply 14 oz. of AzoxyProp when conditions favor disease development. Continue applications on a 7-10 day interval, using the shorter interval when disease conditions are severe. Make no more than one application before alternating to a non-Group 11 fungicides.

Application Instructions: AzoxyProp may be applied by ground (minimum of 15 gals./A) or aerial application (minimum of 5 gals./A).

Specific Use Restrictions:

- 1) Do not apply more than 55 fl. oz./A /season of AzoxyProp per crop.
- 2) Do not apply more than 0.45 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 2.0 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 14 days of harvest.

21/45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Celery	Early Blight (<i>Cercospora apii</i>) Late Blight (<i>Septoria apicola</i>)	14	Apply AzoxyProp on a 7-10 day schedule in alternation with Tilt® alone or another product with a different mode of action than Group 11 fungicides.

Application Instructions: AzoxyProp may be applied by ground, air (5 gals./A minimum) or chemigation.

Specific Use Restrictions:

- 1) Do not apply within 14 days of harvest.
- 2) Do not apply more than 56 fl. oz./A/season.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 4) Do not apply more than 1.5 lbs. a.i. azoxystrobin containing products/A/season.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Wheat See next section for other cereals	Early season suppression of: Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Glume Blotch <i>(Stagonospora nodorum)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	7 - 14	Apply AzoxyProp in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.
	Control of Leaf Diseases: Rust <i>(Puccinia spp.)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Glume Blotch <i>(Stagonospora nodorum)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i> Helminthosporium Leaf Blight <i>(Drechslera tritici-repentis)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Net Blotch <i>(Pyrenophora teres)</i> Kernel Blight <i>(Alternaria spp.)</i>	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when AzoxyProp is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz./A may be applied. Applications may be made no closer than a 14-day interval. AzoxyProp can be applied through full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of AzoxyProp plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 2 applications/A/season.
- 2) Do not apply after Feekes 10.5.
- 3) Do not harvest within 30 days for forage or hay.
- 4) Do not apply more than 28.0 fl. oz./A/season of AzoxyProp.
- 5) Do not apply more than 0.22 lb. a.i. propiconazole containing products/A/season.
- 6) Do not apply more than 0.40 lb. a.i. azoxystrobin containing products/A/season.
- 7) Under certain environmental conditions, tank mixes of AzoxyProp plus herbicides and/or fertilizers may cause crop injury.

5/2

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Barley Triticale	Early season suppression of: Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	7 - 14	Apply AzoxyProp in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.
	Control of Leaf Diseases: Rust (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) Helminthosporium Leaf Blight (<i>Drechslera tritici-repentis</i>) Spot Blotch (<i>Bipolaris sorokiniana</i>) Barley Scald (<i>Rhynchosporium secalis</i>) Barley Stripe (<i>Pyrenophora graminea</i>) Net Blotch (<i>Pyrenophora teres</i>) Kernel Blight (<i>Alternaria</i> spp.)	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when AzoxyProp is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz./A may be applied. Applications may be made no closer than a 14-day interval. AzoxyProp can be applied up to 45 days prior to harvest.
	Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14	Apply full rate of AzoxyProp plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 2 applications/A/season.
- 2) Do not apply within 45 days of harvest¹
- 3) Do not harvest within 30 days for forage or hay.
- 4) Do not apply more than 28.0 fl. oz./A/season of AzoxyProp.
- 5) Do not apply more than 0.22 lb. a.i. propiconazole containing products/A/season.
- 6) Do not apply more than 0.40 lb. a.i. azoxystrobin containing products/A/season.
- 7) Under certain environmental conditions, tank mixes of AzoxyProp plus herbicides and/or fertilizers may cause crop injury²

57/2
42

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
<p>Citrus Fruit (non-bearing)</p> <p>Calamondin Citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma mandarin Tangerine</p> <p>Including all cultivars and/or hybrids of these</p>	<p>Greasy Spot (<i>Mycosphaerella citri</i>)</p>	<p>20.5 - 27.5</p>	<p>Apply AzoxyProp up to 2 times during June, July, or August. The other sprays may be Orbit® alone or another product with a different mode of action than Group 11 fungicides.</p> <p>AzoxyProp may be applied by ground, air or chemigation.</p> <p>Do not use AzoxyProp in citrus plant propagation nurseries.</p>

Specific Use Restrictions:

- 1) To avoid possible illegal residues, do not apply to citrus that will bear harvestable fruit within 12 months.
- 2) Do not apply more than 0.68 lb. a.i. propiconazole containing products/A/year.
- 3) Do not apply more than 1.5 lbs. a.i. azoxystrobin containing products/A/year.

5/5

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Corn Field Pop (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium leaf blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)	7 - 14	<p>Note: Applications prior to tasseling may impose stress on the plant that could inhibit proper kernel development, especially under environmentally stressful conditions.</p> <p>For leaf blights apply AzoxyProp when disease first appears. Continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.</p> <p>Apply no more than 2 applications of AzoxyProp or any other Group 11 fungicide per year.</p>
	Anthracnose leaf blight (<i>Colletotrichum graminicola</i>) Rusts (<i>Puccinia</i> spp.) Gray Leaf Spot (<i>Cercospora zea-maydis</i>) Eye Spot (<i>Aureobasidium zea</i>)	10.5 - 14	<p>Apply AzoxyProp when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.</p> <p>Apply no more than 2 applications of AzoxyProp or any other Group 11 fungicide per year.</p>

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply within 30 days of harvest for forage, grain, or stover.
- 2) Do not apply more than 56 fl. oz./A/season of AzoxyProp.
- 3) Do not apply more than 28 fl. oz. (0.224 lb. a.i. propiconazole) for field corn harvested for forage.
- 4) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 5) Do not apply more than 2.0 lbs. a.i. azoxystrobin containing products/A/season.

26
45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Corn, Sweet Sweet corn (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium leaf blights (<i>H. maydis, H. turcicum, H. carbonum</i>)	7 - 14	For leaf blights apply AzoxyProp when disease first appears. Continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. Alternate applications of AzoxyProp with Tilt or another product with a different mode of action than Group 11 fungicides.
	Anthracnose leaf blight (<i>Colletotrichum graminicola</i>) Rusts (<i>Puccinia</i> spp.) Gray Leafspot (<i>Cercospora zeae-maydis</i>) Eye Spot (<i>Aureobasidium zeae</i>)	10.5 - 14	Apply AzoxyProp when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Alternate applications of AzoxyProp with Tilt or another product with a different mode of action than Group 11 fungicides.

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl. oz./A/season of AzoxyProp.
- 2) Do not apply to sweet corn within 14 days of harvest for ears or forage.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 4) Do not apply more than 2.0 lbs. a.i. azoxystrobin containing products/A/season.

22/5

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cranberries	Cottonball (<i>Monilinia oxycocci</i>)	14 – 20.5	Make the first application at leaf bud break and repeat in 14 days. A third application should be made at early bloom and again in 14 days. Make no more than 2 consecutive sprays before alternating to a non-Group 11 fungicide. For resistance management, make no more than 3 sprays per season using any Group 11 fungicide. Under severe pressure, use the higher rate for control.

Application Instructions: AzoxyProp may be applied by ground (minimum of 10 gals./A) or aerial application (minimum of 20 gals./A).

Specific Use Restrictions:

- 1) Do not apply more than 82.4 fl. oz./A of AzoxyProp per crop.
- 2) Do not apply more than 0.67 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 45 days of harvest.
- 5) Do not treat cranberry fields used for aquaculture of fish and crustacean.
- 6) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators should use care in making applications near non-target aquatic habitats.
- 7) Do not apply to flooded crop.
- 8) Do not allow release of irrigation of flood water to non-target aquatic habitat for at least 14 days after the last application. Use is limited to Oregon, Washington, and Wisconsin only.
- 9) Use is limited to Oregon, Washington, and Wisconsin only.

File

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Filberts	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	14 – 27.5	Begin applications when green leaf tissue becomes visible and continue on a 2-3 week interval. Under severe disease conditions, use the higher rate and shorter interval. NOTE: On certain varieties, AzoxyProp applications may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to AzoxyProp treatments.

Application Instructions: AzoxyProp may be applied by ground or aerial application (minimum of 15 gals./A).

Specific Use Restrictions:

- 1) Do not apply more than 110 fl. oz./A of AzoxyProp per crop.
- 2) Do not apply more than 0.9 lb. a.i. of propiconazole containing products/A/season
- 3) Do not apply more than 1.2 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 60 days of harvest.
- 5) Do not graze livestock in treated areas or cut treated cover crop for feed.

5/12

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Grasses (grown for seed)	Rusts (<i>Puccinia</i> spp.) Powdery Mildew (<i>Erysiphe graminis</i>) Selenophoma Stem Eyespot (<i>Selenophoma</i> spp.) Ergot Stem Diseases	14 - 27.5	Apply AzoxyProp when powdery mildew infections, <i>Selenophoma</i> infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 27.5 fl. oz./A (except bluegrass apply 14 fl. oz./A) and make applications at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin application early in the growing season. Make no more than 2 sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.

Application Instructions: Apply AzoxyProp in a minimum of 20 gals. of water per acre for ground application, or in a minimum of 10 gals. of water per acre for aerial application. AzoxyProp may also be applied by chemigation.

Specific Use Restrictions:

- 1) Do not feed hay cut within 20 days of the last application.
- 2) Do not graze treated areas within 140 days of the last application.
- 3) Do not apply more than 110 fl. oz./A/season of AzoxyProp.
- 4) Do not apply more than 0.90 lb. a.i. propiconazole containing products/A/season.
- 5) Do not apply more than 0.75 lb. a.i. azoxystrobin containing products/A/season.
- 6) Use is limited to Idaho, Minnesota, Nebraska, Oregon, and Washington only.

30
45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Mint Peppermint Spearmint	Rust (<i>Puccinia menthae</i>) Powdery Mildew (<i>Erysiphe</i> spp.)	14	Begin applications when the plants are 2-4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

Application Instructions: AzoxyProp may be applied by ground (minimum of 20 gals./A), air (minimum of 5 gals./A), or chemigation.

CHANGE ON NEXT AMENDMENT – INSTRUCTIONS ON ALL OTHER CROPS.

Specific Use Restrictions:

- 1) Do not apply more than 27.6 fl. oz./A /season of AzoxyProp per crop.
- 2) Do not apply more than 0.22 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 0.75 lb. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 30 days of harvest.

3/15

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Onions Onions (dry) Onions (green) Garlic Scallions Shallots And/or cultivars or hybrids of these. See below for additional onion types.	Purple Blotch (<i>Alternaria porri</i>) Suppression of Botrytis Leaf Blight (<i>Botrytis squamosa</i>) Rust (<i>Puccinia allii</i>) Cladoporium Leaf Blotch (<i>C. allii</i>)	14 – 27.5	Begin applications when conditions favor disease development and continue on a 7-10 day interval. Use the higher rate and shorter interval when disease conditions are severe. Make only 1 application before alternating to a non-Group 11 fungicide.

Onions (dry bulb): Shallots (dry bulb)

Onions (green): Shallots (green), Eschalots (green), Japanese bunching onions, Leeks, Spring onions

Application Instructions: AzoxyProp may be applied by ground (15 gals./A minimum) or aerial application (minimum of 5 gals./A).

Specific Use Restrictions:

- 1) Do not apply more than 55.3 fl. oz./A /season of AzoxyProp per crop.
- 2) Do not apply more than 0.45 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 14 days of harvest on dry bulb onions.
- 5) AzoxyProp may be applied at 0 day PHI (day of harvest) for green onion types.

3/25

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Peanuts	Early Leaf Spot <i>(Cercospora arachidicola)</i> Late Leaf Spot <i>(Cercosporidium personatum)</i> Rust <i>(Puccinia arachidis)</i> Web Blotch <i>(Phoma arachidicola)</i>	10.5 - 14	Apply AzoxyProp beginning 35-40 days after planting or at the first appearance of disease. Continue applications on a 14-day schedule. Under heavy disease pressure use higher recommended application rates. AzoxyProp also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply within 14 days of harvest.
- 2) Do not apply more than 56 fl. oz./A/season.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 4) Do not apply more than 0.80 lb. a.i. azoxystrobin containing products/A/season.

File

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Pecans	Pecan Scab (<i>Cladosporium caryigenum</i>) Anthracnose (<i>Glomerella cingulata</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae pv pecanae</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>) Powdery Mildew (<i>Microsphaera penicillata</i>)	14 – 27.5	Pecan scab: Apply 14 – 27.5 fl. oz./A AzoxyProp on a 14 day schedule during bud break and pre-pollination sprays. Apply 20 – 28 fl. oz./A during nut formation and cover sprays. Use higher rates when disease pressure is heavier. Do not apply after shuck split. Other foliar diseases: AzoxyProp may be applied for control of mid to late season foliar diseases at 14 – 20.5 fl. oz./A with other pecan products labeled for these diseases. Observe all directions, precautions, and limitations for the other products. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides. Do not apply after shuck split.

Application Instructions: AzoxyProp may be applied by ground or air (minimum of 20 gals./A). Ground applications should be applied in sufficient water to provide for full coverage.

Specific Use Restrictions:

- 1) Do not apply more than 122 fl. oz./A/season of AzoxyProp.
- 2) Do not apply after shuck split or within 45 days of harvest, whichever is first.
- 3) Do not graze livestock in treated areas or cut treated cover crops for feed.
- 4) Do not apply more than 0.9 lb. a.i. propiconazole containing products/A/season.
- 5) Do not apply more than 1.2 lbs. a.i. azoxystrobin containing products/A/season.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a bulletin consult http://128.104.224.199/espp_front/view.jsp. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

34
45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Pistachios	Botryspheeria Panicle and Shoot Blight (<i>B. dothidea</i>)	14 – 27.5	Begin applications when green leaf tissue becomes visible and continue on a 14-21 day interval. Under severe disease conditions, use the higher rate and the shorter interval. Make no more than 2 consecutive applications of AzoxyProp before switching to another non-Group 11 fungicide.

Application Instructions: AzoxyProp may be applied by ground or aerial application (15 gals./A minimum).

Specific Use Restrictions:

- 1) Do not apply more than 110 fl. oz./A/season of AzoxyProp.
- 2) Do not apply more than 0.9 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 60 days of harvest.
- 5) Do not graze livestock in treated areas or cut treated cover crop for feed.

25/25

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Rice	Sheath Blight <i>(Rhizoctonia solani)</i> Brown Leaf Spot <i>(Helminthosporium oryzae)</i> Narrow Brown Leaf Spot <i>(Cercospora oryzae)</i> Leaf Smut <i>(Entyloma oryzae)</i> Sheath Spot <i>(Rhizoctonia oryzae)</i> Kernel Smut <i>(Tilletia barclayana)</i> Aggregate Sheath Spot <i>(Rhizoctonia oryzae-sativa)</i> Leaf Blast <i>(Pyricularia grisea)</i> Black Sheath Rot <i>(Gaeumannomyces graminis)</i> Stem Rot <i>(Sclerotium oryzae)</i> For disease suppression of: False Smut <i>(Ustilaginoidea virens)</i>	14 - 34.5	Timing of AzoxyProp application will depend on disease severity, disease complex and rice variety/growth stage. Consult local extension experts for local economic thresholds established for various rice varieties and diseases. AzoxyProp should be applied at 28-34.5 fl. oz./A prior to or in the early stages of disease development. If disease reappears, use another fungicide for the second application. For leaf blast control, use a minimum of 34.5 oz./A. For neck or panicle blast, additional applications of Quadris® may be made (refer to Quadris label). Tank mix option: Apply 14-20.5 fl. oz./A of AzoxyProp in a tank mix with Quadris or other fungicides for control of rice diseases. A minimum of 0.15 lb. a.i./A azoxystrobin should be in the tank mix (see AzoxyProp rate conversion table at the end of the label). Make no more than 2 applications of a Group 11 fungicide per year.

Application Instructions: Applications may be made by air only. For aerial application, volumes should be 5-10 GPA. An adjuvant may be added at recommended rates to improve canopy coverage and penetration while reducing evaporation and drift.

Specific Use Restrictions:

- 1) Do not apply within 35 days of harvest.
- 2) Do not apply to stubble or ratoon crop rice.
- 3) Do not use in rice fields where commercial farming of crayfish will be practiced.
- 4) Do not drain water from treated rice fields into ponds used for commercial fish farming.
- 5) Do not use water drained from treated fields to irrigate other crops.
- 6) Do not apply more than 41 fl. oz./A/season of AzoxyProp.
- 7) Do not apply more than 0.34 lb. a.i. propiconazole containing products/A/season.
- 8) Do not apply more than 0.70 lb. a.i. azoxystrobin containing products/A/season.
- 9) Do not release floodwater within 14 days of an application.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a bulletin consult http://128.104.224.199/espp_front/view.jsp. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

5/2/93

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Sorghum	Anthracnose (<i>Colletotrichum graminicola</i>) Gray leaf spot (<i>Cercospora sorghi</i>) Ergot (<i>Claviceps sorghi</i>)	14	<p>For foliar diseases, AzoxyProp applications should begin prior to disease development. Applications may be made by ground, air or chemigation.</p> <p>For ergot: Make the first application at or just prior to flowering. Repeat on a 5-7 day interval. Apply up to 4 times.</p> <p>Resistance Management: Do not apply more than two sequential applications of AzoxyProp or other Group 11 containing fungicides before alternation with a fungicide that is not in Group 11.</p>

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 55 oz of AzoxyProp/A per season
- 2) Do not apply more than 0.22 lb propiconazole a.i./A (27.5 oz AzoxyProp) on sorghum harvested for forage.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing product/A/season.
- 4) For grain and stover, do not apply more than 0.75 lbs. a.i./A per season of azoxystrobin-containing products.
- 5) For forage, do not apply more than 0.5 lbs. a.i./A per season of azoxystrobin-containing products.
- 6) Do not graze livestock or cut for green chop or silage within 30 days of application.
- 7) Do not apply within 30 days of harvest for forage
- 8) Do not apply within 21 days of harvest for grain and stover.

5/5

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Soybeans	Aerial Web Blight <i>(Rhizoctonia solani)</i> Anthracnose <i>(Colletotrichum truncatum)</i> Brown Spot <i>(Septoria glycines)</i> Frogeye Leafspot <i>(Cercospora sojina)</i> Soybean Rust <i>(Phakopsora pachyrhizi)</i> Alternaria Leafspot <i>(Alternaria spp.)</i> Cercospora Blight and Leaf Spot <i>(C. kickuchii)</i> Pod and Stem Blight <i>(Diaporthe spp.)</i>	14 – 20.5	<p>Aerial Web Blight: Apply 14- 20.5 fl. oz./A at the first appearance of disease and repeat the application 14-21 days later. Under severe disease conditions use the higher rate and shorter interval.</p> <p>Other foliar diseases (except rust): Apply 20.5 fl. oz./A at growth stage R3 (early pod set when pods are 1/8-1/4 inch long) and 14-21 days later at growth stage R5 (pod fill).</p> <p>Soybean Rust: Apply 14-20.5 fl. oz./A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14-21 day interval. Use higher rate and shorter interval when disease is present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p> <p>On certain varieties, AzoxyProp applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to AzoxyProp treatments.</p>

Application Instructions: AzoxyProp is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. AzoxyProp may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl. oz./A/season of AzoxyProp.
- 2) Do not apply more than 0.34 lb. a.i. of propiconazole containing products/A/season.
- 3) Do not apply more than 1.5 lbs. a.i. of azoxystrobin containing products/A/season.
- 4) Do not apply within 21 days of harvest for seed and 0 days for forage and hay.

28
45

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Stone Fruits Apricot Cherry, sweet Cherry, tart Nectarine Peach Plum Plumcot Prune	Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	14	Apply AzoxyProp at early bloom stage. If disease pressure is low, a second application of 14 fl. oz./A may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50-75% bloom and petal fall. AzoxyProp may only be used on 2 of these applications. Orbit may be used alone for one of these applications.
Peach Plum	Brown Rot on Fruit (<i>Monilinia</i> spp.)	14	Apply as needed, a maximum of 2 sprays of AzoxyProp, during the preharvest period up to the day of harvest. Make the two applications no closer than 10 days apart.
Plumcot Prune Including all cultivars and hybrids of these	Powdery Mildew (<i>Podosphaera clandestine</i> , <i>Sphaerotheca pannosa</i>) Cherry Leaf Spot (<i>Blumeriella jaapii</i>)	14	Follow the blossom blight schedule. Make up to 2 additional applications on a 10-14 day interval from the end of petal fall to harvest.

Application Instructions: Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with 2 applications of another product with a different mode of action than Group 11 fungicides. Stone fruit diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. Aerial application (minimum of 15 gals./A) may be used if necessary but disease control may be reduced.

Applications of AzoxyProp during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply AzoxyProp to Stanley plums earlier than 21 days prior to harvest.

Specific Use Restrictions:

- 1) May be applied the day of harvest.
- 2) Do not apply more than 70 fl. oz./A/season of AzoxyProp.
- 3) Do not apply more than 0.56 lb. a.i. propiconazole containing products/A/season.
- 4) Do not apply more than 1.0 lb. a.i. azoxystrobin containing products/A/season.

5/5/19

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Tree Nuts See list below for tree nuts labeled	Foliar Diseases	14 - 27.5	Apply AzoxyProp at first sign of disease. Repeat on a 7-14 day interval. Do not make more than two consecutive applications before switching to a non Group 11 fungicide. Make no more than 4 applications of a AzoxyProp or other Group 11 fungicide containing product.
Additional tree nuts: Almond (see specific directions), Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (see specific directions), Hickory, Macadamia, Pecan (see specific directions), Pistachios (see specific directions), Walnut.			
Application Instructions: AzoxyProp may be applied by ground or aerial application (15 gals./A minimum). For best control of tree nut diseases, ground applications are recommended.			
Specific Use Restrictions: <ol style="list-style-type: none"> 1) Do not apply more than 110 fl. oz./A of AzoxyProp per crop. 2) Do not apply more than 0.9 lb. a.i. of propiconazole containing products/season. 3) Do not apply more than 1.2 lbs. a.i. of azoxystrobin containing products/A/season. 4) Do not apply within 60 days of harvest except for pecan (see specific use directions). 5) Do not graze livestock in treated areas or cut treated cover crop for feed. 			

4/5

AzoxyProp Rate Conversion Table

Fl. oz./A	Lb. a.i./A propiconazole	Lb. a.i./A azoxystrobin	Treated acres per gallon product
7	0.056	0.034	18.3
14	0.113	0.070	9.1
20.5	0.167	0.100	6.2
27.5	0.224	0.134	4.6
34.5	0.280	0.169	3.7

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [Bulk/Mini-Bulk]

Refillable container. Refill this container only with AzoxyProp Fungicide. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Orbit®, Quadris®, Tilt® and the Syngenta logo are trademarks of a Syngenta Group Company

Topsin® trademark of Pennwalt Corporation

©201X Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1178A

43
45

BASE LABEL

AzoxyProp

GROUP 3 11 FUNGICIDES

Fungicide

Broad spectrum fungicide for control of plant diseases

Active Ingredient:

Azoxystrobin	7.0%
Propiconazole	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1178

EPA Est.

SCP 1178A

_____ gallons
Net Contents

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling.

Environmental Hazards

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Do not apply directly to water except as specified on this label. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify state and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

45
45

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

The Syngenta logo is a trademark of a Syngenta Group Company

©201X Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1178A

AzoxyProp Notif - bb - 1-27-11

000100-01178.20110127.QUILT-NOTIF-0111.pdf