



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 14, 2018

Nakia Smith,
Fungicide Registration Manager
Syngenta Crop Protection, LLC
PO Box 18300
410 Swing Road
Greensboro, NC 27419

Subject: Label Amendment – Rate Clarifications
Product Name: Quilt Fungicide
EPA Registration Number: 100-1178
Application Date: January 25, 2018
Decision Number: 538066

Dear Ms. Smith,

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 100-1178
Decision No. 538066

with FIFRA section 6. If you have any questions, please contact Eleanor Thornton by phone at 703-305-6799, or via email at Thornton.eleanor@epa.gov.



Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

ACCEPTED

06/14/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 100-1178

PROPICONAZOLE	GROUP	3	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE

Quilt® Fungicide

Broad spectrum fungicide for control of plant diseases

Active Ingredient:

Azoxystrobin 7.0%

Propiconazole 11.7%

Other Ingredients: 81.3%

Total: 100.0%

Quilt Fungicide is formulated as a suspo-emulsion (SE) and contains 1.04 lb ai of propiconazole and 0.62 lb ai of azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet. See First Aid statement inside booklet or on container label.

EPA Reg. No. 100-1178

EPA Est.

_____ gallons
Net Contents

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Wear appropriate protective eyewear (goggles, face shield, or safety glasses)
- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes and socks

User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Propiconazole is toxic to shrimp.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Protective eyewear (goggles, face shield, or safety glasses)
- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

PRODUCT INFORMATION

Quilt Fungicide is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. Quilt Fungicide is a member of Syngenta's Plant Performance™ product line and may also improve the yield and/or quality of the crop. These additional benefits are due to positive effects on plant physiology. The effects may vary according to other factors such as crop, crop hybrid, or environment. Quilt may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications should be made according to the use directions that follow.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult www.epa.gov/espp/bulletins.htm. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

USE RESTRICTIONS

- Do not use in nurseries, greenhouses or landscape plantings.
- Do not apply in a manner that will result in exposure to humans or animals.

APPLICATION INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Quilt Fungicide has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

Rotational Crops: Soybeans may be planted as a double crop following a cereal crop which has been treated with Quilt Fungicide. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding. Plant wild rice 45 days after Quilt Fungicide application. To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Quilt Fungicide application to the preceding crop, unless the second crop appears on this label.

Rotational Crops	Planting Time From Last Quilt Fungicide Application
Beans, Dry and Succulent Brassica, leafy greens (subgroup 5B) Bulb Vegetables Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale, oat and rye) Corn (field, seed, pop, and sweet) Dill Grasses grown for seed (Cool season grasses only) Mint Peanuts Quinoa Radish Rice (including Wild Rice) Sorghum Soybeans Strawberry and other Low-Growing Berry Subgroup 13-07G Sugar beets Sugarcane Watercress	0 days
Buckwheat Millet	12 Months
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/year)	75 days
All Other Crops Intended for Food, Grazing, or Feed	105 days

Any crop listed and approved for use on this label has a 0 day plant back interval.

Crop Tolerance/Phytotoxicity: Quilt Fungicide demonstrates some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone can contribute to phytotoxicity. Under certain environmental conditions, tank mixes of Quilt Fungicide plus herbicides and/or fertilizers may cause crop injury in barley, triticale and wheat.

ATTENTION

Quilt Fungicide is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray Quilt Fungicide where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply Quilt Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Spray Drift Management: A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 10 mph.

Droplet Size

Apply as a medium or coarser spray (ASAE Standard 572)

Temperature Inversions

Do not apply at wind speeds below 3 mph. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of propiconazole. Where states have more stringent regulations, they must be observed.

Equipment

All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Integrated Pest Management: Quilt Fungicide should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The SPECIFIC USE DIRECTIONS section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area. Quilt Fungicide may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

PROPICONAZOLE	GROUP	3	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE

Quilt Fungicide is a mixture of Group 3 (propiconazole) and Group 11 (azoxystrobin) fungicides. Quilt Fungicide has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (Qol) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Syngenta encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the specific crop recommendations that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups. In situations requiring multiple sprays, develop season long spray programs for Group 11 (Qol) fungicides. The program should meet the goal of no more than $\frac{1}{3}$ of the total sprays per season, when a Group 11 fungicide is used as a solo product, or $\frac{1}{2}$ the total sprays when a Group 11 fungicide is used in a mixture. Programs that include both solo Group 11 products and/or mixes containing Group 11 products should be no more than $\frac{1}{2}$ the total sprays.

Quilt Fungicide should not be alternated or tank mixed with any fungicide to which resistance has already developed.

SPRAY EQUIPMENT

Quilt Fungicide may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be *16-mesh or coarser*.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - (1) Maintain 35-40 psi at nozzles.
 - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

MIXING INSTRUCTIONS

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Quilt Fungicide Alone (no tank mix):

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add Quilt Fungicide to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Quilt Fungicide has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Quilt Fungicide + Tank Mixtures: Quilt Fungicide is usually compatible with all tank mix partners listed on this label. Do not combine Quilt Fungicide in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the

combination is physically compatible, effective, and noninjurious to the crop under your conditions of use. To determine the physical compatibility of Quilt Fungicide with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water dispersible granular products first, then liquid flowables (suspoemulsions), and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank mix partner(s) into the tank in the same order as described above in the "Quilt Fungicide + Tank Mixtures" section.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the Quilt Fungicide to the spray tank.
- Allow Quilt Fungicide to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.

RESTRICTIONS:

- No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product may not be mixed with any product which prohibits such mixing.

APPLICATION PROCEDURES

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.

Ground Application:

- For field crops (non-trees), apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- For field crops (non-trees), apply in a minimum spray volume of 2 gallons per acre unless specified otherwise.
- For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- Thorough coverage is necessary to provide good disease control.
- Quilt Fungicide is extremely phytotoxic to certain apple varieties.

- AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).
- DO NOT spray Quilt Fungicide where spray drift may reach apple trees.
- DO NOT apply via ULV.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches per acre of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Note: Do not inject Quilt Fungicide at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part Quilt Fungicide. Quilt Fungicide is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton®, Buna-N, Neoprene, or PVC seals.

APPLICATION INSTRUCTIONS

Apply Quilt Fungicide at rates and timings as described in this label.

Directions for Use Through Sprinkler and Drip Chemigation Systems (Apply Quilt Fungicide only to crops for which chemigation is specified on this label):

Spray Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Use Precautions for Sprinkler and Drip Irrigation Applications

Sprinkler Irrigation: Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, drip (trickle), or hand move irrigation systems. Do not apply this product through any other type of irrigation system except as specified on this label.

Apply with center pivot or continuous-move equipment distributing ½ acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Do not apply when wind speed favors drift beyond the area intended for treatment. Plant injury, lack of

effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank

prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Crop Use Directions

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Almonds	Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	14 - 27.5	Apply Quilt Fungicide at early bloom stage. If disease pressure is low, a second application of 14 fl oz/A may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50-75% bloom and petal fall. Quilt Fungicide may only be used on only two blossom blight applications. Tilt may be used alone for one of these applications.
	Anthracnose (<i>Collectotrichum acutatum</i>)	14 – 27.5	Apply Quilt Fungicide beginning at bud break on a 7- to 14-day interval. Make no more than two consecutive applications before switching to another fungicide with a different mode of action.

Note: Make no more than two sequential applications of a Group 11 fungicide prior to alternating with two applications of another product with a different mode of action than Group 11 fungicides. Almond diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. Quilt Fungicide may be applied by ground or by air (minimum of 15 gal/A). Aerial application may be used if necessary but disease control may be reduced. Quilt Fungicide may be applied by air only at growth stages prior to and including 5 weeks after petal fall.

Specific Use Restrictions:

- 1) Do not apply more than 110 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.9 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (27.5 fl oz/year) or 7 applications per year at the low rate (14 fl oz product/A).
- 6) Do not apply within 60 days of harvest (60-day PHI).
- 7) Do not graze livestock in treated areas or cut treated cover crop for feed.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Bananas Plantains	Black Sigatoka (<i>Mycosphaerella fijiensis</i>) Yellow Sigatoka (<i>Mycosphaerella musicola</i>)	10.5	Apply Quilt Fungicide before disease symptoms appear at the onset of the rainy season. Apply 10.5 fl oz of Quilt Fungicide per acre in 10-20 gallons of water. Apply no more than 2 consecutive applications on a 21- to 25-day schedule before rotating to another labeled product with a different mode of action for at least 2 sprays. A maximum of 8 applications can be made. If possible, it is recommended to have at least 2 consecutive months 'triazole free' during the period of lower disease pressure.

Note: Quilt Fungicide may be applied by ground (minimum of 15 gal/A) or aerial application (minimum of 5 gal/A).

Specific Use Restrictions:

- 1) Do not apply more than 84 fl oz of Quilt Fungicide during each growing season (this includes any preharvest sprays).
- 2) Do not apply more than 10.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.67 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 1.08 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 8 applications per year.
- 6) Do not apply Quilt Fungicide within 100 yards of non-bagged bananas.
- 7) Do not apply Quilt Fungicide on bananas unless they are protected by polyethylene bags.
- 8) Do not apply Quilt Fungicide on plantains if the fruit present are not protected with polyethylene bags.
- 9) Do not feed whole bananas and plantains to animals.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
<p>Beans, Dry and Succulent</p> <p>Bean (<i>Cicer arietinum</i>) (<i>Lupinus</i> spp.) (<i>Phaseolus</i> spp.) (<i>Vigna</i> spp.) (<i>Vicia faba</i>)</p> <p>See below for complete list of dry and succulent beans.</p>	<p>Alternaria Blight (<i>Alternaria</i> spp.) Alternaria Leaf Spot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum lindemuthianum</i>) Ascochyta Blight (<i>Mycosphaerella pinodes</i>) Ascochyta Leaf and Pod Spot (<i>Ascochyta</i> spp.) Ascochyta Leaf Spot (<i>Ascochyta phaseolorum</i>) Bean Rust (<i>Uromyces appendiculatus</i>) Rust (<i>Phakopsora</i> spp.) Web Blight (<i>Rhizoctonia solani</i>)</p>	<p>14</p>	<p>Apply when conditions are conducive for disease. Up to three applications may be made on a 7- to 14-day interval.</p> <p>NOTE: On certain bean varieties, Quilt Fungicide application may cause crinkled and/or greener leaves. Yields of beans displaying these characteristics have not been reduced.</p>

Dry and Succulent Beans: *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and wax bean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, blackeyed pea and cowpea)

Note: Quilt Fungicide may be applied by ground or by air.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz of Quilt Fungicide/A/year.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.34 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 3 applications per year.
- 6) Do not apply within 7 days of harvest (7-day PHI) for succulent beans.
- 7) Do not apply within 14 days of harvest (14-day PHI) for dry beans.
- 8) Not for use on cowpea cultivars intended for livestock feeding only.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
<p>Berry, Bushberry Subgroup 13-07B</p> <p>Blueberry, Highbush Blueberry, Lowbush Cranberry, Highbush Currant, Black Currant, Red Elderberry Gooseberry</p> <p>And cultivars and/or hybrids of these.</p> <p>See below for complete list of bushberry subgroup.</p>	<p>Mummyberry (<i>Monilinia vaccinicornymbosi</i>)</p>	<p>14 - 20.5</p>	<p>Make first application of Quilt Fungicide beginning at green tip and repeat in 7-10 days. If conditions are favorable for disease development, additional application may need to be made at pink bud and repeating every 7 to 10 days through petal fall. Do not apply more than two consecutive applications before alternating to a non-Group 11 containing fungicide.</p>
	<p>Botryosphaeria Canker (<i>Botryosphaeria spp.</i>) Leaf Spot (<i>Septoria spp.</i>) Leaf Spot and Stem Canker (<i>Septoria albopunctata</i>) Phomopsis Twig Blight, Fruit Rot, and Stem Canker (<i>P. vaccinii</i>) Powdery Mildew (<i>Microsphaera vacinii</i>)</p>	<p>14 - 20.5</p>	<p>Quilt Fungicide applications should begin prior to disease development and continue throughout the season on a 14-day interval. Make no more than two consecutive sprays before alternating to a non-Group 11 fungicide. Make no more than 3 applications of a Group 11 fungicide per acre per year.</p>

Bushberry Subgroup: Aronia berry; Blueberry, highbush; Blueberry, lowbush; Buffalo currant; Chilean guava; Cranberry, highbush; Currant, black; Currant, red; Elderberry; European barberry; Gooseberry; Honeysuckle, edible; Huckleberry; Jostaberry; Juneberry (Saskatoon berry); Lingonberry; Native currant; Salal; Sea buckthorn

Note: Quilt Fungicide may be applied by ground (minimum of 15 gal/A) or by air.

Specific Use Restrictions:

- 1) Do not apply more than 61.5 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 20.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.84 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 0.75 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 3 applications per year.
- 6) Do not apply within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Berry, Caneberry Subgroup 13- 07A Blackberry Bingleberry Boysenberry Dewberry Loganberry Lowberry Marionberry Olallieberry Raspberry, Red and Black Wild Raspberry Youngberry And cultivars and/or hybrids of these.	Leaf and Cane Spot <i>(Septoria rubi)</i> Leaf Spot <i>(Septoria spp.)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i> Rust <i>(Pucciniastrum vaccinii)</i>	14 - 20.5	Quilt Fungicide applications should begin prior to disease development and continue throughout the season on a 7- to 14-day interval. Make no more than two consecutive sprays before alternating to a non-Group 11 fungicide. Make no more than 3 applications of a Group 11 fungicide per acre per year.

Note: Quilt Fungicide may be applied by ground (minimum of 15 gal/A) or by air.

Specific Use Restrictions:

- 1) Do not apply more than 61.5 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 20.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.84 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 3 applications per year.
- 6) Do not apply within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Brassica, leafy greens (subgroup 5B) Cabbage, Chinese (bok choy) Collards Kale Mustard Greens (See additional crops below)	Alternaria Leaf Spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Black Spot <i>(Alternaria spp.)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Powdery Mildew <i>(Erysiphe polygoni)</i> Ring Spot <i>(Mycosphaerella brassicicola)</i>	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. Apply no more than two applications before switching to another fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Leafy Brassica (subgroup 5B): Broccoli raab; Cabbage, Chinese (bok choy); Collards, Kale, Mizuna, Mustard greens; Mustard spinach; Rape greens

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.34 lb ai propiconazole-containing product/A/year.
- 4) Do not apply more than 0.75 lb ai azoxystrobin-containing product/A/year.
- 5) Do not exceed 3 applications per year when applying at the highest rate (14 fl oz/A) or 4 applications at the lowest rate (10.5 fl oz/A).
- 6) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
<p>Bulb Vegetables</p> <p>Dry Bulb Onions</p> <p>Subgroup 3-07A</p> <p>Garlic Onions (dry bulb) Shallots (dry bulb)</p> <p>Green Onions</p> <p>Subgroup 3-07B</p> <p>Leeks Onions (green) Shallots (green)</p> <p>Including all cultivars and/or hybrids of these</p> <p>See below for complete list of bulb vegetables.</p>	<p>Cladosporium Leaf Blotch (<i>C. allii</i>)</p> <p>Purple Blotch (<i>Alternaria porri</i>)</p> <p>Rust (<i>Puccinia allii</i>)</p> <p>Suppression of Botrytis</p> <p>Leaf Blight (<i>Botrytis squamosa</i>)</p>	<p>14 – 27.5</p>	<p>Begin applications when conditions favor disease development and continue on a 7- to 10-day interval. Use the higher rate and shorter interval when disease conditions are severe. Make only 1 application before alternating to a non-Group 11 fungicide.</p>

Complete List of Bulb Vegetables: Chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritillaria, bulb; fritillaria, leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

Note: Quilt Fungicide may be applied by ground (15 gal/A minimum) or aerial application (minimum of 5 gal/A). Mixing with products formulated as an emulsifiable concentrate (EC) may result in phytotoxicity.

Specific Use Restrictions:

- 1) Do not apply more than 55.3 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 2 applications per year when using the highest rate (27.5 fl oz/year) or 3 applications per year at the low rate (14 fl oz product/A).
- 6) Do not apply within 14 days of harvest (14-day PHI) on dry bulb onions.
- 7) Quilt Fungicide may be applied the day of harvest (0-day PHI) for green onion types.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Carrots	Early Blight <i>(Cercospora carotae)</i> Powdery Mildew <i>(Erysiphe polygoni)</i> Suppression of Late Blight <i>(Alternaria dauci)</i>	14	Apply 14 fl oz of Quilt Fungicide when conditions favor disease development. Continue applications on a 7- to 10-day interval, using the shorter interval when disease conditions are severe. Make no more than one application before alternating to a non-Group 11 fungicides.

Note: Quilt Fungicide may be applied by ground (minimum of 15 gal/A), or aerial application (minimum of 5 gal/A), or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 2.0 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 3 applications per year.
- 6) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Celery	Early Blight (<i>Cercospora apii</i>) Late Blight (<i>Septoria apicola</i>)	14	Apply Quilt Fungicide on a 7- to 10-day schedule in alternation with Tilt® alone or another product with a different mode of action than Group 11 fungicides.

Note: Quilt Fungicide may be applied by ground, air (5 gal/A minimum) or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year.
- 6) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Cereals Wheat See next section for other cereals.	Early season suppression of: Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blight <i>(Septoria tritici)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	7 – 14	Apply Quilt Fungicide in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control. If applying early in the season in mixture with fertilizers and herbicides you may see flecking and burning.
	Control of Leaf Diseases: Glume Blotch <i>(Stagonospora nodorum)</i> Helminthosporium Leaf Blight <i>(Drechslera tritici-repentis)</i> Kernel Blight <i>(Alternaria spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Net Blotch <i>(Pyrenophora teres)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Rust <i>(Puccinia spp.)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt Fungicide is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl oz/A may be applied. Applications may be made no closer than a 14-day interval. Quilt Fungicide can be applied through full head emergence (Feekes growth stage 10.5.4). Do not apply after this stage.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of Quilt Fungicide plus half the rate listed of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 28.0 fl oz/A/year of Quilt Fungicide.
 - 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
 - 3) Do not apply more than 0.22 lb ai propiconazole-containing products/A/year.
 - 4) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/year.
 - 5) Do not apply more than 2 applications/A/year when using the highest rate (14 fl oz/A) or 4 applications when using the lowest rate (7 fl oz/A).
 - 6) Do not harvest within 7 days (7-day PHI) for forage or hay.
 - 7) Quilt Fungicide can be applied through full head emergence (Feekes growth stage 10.5.4). Do not apply after this stage.
 - 8) Under certain environmental conditions, tank mixes of Quilt Fungicide plus herbicides and/or fertilizers may cause crop injury.
-

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Cereals Barley Oats Rye Triticale	Early season suppression of: Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blight <i>(Septoria tritici)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	7 – 14	Apply Quilt Fungicide in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control. If applying early in the season in mixture with fertilizers and herbicides you may see flecking and burning.
	Control of Leaf Diseases: Barley Scald <i>(Rhynchosporium secalis)</i> Barley Stripe <i>(Pyrenophora graminea)</i> Glume Blotch <i>(Stagonospora nodorum)</i> Helminthosporium Leaf Blight <i>(Drechslera tritici-repentis)</i> Kernel Blight <i>(Alternaria spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Net Blotch <i>(Pyrenophora teres)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Rust <i>(Puccinia spp.)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt Fungicide is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl oz/A may be applied. Apply no closer than a 14-day interval.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of Quilt Fungicide plus half the rate specified on other EPA registered fungicides such as Topsin M. Apply at tillering but before elongation has occurred.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. An adjuvant may be added at recommended rates to improve canopy coverage and penetration while reducing evaporation and drift. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 28.0 fl oz/A/year of Quilt Fungicide.
 - 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
 - 3) Do not apply more than 0.22 lb ai propiconazole-containing products/A/year.
 - 4) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/year.
 - 5) Do not apply more than 2 applications/A/year when using the highest rate (14 fl oz/A) or 4 applications when using the lowest rate (7 fl oz/A).
 - 6) Do not apply within 7 days of harvest (7-day PHI) for forage and hay.
 - 7) Quilt Fungicide can be applied through full head emergence (Feekes growth stage 10.5.4). Do not apply after this stage.
 - 8) Under certain environmental conditions, tank mixes of Quilt Fungicide plus herbicides and/or fertilizers may cause crop injury.
-

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
<p>Citrus Fruit (non-bearing)</p> <p>Calamondin Citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma mandarin Tangerine</p> <p>Including all cultivars and/or hybrids of these</p>	<p>Greasy Spot (<i>Mycosphaerella citri</i>)</p>	<p>20.5 - 27.5</p>	<p>Apply Quilt Fungicide up to 4 times. The other sprays may be Tilt® alone or another product with a different mode of action than Group 11 fungicides.</p> <p>Quilt Fungicide may be applied by ground, air or chemigation.</p> <p>Do not use Quilt Fungicide in citrus plant propagation nurseries.</p>

Specific Use Restrictions:

- 1) Do not apply more than 82.5 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.68 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 3 applications per year when using the highest rate (27.5 fl oz/A) or 4 applications when using the lowest rate (20.5 fl oz/A).
- 6) Do not apply to citrus that will bear harvestable fruit within 12 months.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Corn Field Pop (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium Leaf Blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)	7 - 14	For leaf blights apply Quilt Fungicide when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate. Apply no more than 2 applications of Quilt Fungicide or any other Group 11 fungicide per year. Do not use adjuvants or other additives after the V8 growth stage and prior to the VT growth stage, as use during these development times may impose stress on the plant that could inhibit proper kernel development. VT is defined as when the last branch of the tassel is completely visible but silks have not yet emerged from the ear shoot.
	Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Gray Leaf Spot (<i>Cercospora zeae- maydis</i>) Rusts (<i>Puccinia spp.</i>)	10.5 - 14	Apply Quilt Fungicide when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate. Apply no more than 2 applications of Quilt Fungicide or any other Group 11 fungicide per year.

Note: Quilt Fungicide may be applied by ground, air, or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 28 fl oz (0.224 lb ai propiconazole) for field corn harvested for forage.
- 3) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 4) Do not apply more than 0.45 lb ai propiconazole-containing products/A/year.
- 5) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/year.
- 6) Do not exceed 2 applications per year.
- 7) Do not apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Corn Sweet corn (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium Leaf Blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)	7 - 14	For leaf blights apply Quilt Fungicide when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. Alternate applications of Quilt Fungicide with Tilt or another product with a different mode of action than Group 11 fungicides.
	Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Gray Leaf Spot (<i>Cercospora zeae- maydis</i>) Rusts (<i>Puccinia spp.</i>)	10.5 - 14	Apply Quilt Fungicide when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Alternate applications of Quilt Fungicide with Tilt or another product with a different mode of action than Group 11 fungicides.

Note: Quilt Fungicide may be applied by ground, air, or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (14 fl oz/A) or 8 applications when using the lowest rate (7 fl oz/A).
- 6) Do not apply to sweet corn within 14 days of harvest (14-day PHI) for ears or forage.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Cranberries	Cottonball (<i>Monilinia oxycocci</i>)	14 – 20.5	<p>Make the first application at leaf bud break and repeat in 14 days. A third application should be made at early bloom and again in 14 days. Make no more than 2 consecutive sprays before alternating to a non-Group 11 fungicide.</p> <p>For resistance management, make no more than 3 sprays per season using any Group 11 fungicide. Under severe pressure, use the higher rate for control.</p>

Note: Quilt Fungicide may be applied by ground (minimum of 10 gal/A) or aerial application (minimum of 20 gal/A).

Specific Use Restrictions:

- 1) **For use only in Oregon, Washington, and Wisconsin.**
- 2) Do not apply more than 61.5 fl oz/A/year of Quilt Fungicide.
- 3) Do not apply more than 20.5 fl oz/A/application of Quilt Fungicide.
- 4) Do not apply more than 0.67 lb ai of propiconazole-containing products/A/year.
- 5) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 6) Do not exceed 3 applications per year.
- 7) Do not apply within 45 days of harvest (45-day PHI).
- 8) Do not allow release of irrigation of flood water to non-target aquatic habitat for at least 14 days after the last application.
- 9) Do not treat cranberry fields used for aquaculture of fish or crustacean.
- 10) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators should use care in making applications near non-target aquatic habitats.
- 11) Do not apply to flooded crop.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Dill	Cercospora Leaf Blight (<i>Cercosporidium punctum</i>) Powdery Mildew (<i>Erysiphe heraclei</i>)	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.34 lb ai propiconazole-containing product/A/year.
- 4) Do not apply more than 1.5 lb ai azoxystrobin-containing product/A/year.
- 5) Do not exceed 3 applications per year when using the highest rate (14 fl oz/A) or 4 applications at the lowest rate (10.5 fl oz/A).
- 6) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Filberts	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	14 – 27.5	<p>Begin applications when green leaf tissue becomes visible and continue on a 2- to 3-week interval. Under severe disease conditions, use the higher rate and shorter interval.</p> <p>NOTE: On certain varieties, Quilt Fungicide applications may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to Quilt Fungicide treatments.</p>

Note: Quilt Fungicide may be applied by ground or aerial application (minimum of 15 gal/A).

Specific Use Restrictions:

- 1) Do not apply more than 110 fl oz/A of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.9 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.2 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (27.5 fl oz/A) or 7 applications when using the lowest rate (14 fl oz/A).
- 6) Do not apply within 60 days of harvest (60-day PHI).
- 7) Do not graze livestock in treated areas or cut treated cover crop for feed.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Grasses Grown for seed (Cool season grasses only)	Ergot Stem Diseases Powdery Mildew <i>(Erysiphe graminis)</i> Rusts <i>(Puccinia spp.)</i>	14 - 27.5	<p>Apply Quilt Fungicide when powdery mildew infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 27.5 fl oz/A (except bluegrass, apply 14 fl oz/A) and make applications at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin application early in the growing season.</p> <p>Make no more than 2 sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.</p>

Note: Apply Quilt Fungicide in a minimum of 20 gal of water per acre for ground application, or in a minimum of 10 gal of water per acre for aerial application. Quilt Fungicide may also be applied by chemigation.

Specific Use Restrictions:

- 1) **For use only in Idaho, Minnesota, Nebraska, Oregon, and Washington.**
- 2) Do not apply more than 110 fl oz/A/year of Quilt Fungicide.
- 3) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 4) Do not apply more than 0.90 lb ai propiconazole-containing products/A/year.
- 5) Do not apply more than 0.75 lb ai azoxystrobin-containing products/A/year.
- 6) Do not exceed 4 applications per year when using the highest rate (27.5 fl oz/A) or 7 applications when using the lowest rate (14 fl oz/A).
- 7) Do not apply within 20 days of harvest (20-day PHI) of seed.
- 8) Do not feed hay cut within 20 days of the last application.
- 9) Do not graze treated areas within 140 days of the last application.
- 10) Do not apply to bermudagrass grown for seed.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Mint Peppermint Spearmint	Powdery Mildew (<i>Erysiphe</i> spp.) Rust (<i>Puccinia menthae</i>)	14	Begin applications when the plants are 2 - 4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

Note: Quilt Fungicide may be applied by ground (minimum of 20 gal/A), air (minimum of 5 gal/A), or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A /year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.22 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 0.75 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 2 applications per year.
- 6) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Peanuts	Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Rust (<i>Puccinia arachidis</i>) Web Blotch (<i>Phoma arachidicola</i>)	10.5 - 14	Apply Quilt Fungicide beginning 35-40 days after planting or at the first appearance of disease. Continue applications on a 14-day schedule. Under heavy disease pressure use higher listed application rates. Quilt Fungicide also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt Fungicide may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 0.80 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (14 fl oz/A) or 5 applications when using the lowest rate (10.5 fl oz/A).
- 6) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Pecans	Anthracnose (<i>Glomerella cingulata</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae</i> pv <i>pecanae</i>) Pecan Scab (<i>Cladosporium caryigenum</i>) Powdery Mildew (<i>Microsphaera penicillata</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>)	14 – 27.5	Pecan scab: Apply 14 – 27.5 fl oz/A Quilt Fungicide on a 14-day schedule during bud break and pre-pollination sprays. Apply 20 – 28 fl oz/A during nut formation and cover sprays. Use higher rates when disease pressure is heavier. Do not apply after shuck split. Other foliar diseases: Quilt Fungicide may be applied for control of mid to late season foliar diseases at 14 – 20.5 fl oz/A with other pecan products labeled for these diseases. Observe all directions, precautions, and limitations for the other products. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides. Do not apply after shuck split.

Note: Ground applications should be applied in sufficient water to provide for full coverage. Quilt Fungicide may be applied by ground or air (minimum of 20 gal/A).

Specific Use Restrictions:

- 1) Do not apply more than 122 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.9 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 1.2 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (27.5 fl oz/A) or 8 applications when using the lowest rate (14 fl oz/A).
- 6) Do not graze livestock in treated areas or cut treated cover crops for feed.
- 7) Do not apply after shuck split or within 45 days of harvest (45-day PHI), whichever is first.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult www.epa.gov/espp/bulletins.htm. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Pistachios	Botryosphaeria Panicle and Shoot Blight (<i>B. dothidea</i>)	14 – 27.5	Begin applications when green leaf tissue becomes visible and continue on a 14- to 21-day interval. Under severe disease conditions, use the higher rate and the shorter interval. Make no more than 2 consecutive applications of Quilt Fungicide before switching to another non-Group 11 fungicide.

Note: Quilt Fungicide may be applied by ground or aerial application (15 gal/A minimum).

Specific Use Restrictions:

- 1) Do not apply more than 110 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.9 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (27.5 fl oz/A) or 7 applications when using the lowest rate (14 fl oz/A).
- 6) Do not graze livestock in treated areas or cut treated cover crop for feed.
- 7) Do not apply within 60 days of harvest (60-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Quinoa	Leaf Spot (<i>Ascochyta hyalospora</i>) Stalk Rot (<i>Phoma exigua</i>)	10.5 – 14	Apply prior to disease development. An adjuvant may be added at recommended rates.

Note: Quilt Fungicide can be applied by ground, chemigation, or aerial application.

Specific Use Restrictions:

- 1) Do not apply more than 28.0 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.22 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/year.
- 5) Do not apply more than 2 applications/A/year.
- 6) Under certain environmental conditions, tank mixes of Quilt Fungicide plus herbicides and/or fertilizers may cause crop injury.
- 7) Do not apply within 7 days of harvest (7-day PHI) for forage and hay.
- 8) Do not apply within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Radish	Cercospora Leaf Spot (<i>Cercospora</i> spp.)	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7- to 10-day interval. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing product/A/year.
- 4) Do not apply more than 2.0 lb ai azoxystrobin-containing product/A/year.
- 5) Do not exceed 4 applications per year when using the highest rate (14 fl oz/A) or 5 applications per year at the lowest rate (10.5 fl oz/A).
- 6) Do not apply within 14 days of harvest (14-day PHI)

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Rice	Aggregate Sheath Spot <i>(Rhizoctonia oryzae-sativa)</i> Black Sheath Rot <i>(Gaeumannomyces graminis)</i> Brown Leaf Spot <i>(Helminthosporium oryzae)</i> Kernel Smut <i>(Tilletia barclayana)</i> Leaf Blast <i>(Pyricularia grisea)</i> Leaf Smut <i>(Entyloma oryzae)</i> Narrow Brown Leaf Spot <i>(Cercospora oryzae)</i> Sheath Blight <i>(Rhizoctonia solani)</i> Sheath Spot <i>(Rhizoctonia oryzae)</i> Stem Rot <i>(Sclerotium oryzae)</i> For disease suppression of: False Smut <i>(Ustilaginoidea virens)</i>	14 - 34.5	Timing of Quilt Fungicide application will depend on disease severity, disease complex and rice variety/growth stage. Consult local extension experts for local economic thresholds established for various rice varieties and diseases. Quilt Fungicide should be applied at 28 - 34.5 fl oz/A prior to or in the early stages of disease development. If disease reappears, use another fungicide for the second application. For leaf blast control, use a minimum of 34.5 fl oz/A. For neck or panicle blast, additional applications of Quadris® may be made (refer to Quadris label). Tank mix option: Apply 14 - 20.5 fl oz/A of Quilt Fungicide in a tank mix with Quadris or other fungicides for control of rice diseases. A minimum of 0.15 lb ai/A azoxystrobin should be in the tank mix (see Quilt Fungicide rate conversion table at the end of the label). Make no more than 2 applications of a Group 11 fungicide per year.

Note: Applications may be made by air only. For aerial application, volumes should be 5 - 10 GPA. An adjuvant may be added at specified rates to improve canopy coverage and penetration while reducing evaporation and drift.

Specific Use Restrictions:

- 1) Do not apply more than 41 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 34.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.34 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 0.70 lb ai azoxystrobin-containing products/A/year.
- 5) Do not make more than 1 application per year at the highest rate (34.5 fl oz/A) or 2 applications per year at the lowest rate (14 fl oz/A).
- 6) Do not apply within 35 days of harvest (35-day PHI).
- 7) Do not release floodwater within 14 days of an application.
- 8) Do not apply to stubble or ratoon crop rice.
- 9) Do not use in rice fields where commercial farming of crayfish will be practiced.
- 10) Do not drain water from treated rice fields into ponds used for commercial fish farming.
- 11) Do not use water drained from treated fields to irrigate other crops.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult www.epa.gov/espp/bulletins.htm. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Sorghum	Anthracnose <i>(Colletotrichum graminicola)</i> Ergot <i>(Claviceps sorghi)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i>	14	<p>For foliar diseases, Quilt Fungicide applications should begin prior to disease development. Applications may be made by ground, air or chemigation.</p> <p>For ergot: Make the first application at or just prior to flowering. Repeat on a 5- to 7-day interval. Apply up to 4 times.</p> <p>Resistance Management: Do not apply more than two sequential applications of Quilt Fungicide or other Group 11 containing fungicides before alternation with a fungicide that is not in Group 11.</p>

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt Fungicide may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz of Quilt Fungicide/A/year.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.22 lb propiconazole ai/A (27.5 fl oz Quilt Fungicide) on sorghum harvested for forage.
- 4) Do not apply more than 0.45 lb ai propiconazole-containing product/A/year.
- 5) For grain and stover, do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 6) For forage, do not apply more than 0.5 lb ai/A/year of azoxystrobin-containing products.
- 7) Do not exceed 4 applications per year.
- 8) Do not graze livestock or cut for green chop or silage within 30 days of application.
- 9) Do not apply within 30 days of harvest (30-day PHI) for forage
- 10) Do not apply within 21 days of harvest (21-day PHI) for grain and stover.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Soybeans	Aerial Web Blight (<i>Rhizoctonia solani</i>) Alternaria Leaf Spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Blight and Leaf Spot (<i>C. kickuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe</i> spp.) Soybean Rust (<i>Phakopsora pachyrhizi</i>)	14 – 20.5	<p>Aerial Web Blight: Apply 14 - 20.5 fl oz/A at the first appearance of disease and repeat the application 14 - 21 days later. Under severe disease conditions use the higher rate and shorter interval.</p> <p>Other foliar diseases (except rust): Apply 20.5 fl oz/A at growth stage R3 (early pod set when pods are 1/8-1/4 inch long) and 14 - 21 days later at growth stage R5 (pod fill).</p> <p>Soybean Rust: Apply 14 - 20.5 fl oz/A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter interval when disease is present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p> <p>On certain varieties, Quilt Fungicide applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to Quilt Fungicide treatments.</p>

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions will not provide good coverage. Quilt Fungicide may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 20.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.34 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 2 applications per year when using the highest rate (20.5 fl oz/A) or 3 applications when using the lowest rate (14 fl oz/A).
- 6) Apply up to Stage R6.

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Stone Fruits Crop Group 12-12 Apricot Cherry, sweet Cherry, tart Nectarine Peach Plum Plumcot Prune	Brown Rot Blossom Blight <i>(Monilinia spp.)</i>	14	Apply Quilt Fungicide at early bloom stage. If disease pressure is low, a second application of 14 fl oz/A may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50-75% bloom and petal fall. Quilt Fungicide may only be used on 2 of these applications. Tilt may be used alone for one of these applications.
Including all cultivars and hybrids of these	Brown Rot on Fruit <i>(Monilinia spp.)</i>	14	Apply as needed, a maximum of 2 sprays of Quilt Fungicide, during the preharvest period up to the day of harvest. Make the two applications no closer than 10 days apart.
	Cherry Leaf Spot <i>(Blumeriella jaapii)</i> Powdery Mildew <i>(Podosphaera clandestine,</i> <i>Sphaerotheca pannosa)</i>	14	Follow the blossom blight schedule. Make up to 2 additional applications on a 10- to 14-day interval from the end of petal fall to harvest.

Note: Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with 2 applications of another product with a different mode of action than Group 11 fungicides. Stone fruit diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. Aerial application (minimum of 15 gal/A) may be used if necessary but disease control may be reduced.

Applications of Quilt Fungicide during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply Quilt Fungicide to Stanley plums earlier than 21 days prior to harvest.

Specific Use Restrictions:

- 1) Do not apply more than 70 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.56 lb ai propiconazole-containing products/A/year.
- 4) Do not apply more than 1.0 lb ai azoxystrobin-containing products/A/year.
- 5) Do not exceed 5 applications per year.
- 6) Quilt Fungicide may be applied the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
<p>Tree Nuts Crop Group 14-12</p> <p>See list below for tree nuts labeled</p> <p>See specific directions for Almond Filbert Pecan Pistachios</p>	<p>Foliar Diseases</p>	<p>14 - 27.5</p>	<p>Apply Quilt Fungicide at first sign of disease. Repeat on a 7- to 14-day interval. Do not make more than two consecutive applications before switching to a non-Group 11 fungicide. Make no more than 4 applications per year of a Quilt Fungicide or other Group 11 fungicide containing product.</p>

Additional tree nuts: Almond (see specific directions), Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (see specific directions), Hickory, Macadamia, Pecan (see specific directions), Pistachios (see specific directions), Walnut

Note: For best control of tree nut diseases, ground applications are recommended. Quilt Fungicide may be applied by ground or aerial application (15 gal/A minimum).

Specific Use Restrictions:

- 1) Do not apply more than 110 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 27.5 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.9 lb ai of propiconazole-containing products/A/year.
- 4) Do not apply more than 1.2 lb ai of azoxystrobin-containing products/A/year.
- 5) Do not exceed 4 applications per year.
- 6) Do not graze livestock in treated areas or cut treated cover crop for feed.
- 7) Do not apply within 60 days of harvest (60-day PHI) except for pecan (see specific use directions).

Crop	Target Diseases	Use Rate fl oz product/A	Application Instructions
Watercress	Alternaria Leaf Spot (<i>Alternaria</i> spp.) Cercospora Leaf Spot (<i>C. nasturtii</i>)	10.5 - 14	Begin applications at first sign of disease. Repeat on a 7-10 day interval. Make no more than 2 applications before harvesting leaves. Up to 4 applications can be made per year. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Note: Quilt Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt Fungicide may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year of Quilt Fungicide.
- 2) Do not apply more than 14 fl oz/A/application of Quilt Fungicide.
- 3) Do not apply more than 0.45 lb ai propiconazole-containing product/A/year.
- 4) Do not apply more than 1.5 lb ai azoxystrobin-containing product/A/year.
- 5) Do not exceed 4 applications per year.
- 6) Do not apply within 7 days of harvest (7-day PHI).

Quilt Fungicide Rate Conversion Table

Fl oz/A	Lb ai/A Propiconazole	Lb ai/A Azoxystrobin	Treated acres per gallon product
3	0.025	0.015	42.6
4	0.033	0.020	32.0
7	0.056	0.034	18.3
10.5	0.085	0.051	12.2
14	0.113	0.070	9.1
15	0.122	0.073	8.5
20.5	0.167	0.100	6.2
27.5	0.224	0.134	4.6
34.5	0.280	0.169	3.7

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [equal to or less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Quilt Fungicide®, Plant Performance™, Quadris®, Tilt®, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company.

Topsin® is a trademark of Pennwalt Corporation.

Viton® is a trademark of E.I. DuPont de Nemours and Company.

©20XX Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

Quilt Fungicide 1178 MAS 1217 AMEND-D 0118 –ep- 6/14/18
000100-01178.20180126D.QUILT-FUNGICIDE_AMEND_0118