

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms Pat Dinnen Syngenta Crop Protection P.O.BOX 18300 Greensboro, NC 27419

NOV 27 2009

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated September 21, 2009 for:

EPA Registration 100-1187

Uniform®

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

Registration Division (7505P)

Office of Pesticide Programs

I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or occord

Title Regulatory Specialist

Date

September 21, 2009

both under applicable law.

Signature

Pat dinnen

Typed Name

Pat Dinnen

c(Stamped)

Certification with Respect to Label Integrity

Version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL						
EPA Registration # Date Submitted to EPA Electronic file name						
100-1187	9/21/2009	000100-01187.20090917.UNIFORM_PRN2007-4_SEP2009.pdf				

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Pat oblinsen	September 21, 2009
Signature	Date
Pat Dinnen	_
Name (typed)	
Label Group Leader	_
Title	_



40x Ve

Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, NC 27419-8300 www.syngenta.com

syngenta

FedEx

September 21, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attention: Ms. Linda Arrington

SUBJECT: UNIFORM®

EPA REG. NO. 100-1187

NOTIFICATION OF LABEL CHANGE PER PR NOTICE 2007-4

Syngenta Crop Protection, Inc. is submitting Notification for Uniform, EPA Reg. No. 100-1187. Syngenta is amending the Storage and Disposal section of the label by Notification according to the directions stated in PR Notice 2007-4.

Attached are:

- One copy of the label with the changes clearly marked
- One unmarked copy of the label
- A CD of the unmarked copy of the label for "Electronic Comparison and Review"
- Certificate with Respect to Label Integrity Form
- Completed EPA Form 8570-1

Thank you in advance for approving this Notification. If you have any questions, please contact me at 336-632-2494.

Sincerely,

Pat dinnen

Pat Dinnen Label Group Leader Regulatory Affairs

Enclosures



Uniform®

NOV 2 7 2009

NOTIFICATION

Fungicide

A premix of azoxystrobin and mefenoxam for broad-spectrum disease control in cotton.

Active Ingredients:

Azoxystrobin*: methyl (E)-2-{2-[6-(2-cyanophenoxy)

 pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate
 28.2%

 Mefenoxam**:
 10.9%

 Other Ingredients:
 60.9%

 Total:
 100.0%

Contains 2.68 lbs. of azoxystrobin per gallon Contains 1.04 lbs. of mefenoxam per gallon

*IUPAC

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.


EPA Reg. No. 100-1187

EPA Est. 100-NE-001

SCP 1187A-L1A 0406

1 gallon [Product ID 17956]

Net Contents



^{**}CAS Nos. 70630-17-0 and 69516-34-3

	(20816)	
1	WON 14	

FIRST AID					
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 				
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 				
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 				
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. 				
Have the product doctor, or going	t container or label with you when calling a poison control center or for treatment.				
For 24	HOT LINE NUMBER Hour Medical Emergency Assistance (Human or Animal) or nical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372				

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.
- · Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Notify state and/or federal authorities and SYNGENTA immediately if you observe any adverse environmental effects due to use of this product.




Uniform Page 4 of 13

SOFTU"

Azoxystrobin, can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use. Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use.

Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. This product is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas.



CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY


NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. To the fullest extent allowed by State law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the fullest extent allowed by State law, SYNGENTA or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Uniform is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray Uniform where spray drift may reach apples.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Worker entry into treated areas is allowed immediately (0 hours) after application.

This product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONT	ROL.	
[STORAGE AND DISPOSAL section has been moved to the end of the	lapel.↓	: · .
	••••	•••
	•	••••

GENERAL INFORMATION

Uniform is a broad spectrum preventative fungicide with systemic properties for control of many soilborne plant diseases. All applications should be made according to the use directions that follow.

Uniform can provide control of many soilborne diseases when applied as an in-furrow application at planting. In-furrow applications will provide control of pre- or post-emergence damping off and diseases that infect plants at the soil-plant interface.

Chemigation

Do not apply this product through any type of irrigation equipment.

GENERAL USE PRECAUTIONS

Rotational Crop Restrictions

	Planting Time From	
Pototional Cron	Last Uniform Application	
Rotational Crop		
Brassica, Head and Stem subgroup	0 Days	
Carrot		
Cotton		
Cucurbits		
Eggplant		
Bulb Vegetables		
Legume Vegetables		
Peanuts	·	
Potatoes		
Strawberries		
Sugar Beets		
Tomatoes		
Soybeans		
Cereal Grains (other than Corn)	14 days	
Leafy Vegetables (except Brassica)	36 Days	
Brassica, Leafy Greens subgroup; Vegetables,		••••
Root subgroup; Vegetables, Tuberous and Corm		•••••
subgroup; and Vegetables Leaves of Root and		: .**.
Tuber group.		••••
Corn	9 months	
All Other Rotated Food or Feed Crops	12 months	
		• • •

RESISTANCE MANAGEMENT



Uniform fungicide is a mixture of a Group 4 (mefenoxam) and a Group 11 (azoxystrobin) fungicide. Uniform has two modes of action: (1) Phenylamides- affects RNA Synthesis [Group 4], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. SYNGENTA encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.


Uniform should not be alternated or tank mixed with any fungicide to which resistance has already developed.

INTEGRATED PEST (DISEASE) MANAGEMENT

Uniform should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The **SPECIFIC USE DIRECTIONS** section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area.

APPLICATION PROCEDURES

Apply Uniform as an in-furrow spray in 5-15 gals. of water per acre at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place. Uniform may be applied with all types of spray equipment commonly used for making in-furrow applications. Proper adjustments and calibration of spraying equipment is essential for good disease control.




Uniform DOF (Q Page 9 of 13

MIXING/SPRAYING

To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Uniform to the tank, allowing time for good dispersion. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operations. Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers should be thoroughly cleaned immediately after application.

Uniform is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used. Consult compatibility charts or your local or state agricultural authorities for compatibility information. Conduct a physical compatibility test as described in the paragraph below before making a field application.

Do not combine Uniform in the spray tank with pesticides, surfactants or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective and non-injurious under your conditions of use. If physical compatibility is unknown, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least twenty (20) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.



SPECIFIC USE DIRECTIONS

Crop	Target Diseases	Use Rate fl. oz./1000 ft. of row	Remarks
Cotton	Southern Blight (Sclerotium rolfsii) Rhizoctonia Seedling Blight (Rhizoctonia solani) Pythium Seedling Blight (Pythium aphanidermatum) Pythium Root Rot (Pythium ultimum)	0.32-0.48	Integrated Pest (Disease) Management: Uniform should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, plant residue management, crop rotation and proper water management. Apply Uniform as an in-furrow spray in 5-15 gals. of water per acre at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place. Make only one application per use season.




Table 1. Amount of Uniform (fluid ounces) per Acre Based on Row Spacing

Rate per 1000 Row Feet		22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
Fluid ounces of Product	Ounces of active ingredient (mefenoxam, (azoxystrobin)	Product per Acre (fluid ounces)						
0.32	0.05, 0.10	7.6	5.6	5.2	4.9	4.6	4.4	4.2
0.48	0.07, 0.15	11.4	8.4	7.8	7.4	7.0	6.6	6.3

40" = 13,068 row ft., 38" = 13,754 row ft., 36" = 14,520 row ft., 34" = 15,374 row ft., 32" = 16,315 row ft., 30" = 17,424 row ft., and 22" = 23,760 row ft./Acre

Table 2. Number of Acres per Gallon Product Based on Row Spacing

Use Rate	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
·	Acres Treated						
0.32 fl. oz.	16.8	22.8	24.6	26.1	27.8	29.0	30.5
0.48 fl. oz.	11.2	15.2	16.4	17.3	18.2	19.4	20.3

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original containers only. Store in a cool place. Protect from excessive heat. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Uniform
Page 12 of 13

Container Handling

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent); promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Uniform®, the Syngenta logo and the CP FRAME [frame] are trademarks of a Syngenta Group Company.

©2006 Syngenta

For non-emergency (e.g., current product information) call Syngenta Crop Protection at 1-800-334-9481

Manufactured for: Syngenta Crop Protection, Inc. Greensboro, North Carolina 27409 www.syngenta-us.com

UNI 1187A-L1A 0406 PRN2007-4 NOTIF-SEP2009 CLEAN - pl - 9/17/09 000100-01187.20090917.UNIFORM_PRN2007-4 SEP2009.pdf

