



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 23, 2010

Carolyn Brinkley
Syngenta Crop Protection, Inc.
PO Box 18300
Greensboro, NC 27419

re: Avicta 500 FS, EPA Reg. # 100-1204
label amendment submitted 3/4/2009, revised 5/5/2010, 7/15/2010 (D# 406876)
accepted with label comments (7/15/2010 version) and data conditions

Dear Ms. Brinkley:

This label amendment adds soybeans to this product. The Agency has determined that use as a soybean seed treatment would be a non-food use.

The revised labeling reference above, submitted in connection with the registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is conditionally acceptable provided that you comply with the following:

1. Make the following label changes before you release the product for shipment:
 - a. Directions for Use, first note - Add closed system concept so statement reads:
"Avicta 500 FS is for use only in Syngenta-certified seed treatment facilities using closed transfer and application equipment. Use in hopper box, ... is not permitted."
2. Submit studies to support the following data gaps. All studies are due within 18 months of this acceptance letter. *Note: Requirements (d) - (k) were previously levied under EPA Reg. # 100-1351 dated 4/6/2010.*
 - a. Conduct a survey of appropriate facilities and staff in order to provide information regarding the process of cleaning closed system seed treatment facilities. Survey questions and methodology should be discussed with EPA before the survey is conducted.
 - b. "Dust-off" studies on both untreated and treated soybean seeds using the Heubach apparatus. Since this is a new non-guideline study, a protocol outlining a standard methodology to be used for "dust-off" studies should be submitted to and approved by OPP before the study is run.
 - c. OPPTS 875.1200 and 875.1400 worker exposure monitoring studies using Avicta 500FS.

- d. OPPTS 850.1400- Early Life-Stage Toxicity Test
- e. OPPTS 850.4225 – Seedling Emergence, Tier II and OPPTS 850.4250 – Vegetative Vigor, Tier II.
- f. OPPTS 850.2300 - Avian reproduction Study.
- g. OPPTS 850.2100 - Acute Oral Toxicity with a Passerine Bird.
- h. Whole Sediment Toxicity Test: Chronic Invertebrates Freshwater and Marine. A protocol should be submitted to the Agency for review prior to testing.
- i. OPPTS 850.1075 – Fish Acute Toxicity Test, freshwater and marine; 850.1010- Aquatic Invertebrate Acute Toxicity test with Daphnia; 850.1025 or 1055 – Oyster Acute Toxicity Test (shell deposition) or Bivalve Acute Toxicity Test (embryo-larvae).
- j. OPPTS 850.1300 – Daphnia Chronic Toxicity Test.
- k. OPPTS 850.5400 – Algal Toxicity and 850.4400 Aquatic Plant Toxicity Test using Lemna spp.

3. Submit two copies of the revised final printed label for the record before you release the product for shipment.

If the above provisions are not complied with the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A copy of the label stamped "accepted with comments" is enclosed for your records. If you have any questions please contact Tom Harris at (703) 308-9423 or harris.thomas@epa.gov.

Sincerely yours,



John Hebert
 Product Manager
 Insecticide Rodenticide Branch
 Registration Division (7505C)
Hebert.John@epa.gov
 (703) 308-6249

enclosure

(Booklet)

RESTRICTED USE PESTICIDE
DUE TO TOXICITY.
FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Avicta® 500 FS

Nematicide

A Seed Treatment Product to Protect Cotton, Corn Grown for Seed, Field Corn, Popcorn, Soybean, and Sweet Corn Seedlings from Early Season Nematode Damage

For use only in Syngenta-certified seed treatment facilities and only in seed treatment equipment with closed transfer and application systems. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

Active Ingredient:	
Abamectin (CAS No. 71751-41-2).....	46.3%*
Other Ingredients:	53.7%
Total:	100.0%

*Contains 4.17 lbs. abamectin per gal.

KEEP OUT OF REACH OF CHILDREN.



DANGER / PELIGRO
POISON
(red type)

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1204
EPA Est. XXXX

SCP 1204-L (draft-c soybeans 7-15-10)

Net Contents

ACCEPTED
With COMMENTS
In EPA Letter, Dated:
JUL 23 2010

Under the Federal Insecticide, Fungicide and Rodenticide Act, As amended, for the pesticide Registered under EPA Reg. No:

100-1204

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
<p>Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).</p> <p>If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
HOT LINE NUMBER	
<p>For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals



DANGER / PELIGRO
POISON
(red type)

Fatal if inhaled. Do not breathe vapors. May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

The following personal protective equipment and clothing are required when using this product:

Corn and Cotton Seed Treatment - Personal Protective Equipment

I. Seed Treatment Workers

A. Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) must wear:

- Chemical resistant gloves made of any waterproof material – Category A (e.g., natural rubber \geq 14 mils)
- Protective eyewear
- Coveralls over long sleeved shirt and long pants
- Chemical-resistant apron
- Shoes and socks

B. Baggers and Bag Sewers must wear:

- Dust/mist filtering respirator (NIOSH-approved respirator with any N, R, P, or HE filter)
- Long sleeved shirt and long pants
- Shoes and socks

C. Workers involved with the clean up and maintenance of seed treatment equipment must wear:

- Chemical resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Protective eyewear
- A NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter
- Chemical-resistant coveralls over long sleeved shirt and long pants
- Shoes and socks

II. Multiple Task Workers must wear:

(Multiple task workers perform multiple tasks in one day such as mixing, product application, bagging, bag sewing, and clean up)

- Chemical resistant gloves (when handling treated seeds and equipment clean-up) made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Protective eyewear (when treating seed and during clean-up)
- Dust/mist filtering respirator (dust mask)
- Long sleeved shirt and long pants
- Chemical-resistant coveralls over long sleeved shirt and long pants (required when performing seed treatment equipment clean-up procedures)
- Shoes and socks

Soybean Seed Treatment - Personal Protective Equipment

I. Seed Treatment Workers

A. Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) must wear:

- Chemical resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Protective eyewear
- Coveralls over long sleeved shirt and long pants
- A quarter face cup-style respirator with a dust mist filter with MSHA/NIOSH approval number prefix TC21 or NIOSH approved respirator with any N, R, P, or HE filter
- Shoes and socks

B. Workers involved in packaging:

baggers, bag sewers, bag stackers or workers involved in filling, closing, and handling seed containers must wear:

- Chemical resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Protective eyewear
- A quarter face cup-style respirator with a dust mist filter with MSHA/NIOSH approval number prefix TC21 or NIOSH-approved respirator with any N, R, P, or HE filter
- Coveralls over long sleeved shirt and long pants
- Shoes and socks

C. Workers involved with the clean up and maintenance of seed treatment equipment must wear:

- Chemical resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Protective eyewear
- A half or full-face respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-21C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter
- Coveralls over long sleeved shirt and long pants
- Shoes and socks

Environmental Hazards

This pesticide is toxic to fish and wildlife. Exposed treated seeds may be hazardous to wildlife. Do not contaminate water when disposing of equipment wash water.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE:

Avicta 500 FS is only for use in Syngenta-certified seed treatment facilities; use in hopper box, planter box, slurry box, or other farmer-applied applications is not permitted.


- Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Conduct germination tests with a representative sample of the seed lot before committing the total seed lot to a selected seed treatment. Because seed quality and seed storage conditions are beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crops.
- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 180.920. Refer to the section on this label **REQUIRED SEED CONTAINER STATEMENTS** for language that must appear on the label that is attached to the bag of treated seed.

PRODUCT INFORMATION

To provide cotton, corn grown for seed, field corn, popcorn, sweet corn, and soybean seedlings early season protection from nematode damage, apply Avicta 500 FS to the seed of these crops. To add early season insect and disease protection, the seed treatment fungicides and insecticides specified in the Crop Use Directions can be mixed with Avicta 500 FS and applied to these seed.

REQUIRED SEED CONTAINER STATEMENTS

The Federal Seed Act requires that bags containing Avicta 500 FS-treated seeds shall be labeled with the following statements:

- **"Danger -- Poison **
- This seed has been treated with abamectin nematicide.
- Do not use treated seed for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing Avicta 500 FS-treated seed.

- Store away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and rubber gloves when handling treated seed.
- Plant Avicta 500 FS-treated seed based upon the local planting dates and soil temperatures that are recommended by your state agricultural extension service. In areas with a history of high nematode populations, a soil fumigant is recommended.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds that are spilled during loading and in areas such as row ends.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging in accordance with local requirements.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. **Exception:** Once the seeds are planted into soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the soil/media subsurface.

MIXING AND APPLICATION INSTRUCTIONS

- Consult the Syngenta Seed Treatment representative or the manufacturer of the application equipment you plan to use regarding the suitability of this application, equipment calibration, and operating instructions.
- Always treat seed in a well-ventilated area. Refer to the **Precautionary Statements** section of this label for personal protective equipment requirements and other safety precautions.
- Prepare no more mixture than is needed for the immediate operation.
- Apply Avicta 500 FS and its tank mix products as a water-based slurry utilizing standard slurry seed treatment equipment that will provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of nematode, insect, or disease control.
- Thoroughly mix the Avicta 500 FS and any other recommended seed treatment products with the amount of water required for the slurry treater and the dilution rate. (See **Avicta 500 FS in Tank Mixtures.**)
- Allow seed to dry before bagging/filling containers and store treated seed away from feed and foodstuffs.

Avicta 500 FS Alone:

1. To dispense Avicta 500 FS into a mix tank, first attach a pump with fittings (example: Micro Matic) that match the fittings on the Avicta 500 FS container.
2. Recirculate product. Recirculation time should be calculated based upon the ability of the pump to completely recirculate the volume of the tank 5-10 times within a 30-minute time period.
3. After the initial recirculation, a short recirculation period of 5 minutes per day is recommended.
4. Attach the dispensing hose to a dry-lock type of fitting that is permanently attached to the mix tank lid.
5. Add $\frac{1}{4}$ - $\frac{1}{2}$ of the required amount of water to the mix tank.
6. With the agitator running, add the Avicta 500 FS to the tank.
7. Continue agitation while adding the remainder of the water.
8. Begin application of the solution after the Avicta 500 FS has completely dispersed into the mix water.
9. Maintain agitation until all of the mixture has been applied.

Avicta 500 FS in Tank Mixtures:

If using Avicta 500 FS in a tank mixture with other seed treatment products,

- Confirm that the tank mix product is approved for the uses that are on the Avicta 500 FS label.
- Do not mix this product with any product that prohibits such mixing.
- Follow all directions for use, dilution ratios, precautions, and limitations that appear on the tank mix partner's label.
- Compare all of the product labels and follow the most restrictive label precautions and limitations. Do not exceed the maximum application rate on any of these labels.

Tank Mixing Instructions

- 1) Add $\frac{1}{2}$ of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
- 2) If using wettable powders and/or products in water-soluble packaging, add water-soluble packages first to clean water and allow them to completely disperse before adding Avicta 500 FS or other products.
- 3) Allow each tank mix partner to completely disperse before adding the next product.
- 4) Add the remaining amount of water and agitate.
- 5) Maintain agitation until the entire slurry mixture has been used.

CROP USE DIRECTIONS

Avicta 500 FS on Corn Seed

To provide the seedlings of corn grown for seed, field corn, popcorn, and sweet corn early season protection against nematode damage, Avicta 500 FS may be applied to the seed as specified in the following table. To add early season insect and disease protection Avicta 500 FS may be mixed with the products specified in the following table.

NOTE: In order to minimize dust off and loss of active ingredient from treated seed, all corn seed treated with Avicta 500 FS must receive an application of a dust reducing agent. Contact Syngenta for specific recommendations to insure acceptability of the dust-reducing agent with respect to retention of the active ingredient, seed germination and viability, and compatibility with other treatments that might be applied to the seed.

Pest/Disease	Amount of Product to Apply	Instructions
Nematodes	Apply Avicta 500 FS at the rate needed to achieve: 0.20 – 0.25 mg. abamectin per seed	<ul style="list-style-type: none"> To control early season nematodes, apply the amount of Avicta 500 FS specified in the center column. Use the higher rates if seed will be planted in areas where heavy nematode pressure is known to occur. <p>NOTE:</p> <ul style="list-style-type: none"> 1 fl. oz. of Avicta 500 FS contains 14.79 grams of abamectin The typical density of Avicta 500 FS is 9.01 lbs./gal.
Disease & Insect Protection – Early Season	For the amount to apply, refer to the labels for Apron XL®, Cruiser® 5FS, Dynasty®, and Maxim® XL.	To add early season disease and insect protection, treat corn seed with a mixture of Avicta 500 FS plus Apron XL, Cruiser 5FS, Dynasty, and/or Maxim XL

Avicta 500 FS on Cottonseed

To provide cotton seedlings early season protection against nematode damage, Avicta 500 FS may be applied to the seed as specified in the following table. To add early season insect and disease protection Avicta 500 FS may be mixed with the products specified in the following table.

NOTE: In order to minimize dust off and loss of active ingredient from treated seed, all cottonseed treated with Avicta 500 FS must receive an application of a dust reducing agent. Contact Syngenta for specific recommendations to insure acceptability of the dust-reducing agent with respect to retention of the active ingredient, seed germination and viability, and compatibility with other treatments that might be applied to the seed.

Pest/Disease	Amount of Product to Apply	Instructions
Nematodes	Apply Avicta 500 FS at the rate needed to achieve: 0.10– 0.15 mg. abamectin per seed	<ul style="list-style-type: none"> • To control early season nematodes, apply the amount of Avicta 500 FS specified in the center column. • Use the higher rates if seed will be planted in areas where heavy nematode pressure is known to occur. <p>NOTE:</p> <ul style="list-style-type: none"> • 1 fl. oz. of Avicta 500 FS contains 14.79 grams of abamectin • The typical density of Avicta 500 FS is 9.01 lbs./gal.
Disease & Insect Protection – Early Season	For the amount to apply, refer to the labels for Cruiser® 5FS* and Dynasty® CST.	To add early season disease and insect protection, treat cottonseed with a mixture of Avicta 500 FS plus Cruiser 5FS* and Dynasty CST.

*For use only on seed of delinted cotton

Avicta 500 FS on Soybean Seed

To provide soybean seedlings early season protection against damage from non-cyst forming nematodes, apply Avicta 500 FS to soybean seed as specified in the following table.

Use Restrictions

- Do not apply more than 177 gallons of this product per 8-hour day.
- In order to minimize dust off and loss of active ingredient from treated seed, all soybean seed treated with Avicta 500 FS must receive an application of a dust reducing agent. Contact Syngenta for specific recommendations to insure acceptability of the dust-reducing agent with respect to retention of the active ingredient, seed germination and viability, and compatibility with other treatments that might be applied to the seed.

To add early season insect and disease protection mix Avicta 500 FS with the specified insecticide and fungicide products that are also listed in this table.

Pest/Disease	Amount of Product to Apply	Instructions
<p>Non-Cyst Forming Nematodes¹</p>	<p>Apply Avicta 500 FS at the rate needed to achieve:</p> <p>0.10 – 0.15 mg. abamectin per seed</p>	<ul style="list-style-type: none"> • To control early season non-cyst forming nematodes, apply the amount of Avicta 500 FS specified in the center column. • Use the higher rates if seed will be planted in areas where heavy nematode pressure is known to occur. <p>NOTE:</p> <ul style="list-style-type: none"> • 1 fl. oz. of Avicta 500 FS contains 14.79 grams of abamectin • The typical density of Avicta 500 FS is 9.01 lbs./gal.
<p>Disease & Insect Protection – Early Season</p>	<p>For the amount to apply, refer to the labels for: ApronMaxx® brands, Apron XL, Cruiser 5FS, CruiserMaxx® and Maxim 4 FS,</p>	<p>To add early season disease and insect protection, treat soybean seed with a mixture of Avicta 500 FS plus ApronMaxx® brands, Apron XL, Cruiser 5FS, CruiserMaxx® and/or Maxim 4 FS</p>

¹Do not use Avicta 500 FS for protection against the soybean cyst nematode

CLEANING SEED TREATMENT EQUIPMENT

Do not use compressed air hoses to remove the dust from the seed treating equipment or the area around the equipment; use a central vacuum system or a HEPA filter portable vacuum cleaner.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal


Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. Do not reuse or refill this container.

Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, triple rinse the container as follows: empty the remaining contents into a rinsate collection system and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into a rinsate collection system or store rinsate for later disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration or by other procedures allowed by state and local authorities.

Apron®, ApronMaxx®, Avicta®, Cruiser®, CruiserMaxx®, Dynasty®, Maxim® the Syngenta logo, and the CP FRAME  are trademarks of a Syngenta Group Company Protected by U.S. Patent No. 6,875,727. Other patents pending.
©2010 Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1204A-L (draft-c soybeans 7-15-10)

AVI 500 FS1204A-M(draft-c soybeans 7-15-10) clean-lg-7-15-10

000100-01204.20100715.soybeansC.pdf

(Non-detachable Container Label)

RESTRICTED USE PESTICIDE
DUE TO TOXICITY.
FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S
CERTIFICATION.

Avicta™ 500 FS

Nematicide

A Seed Treatment Product to Protect Cotton, Corn Grown for Seed, Field Corn, Popcorn, Soybean and Sweet Corn Seedlings from Early Season Nematode Damage

For use only in Syngenta-certified seed treatment facilities and only in seed treatment equipment with closed transfer and application systems. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

Active Ingredient:	
Abamectin (CAS No. 71751-41-2).....	46.3%*
Other Ingredients:	53.7%
Total:	100.0%

*Contains 4.17 lbs. abamectin per gal.

KEEP OUT OF REACH OF CHILDREN.



DANGER / PELIGRO
POISON
(red type)

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements, PPE, and Directions for Use in booklet.

EPA Reg. No. 100-1204
EPA Est. XXXX

SCP 1204A-L (draft-c soybeans 7-15-10)

Net Contents

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals



DANGER / PELIGRO
POISON
(red type)

Fatal if inhaled. Do not breathe vapors. May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)
Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)
Call
1-800-888-8372

Environmental Hazards

This pesticide is toxic to fish and wildlife. Exposed treated seeds may be hazardous to wildlife. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. Do not reuse or refill this container.

Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, triple rinse the container as follows: empty the remaining contents into a rinsate collection system and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into a rinsate collection system or store rinsate for later disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration or by other procedures allowed by state and local authorities.

Avicta®, the Syngenta logo and the CP FRAME  are trademarks of a Syngenta Group Company

Protected by U.S. Patent No. 6,875,727. Other patents pending.

©2010 Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1204-M(draft-c soybeans 7-15-10)

AVI 500 FS 1204A-M(draft-c soybeans 7-15-10)clean-lg-7-15-10

000100-01204.20100715.soybeansC.pdf