



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

March 21, 2017

Patrick McCain  
Senior Regulatory Product Manager  
Regulatory Affairs, Seed Care  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419-8300

Subject: Label Amendment – revisions to PPE, seed bag tag, etc.  
Product Name: Avicta 400 FS  
EPA Registration Number: 100-1211  
Application Date: 9/9/2015  
Decision Number: 509989

Dear Mr. McCain:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 100-1211  
Decision No. 509994

with FIFRA section 6. If you have any questions, please contact Jacquelyn Herrick by phone at 703-347-0559, or via email at [herrick.jacquelyn@epa.gov](mailto:herrick.jacquelyn@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to be 'MS' with a large loop and a horizontal line extending to the right.

Mark Suarez  
Product Manager 07  
Invertebrate & Vertebrate Branch 3  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

**RESTRICTED USE PESTICIDE**  
**DUE TO ACUTE INHALATION TOXICITY.**  
**FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.**

**GROUP 6 INSECTICIDE**

**Avicta® 400 FS**

Seed Treatment

For Early Season Protection Against Nematodes in Cucurbits and Tomatoes

For use by Syngenta Crop Protection- certified professional vegetable seed treatment facilities only. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

Active Ingredient:

Abamectin (CAS No. 71751-41-2).....	37.0%*
Other Ingredients:	63.0%
<hr/> Total:	<hr/> 100.0%

\* Avicta 400 FS is a flowable concentrate containing 3.33 lb abamectin per gal.

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON**  
**PELIGRO / TOXICO**  
**POISON**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1211

EPA Est.

SCP

2.5 gallons  
\_\_\_\_\_gallons  
Net Contents

**ACCEPTED**  
Mar 21, 2017  
Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 100-1211

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<p style="text-align: center;"><b>NOTE TO PHYSICIAN</b></p> <p>Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).</p> <p>If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.</p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p style="text-align: center;"><b>HOT LINE NUMBER</b></p> <p style="text-align: center;">For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-888-8372</b></p>	

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals



**DANGER / POISON**  
**PELIGRO / TOXICO**  
**POISON**

Fatal if swallowed. May be fatal if inhaled. Harmful if absorbed through skin. Causes moderate eye irritation. Do not breathe vapors. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Avicta 400 FS must be used only in seed treatment equipment with closed transfer and application systems. The following personal protective equipment and clothing are required when using this product.

### Personal Protective Equipment (PPE)

**Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) must wear:**

- Chemical-resistant gloves made of: barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, or Viton®  $\geq$  14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant apron
- Shoes and socks
- NIOSH-approved respirator with any N, R, P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved power air purifying respirator with HE filter with NIOSH approval number prefix TC-21C

**Workers involved with packaging the treated seed (e.g. bagging, bag sewing, filling seed containers) must wear:**

- Chemical-resistant gloves made of: barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, or Viton®  $\geq$  14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants

- Shoes and socks
- NIOSH-approved respirator with any N, R, P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved power air purifying respirator with HE filter with NIOSH approval number prefix TC-21C

**Workers involved with the clean-up and maintenance of the seed treatment equipment must wear:**

- Chemical-resistant gloves (minimum length 12 inches) made of: barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, or Viton®  $\geq$  14 mils Important: Tape down gloves to the sleeves of the coveralls.
- Protective eyewear
- Chemical-resistant, hooded coveralls worn over long-sleeved shirt and long pants
- Shoes and socks
- A NIOSH-approved respirator with an organic vapor (OV) cartridge with a combination N, R, or P filter with NIOSH approval number prefix TC-84A; **OR** NIOSH approved gas mask with an organic vapor canister with NIOSH approval number prefix TC-14G; **OR** a NIOSH approved powered air purifying respirator with organic vapor (OV) cartridge and combination HE filter with NIOSH approval number prefix TC-23C

**Multiple Task Workers must wear:**

(Multiple task workers perform multiple tasks in one day such as mixing, product application, bagging/filling seed containers, bag sewing, and clean up.)

- Chemical resistant gloves made of: barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, or Viton®  $\geq$  14 mils. [Important: when performing seed treatment equipment clean-up procedures, gloves must be at least 12 inches long and should be taped down to the sleeves of the coveralls.
- Protective eyewear
- Long-sleeved shirt and long pants
- When performing seed treatment equipment clean-up procedures, chemical-resistant, hooded coveralls over long-sleeved shirt and long pants are required.
- Shoes and socks
- A NIOSH-approved respirator with an organic vapor (OV) cartridge with a combination N, R, or P filter with NIOSH approval number prefix TC-84A; **OR** NIOSH approved gas mask with an organic vapor canister with NIOSH approval number prefix TC-14G; **OR** a NIOSH approved powered air purifying respirator with organic vapor (OV) cartridge and combination HE filter with NIOSH approval number prefix TC-23C

## Environmental Hazards

This pesticide is toxic to fish and wildlife. Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading. Do not contaminate water when disposing of equipment washwater or rinsate.

Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 180.1001. Refer to the section on this label **SEED CONTAINER LABEL REQUIREMENTS** for language that must appear on the label that is attached to the container of treated seed.

---

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

---

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES**

**OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

### **DIRECTIONS FOR USE RESTRICTED USE PESTICIDE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS:**

- Only for use in Syngenta-certified professional seed treatment facilities. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.
- Must only be mixed with other seed treatments that are approved by the EPA for use on these crops and are compatible with Avicta 400 FS.

---

#### **PRODUCT INFORMATION**

---

Avicta 400 FS may be applied to the seed of tomatoes and crops in the cucurbits crop group.

When Avicta 400 FS is applied as a seed treatment, it provides protection from various nematodes during the early stages of plant growth and development.

#### **Avicta 400 FS:**

- Apply to primed or unprimed seed using standard and commonly accepted vegetable seed treatment processes that have confirmed efficacy and seed safety (i.e. slurry application, film coating, encrusting, or pelleting). Uneven or incomplete seed coverage may not give the desired level of early season protection against nematode damage. Depending upon the seed size and shape, highly specialized batch seed treatment application equipment may be required to ensure uniform treatment of the seed, to ensure safety to the seed, and to meet quality standards.



- Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Conduct germination tests on a representative sample of the seed lot before committing the total seed lot to a selected seed treatment. Because seed quality and seed storage conditions are beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crops.

---

## RESISTANCE MANAGEMENT

---

<b>GROUP</b>	<b>6</b>	<b>INSECTICIDE</b>
--------------	----------	--------------------

Avicta 400 FS contains abamectin, a Group 6 insecticide. Abamectin belongs to the avermectin class of chemistry which affects nerves and muscles.

Insect populations may contain individuals naturally resistant to Group 6 insecticides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refugia (untreated areas). Sound management programs also consider cultural and biological control practices.

### **In order to maintain susceptibility to this class of chemistry:**

- Use products at their full, specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless otherwise excluded by product labels. Do not exceed the maximum amount of this insecticide's mode of action allowed per growing season.
- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

Syngenta encourages responsible product stewardship to ensure effective long term control of the insect pests on this label.

**For additional information on Insect Resistance Management:**

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor

Visit the Insecticide Resistance Action Committee (IRAC) on the web at:


<http://www.irac-online.org>

---

## SEED CONTAINER LABEL REQUIREMENTS

---

The Federal Seed Act requires that containers containing treated seeds must be labeled with the following statements:

- **“Danger -- Poison 

The U.S. Environmental Protection Agency requires the following statements on containers containing vegetable seed treated with Avicta (abamectin) 400 FS:**

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting in areas (such as in row ends).
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Do not re-use bags from treated seed to handle food or feed products.
- This seed has been treated with x mg abamectin/seed.

- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. **Exception:** Once the seeds are planted into soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the soil/media substance.
- Treated seed must be planted at a depth of at least 0.5 inches.
- Do not use at a rate that will result in more than 0.17 lb of abamectin (81.6 g ai) as a cucurbits seed treatment application or 0.18 lb of abamectin (77.1 g ai) as a tomato seed treatment application per acre per calendar year.

---

## MIXING AND APPLICATION INSTRUCTIONS

---

**IMPORTANT:** Always treat seed in a well-ventilated area. Refer to the **Precautionary Statements** section of this label for personal protective equipment requirements and other safety precautions.

### Maximum Amount Allowed Per Day

The maximum amount of Avicta 400 FS that can be used in a seed treatment facility in one day is limited to 50 gallons of product.

### Important:

1. Consult the Syngenta Seed Treatment Representative or the manufacturer of the application equipment you plan to use regarding application, equipment calibration, and operating instructions.
2. Avicta 400 FS should be thoroughly recirculated/mixed in its original container before using.
3. Avicta 400 FS already contains an EPA-approved dye that will impart an unnatural color to the seed; adding additional dye or colorant to the seed treatment mixture is not required.
4. If carriers, colorants, or other materials are used when treating the seed, they must be EPA approved and known to be safe to the seed and compatible with Avicta 400 FS.
5. Thoroughly mix the recommended amount of Avicta 400 FS and other seed treatment products (i.e. fungicides, insecticides) with the amount of water required for the dilution rate and the slurry treater.
6. Prepare no more mixture than is needed for the immediate operation.
7. Store treated seed away from feed and foodstuffs.

### Applying Avicta 400 FS with compatible fungicides and insecticides

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions

and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Mixing Avicta 400 FS with fungicides and insecticides in the seed treating system is permissible. These fungicides and insecticides must be approved by the EPA for application to the vegetable seed specified in this label and recommended by Syngenta for use with Avicta 400 FS. These products can either be applied to the seed in slurry mixtures or applied in sequence, and applied continuously or in batches.

**Important:**

- **Do not mix Avicta 400 FS with any product that prohibits such mixing.**
- **The combination of Avicta 400 FS and compatible products must be tested to determine safety to all intended varieties or lines of seed.**
- **Follow all directions for use, crop/sites, dilution rates, precautions and limitations that are prescribed in the label for each product.**
- **Follow the most restrictive label precautions and limitations and do not apply more than the label permits.**

**A. Compatibility**

Before mixing Avicta 400 FS with other seed treatment products conduct a compatibility test before use.

1. Using a clear glass container, thoroughly mix all intended seed treatments with the appropriate amount of water.
2. Allow the mixture to sit for one hour.
3. Remix and observe for incompatibility.

**B. Tank Mixing Instructions**

1. To dispense Avicta 400 FS into a mix tank, first attach a pump with fittings that match the fittings on the closed system container for Avicta 400 FS.
2. Recirculate the product. Recirculation time should be calculated based upon the ability of the pump to completely recirculate the volume of the tank 5-10 times within a 30-minute time period.
3. After the initial recirculation, a short recirculation period of 5 minutes per day is recommended.
4. Attach the dispensing hose to a dry-lock type of fitting that is permanently attached to the mix tank lid.
5. Add ½ of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
6. If using wettable powders and/or products in water-soluble packaging, add water-soluble packages first to clean water and allow them to completely disperse before adding Avicta 400 FS or other products.
7. Allow each tank mix partner to completely disperse before adding the next product.

8. Add the remaining amount of water and agitate.
9. Maintain agitation until the entire slurry mixture has been used.

### **Applying Avicta 400 FS Alone:**

Although mixing Avicta 400 FS with fungicides and insecticides in the seed treating system is permissible, Avicta 400 FS may be applied alone as a slurry treatment to the seed specified in this label

### **Mixing Instructions**

1. First attach a pump with fittings that match the fittings on the closed system container for Avicta 400 FS.
2. Recirculate the product. Recirculation time should be calculated based upon the ability of the pump to completely recirculate the volume of the tank 5-10 times within a 30-minute time period.
3. After the initial recirculation, a short recirculation period of 5 minutes per day is recommended.
4. Attach the dispensing hose to a dry-lock type of fitting that is permanently attached to the mix tank lid.
5. Add  $\frac{1}{2}$  of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
6. Add Avicta 400 FS.
7. Add the remaining amount of water and agitate.
8. Maintain agitation until the entire slurry mixture has been used.

### Application Rate – Avicta 400 FS

Vegetable Seed	Amount of ai per Seed <sup>1</sup>	Instructions
<p><b>Cucurbits Crop Group</b></p> <p><u>Some of the crops in this group are:</u></p> <p>Cucumbers, all types Edible Gourds Gherkin Melons, all types (such as cantaloupes, casaba, honeydew, watermelons) <i>Momordica</i> spp. Pumpkins Summer Squash Winter Squash</p>	0.3 - 0.6 mg	To provide early season protection from nematode damage to seedlings, apply Avicta 400 FS to seed at a rate to achieve 0.3 - 0.6 mg ai per seed. <sup>1</sup>
<p><b>Tomatoes</b></p>		

<sup>1</sup> 1 fl oz of Avicta 400 FS contains 11.81 grams of abamectin and the typical density of Avicta 400 FS is 9.01 lb/gal.

---

### CLEANING SEED TREATMENT EQUIPMENT

---

Do not use compressed air hoses to remove the dust from the seed treating equipment or the area around the equipment; use a central vacuum system or a HEPA filter portable vacuum cleaner.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store in a cool, dry place.

#### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinse water is a violation of Federal Law. If these wastes cannot be disposed

of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### **Container Handling**

#### **Nonrefillable Container. Do not reuse or refill this container.**

Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Triple rinse the container as follows: Empty the remaining contents into application equipment or mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or mix tank or store rinsate for later disposal. Repeat this procedure two more times. Then contact Syngenta or its authorized agent for disposal.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

Avicta® and the SYNGENTA logo are Trademarks of a Syngenta Group Company

Protected by U.S. Patent No. 6875727. Other patents pending.

©200X Syngenta

For non-emergency (e.g. current product information) call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27409-8300