
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 100-1219	Date of Issuance: JAN 19 2005
	Term of Issuance: Conditional	
	Name of Pesticide Product: Azoxytrobin H&G	
Name and Address of Registrant (include ZIP Code): Syngenta Crop Protection P.O. Box 18300 Greensboro, NC 27419		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is <u>conditionally</u> registered in accordance with FIFRA Section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> Submit and/ or cite all data required for registration of your product under FIFRA section 3 (c) (5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA Section 4. Before releasing the product for shipment revise the EPA Registration Number to read, "EPA Reg. No. 100-1219" to your label before you release the product for shipment. Under the section "Precautionary Statement" add: "...and before eating, drinking, chewing gum or using tobacco" to the last sentence of the first paragraph. Move the remaining paragraphs to the section "Directions for Use". Delete the "User Safety Requirements" and "User Safety Recommendations" sections as they appear throughout the label. Since this product is intended for homeowner use Worker Protection Standard statements are not required. Under "Environmental Hazards" delete the following sentence from the second paragraph: "Do not apply directly to water except as specified on this label". Under the section "Conditions of Sale and Limitation of Warranty..." revise the statement beginning with: "In no event shall Syngenta or seller..." to read: "To the extent permitted by law Syngenta or seller shall not be liable...". As currently written the statement negates and/or detracts from labeling requirements as set forth by the Agency in 40 CFR Part 156, and is, therefore, considered misbranding." 		

- 7. Add a period at the end of the fourth sentence under "Mixing Procedures/Pump-up Sprayer (for all uses)."
- 8. Delete "...and more" from the label claim "Rust and more." Vague claims or claims implying a broader range of control than labeled are not acceptable.
- 9. On page 37, under Stone Fruits/Comments "This" should not be capitalized.
- 10. On page 70, under Corn(Sweet and Pop) /Disease "Leaf Blight" recurs twice and should appear once.
- 11. Please submit two (2) copies of your final printed label for before your release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the Master label is enclosed for your records.

If you have any questions regarding this action, please contact Bryant Crowe of my team at (703) 305-0025.

<p>Signature of Approving Official:</p>  <p>Olga Odion, Acting Product Manager (22) Fungicide Branch, Registration Division (7505C)</p>	<p>Date:</p> <p>JAN 19 2005</p>
--	---------------------------------

Enclosure

3/73

**Azoxystrobin H&G
Fungicide Concentrate**

Active Ingredient:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)
pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate^{1*} 22.9%

Inert Ingredients: 71.1%

Total: 100.00%

Contains 2.08 lbs. of active ingredient per gallon

¹CAS No. 131860-33-8

*IUPAC

Product of the United Kingdom

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on label.

EPA Reg. No. 100-

EPA Est.

SCP


Net Contents/Net Weight

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN 19 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

100-1219



Appendix 1 - Consolidated List of Label Claims

- Broad spectrum and systemic disease control on ornamentals, flowers, home lawns, vegetables, fruits, trees, and shrubs
- Broad spectrum and systemic disease control on ornamentals, flowers, trees, and shrubs
- Broad spectrum and systemic disease control on home lawns
- For use on home lawns.
- Broad spectrum and systemic disease control on vegetables and fruits
- Controls brown patch, powdery mildew, anthracnose, rust, blights, Scab, and more
- Prevents and stops major diseases on vegetables, flowers, lawns, trees, and shrubs
- Concentrate - Mix with water and spray
- For use in pump-up and hose-end sprayers
- 4 oz. will treat up to 12,000 sq. ft. of lawn
- 8 oz. will treat up to 24,000 sq. ft. of lawn
- 16 oz. will treat up to 48,000 sq. ft. of lawn
- Absorbed into the plant - Can't be washed off by rain or sprinklers after spray has dried
- Rainfast shortly after spray has dried
- Not affected by rain after spray has dried
- Systemic action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on home lawns and ornamentals
- Works up to 4 weeks for certain diseases
- Prevents and stops ...
 - Anthracnose Brown Patch
 - Blights Powdery Mildew
 - Rust Leaf Spot
 - and more

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash thoroughly with soap and water after handling.

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.



People and pets should not be allowed in treated areas until spray has dried.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available; for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Alternate Warranty Statement

Notice: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The buyer assumes responsibility for lack of performance or safety if not used according to the instructions on this label.

DIRECTIONS FOR USE

For Outdoor Residential Use Only

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR BEST RESULTS READ AND FOLLOW ALL LABEL DIRECTIONS

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-888-8372, day or night.

Pesticide/Container Disposal

If container is partially filled:

Important: It is best to use all of the product in accordance with label directions.

If it is necessary to dispose of unused product, please follow any applicable state or local guidelines or call your local solid waste agency.

Never place unused product down any indoor or outdoor drain.

If Container is Empty:

Do not reuse the container.

Place container in the trash.

GENERAL INFORMATION

This water-based systemic fungicide effectively prevents and controls major diseases on flowers, fruits, lawns, ornamentals, shrubs, trees, and vegetables.

USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.
- **DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.**
- If you need more information about fruit, vegetable, lawn and ornamentals diseases, contact your local County Extension Service.

To control diseases on plants other than turfgrass, apply product to the plant until it begins to run off of the leaves.

- Allow spray to completely dry before watering.

MIXING PROCEDURES

Shake well before using.

NOTE: ¼ fl. oz = 1½ teaspoons
 ½ fl. oz. = 1 Tablespoon = 3 teaspoons
 1 fl. oz. = 2 Tablespoons = 6 teaspoons

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides.

Move indoor houseplants outdoors before treating. Allow to dry before handling or moving indoors.

Pump-Up Sprayer (for all uses)

- Fill with ½ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

Concentrate Plus Water Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).

- From the following table, determine the amount of concentrate required.
- Pour concentrate into sprayer jar to the required "fl. oz." level.
- Then add the required amount of water to the "gals." level.
- Replace sprayer top on jar and shake well.
- Attach sprayer to hose and apply.
- For Lawns, apply to the measured square foot area to be sprayed.
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

Dial-Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).

- Fill the sprayer jar with only enough product to cover the plants or the area to be treated.
- Set Dial on the "fl. oz./gallon" level specified in the following table.
- Close sprayer, attach to hose and apply.
- For Lawns, apply to the measured square foot area to be sprayed.
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

**(Optional language for product sold in a HOSE-END SPRAYER package)
HOSE-END SPRAYER: Easy to Use. (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).**

- Make sure knob control is in the Off position. Shake well and attach to hose.
- Turn water on at faucet, aim nozzle toward lawn or desired plant and turn control lever to ON to begin spraying.
- When finished turn control lever to OFF position. Discharge residual water pressure by turning ON and OFF again.
- For Lawns, apply to the [xxx] square foot area to be sprayed. [Square feet value to be inserted per specific package size].
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

APPLICATION INSTRUCTIONS



People and pets should not be allowed in treated areas until spray has dried.

Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

Apply sufficient water to obtain coverage without excessive runoff.

13/73

FLOWERS (annuals and perennials)

General Directions

Do not apply more than two applications before alternating with a fungicide that has a different mode of action.

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Roses	Downy Mildew Leaf Spot Powdery Mildew Rusts	1/4-1/2 tsp.	Thorough coverage needed for best results. Begin application at bud break. Treat every 7-21 days.
Annual and Perennial Flowers-Bedding Plants	Downy Mildew Leaf Spots Leaf Blights Powdery Mildew Rusts Scab	1/8-1/4 tsp.	Apply every 7-14 days prior to infection. Do not apply the 3 teaspoon rate on less than 14 day spray intervals.
Annual and Perennial Flowers/Mature Plants	Downy Mildew Leaf Spots Leaf Blights Powdery Mildew Rusts Scab	1/4-1/2 tsp.	Apply before disease begins. Apply every 7-28 days.
Geraniums	Flower Blights	1/8-1/2 tsp.	Apply before disease begins. Apply every 7-21 days.
Petunias Vinca	Shoot/Stem Diseases	1/8-1/4 tsp.	Apply every 7-28 days.
Chrysanthemums Cyclamen Impatiens Sweet Alyssum	Soil-Borne Diseases	1/8-1/4 tsp.	Use as a directed spray. Apply every 7-21 days.

14/73

LAWNS

- 1) Apply after mowing or allow sprayed area to completely dry before mowing.
- 2) Measure the length and width of area to be treated before application. Length x width = square feet. For example, an area 25 ft. long and 20 ft. wide is 500 sq. ft.
- 3) Begin applications when conditions are favorable for disease infection, prior to disease symptoms.
- 4) Can be safely applied after seeding or at seed germination.

Disease	How Much to Apply (tsp. per 250 sq. ft.)	Comments
Brown Patch	½-1 tsp.	First application should be in May or June before disease is present. If needed, repeat treatment every 14-28 days.
Anthracnose Fusarium Patch Pink Patch Summer Patch Gray Leaf Spot Leaf Spot Rhizoctonia Leaf Spot Necrotic Ring Spot Leaf Rust Stem Rust Stripe Rust Melting Out Powdery Mildew Southern Blight Red Thread	½-1 tsp.	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Apply every 14-28 days.
Cool Weather Brown Patch Yellow Patch	½-1 tsp.	Make 1 or 2 applications in fall (14-28 days interval) or when conditions are favorable for disease development.
Take-All Patch	½-1 tsp.	Apply twice in the Fall (Sept. and Oct.) and twice in the Spring (April and May). Southern lawns also need to be treated twice in the summer (June-August).

15/73

Disease	How Much to Apply (tsp. per 250 sq. ft.)	Comments
Snowmold (Gray or Pink)	1½ tsp.	Make a single application in late fall. or
	1 tsp.	Apply twice in the fall prior to snow cover waiting 10 -14 days between each application.
Pythium Blight Pythium Root Rot	½-1 tsp.	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Apply every 10-14 days. For use on newly seeded as well as established lawns.
Zoysia Patch	½-1 tsp.	Apply 1-2 applications approximately one month prior to zoysiagrass dormancy (14-28 day interval)

10/13

TREES, ORNAMENTALS AND SHRUBS

General Directions

Do not apply more than two applications before alternating with a fungicide that has a different mode of action.

Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Conifers Cedar Christmas Trees Cypress Fir Juniper Pine Spruce	Conifer Blights	1/8-1/2 tsp.	Apply every 7-28 days.

17/73

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Abelia Alder Arborvitae Aspen Azalea Bougainvillea Butterfly bush (<i>Buddleia</i>) Boxwood Camellia Cedar Clethra Coleus Crapemyrtle Dogwood Dumbcane (<i>Diffenbachia</i>) Euonymus Fatsia Fig* Forsythia Grasses Hawthorn Heather Hibiscus Hydrangea Ivy Japanese andromeda Lirope Magnolia Maples Nandina Oak Oleander Palms Philodendron Photinia Plum (flowering or purple-leaf) Poinsettia Pothos Red Bud Rhododendron Rose of Sharon Rubber tree Sedum Viburnum Wiegela Zebra plant	Downy Mildew Leaf Spot Scab	1/8-1/2 tsp.	Apply every 7-28 days. *Figs: Upon treating, do not use for food and/or feed purposes.
Azalea Dogwood Rhododendron	Anthracnose	1/2-1 tsp.	Apply every 7-28 days.

18/73

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Ajuga Azalea Barberry Birch Bradford's pear Ceanothus Crapemyrtle Dogwood Gardenia Hawthorn Hibiscus Hydrangea Holly Lilac Laurel Mock-orange Oak Pampas & Ribbon grasses Photinia Rhododendron Sage Spirea Viburnum Willow	Powdery Mildew	1/8-1/2 tsp.	Apply every 7-28 days.
Aucuba Azalea Boxwood Holiday cactus Caladium Cotoneaster Japanese andromeda Laurel Palm Rhododendron Rubber plant (<i>Peperomia</i>) Yew Yucca	Soil-Borne Diseases	1/8-1/2 tsp.	Use as a directed spray. Apply every 7-21 days.
Azalea Rhododendron	Shoot/Stem Diseases	1/8-1/4 tsp.	Apply every 7-28 days

19/73

FRUIT AND NUT TREES, BERRIES AND VINES: NOT FOR USE ON APPLES, CRAB-APPLES, OR FLOWERING (ORNAMENTAL) CHERRIES

General Directions

- 1) Do not apply more than two applications before alternating with a fungicide that has a different mode of action.
- 2) Do not make more than 4 applications of this product per crop (unless specified in use directions).
- 3) Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Almonds	Alternaria Leaf And Fruit Spot Anthracnose Brown Rot Blossom Blight Leaf Blight Leaf Rust Scab Shothole	1½-2 tsp.	28	Start applications during early bloom. Apply every 7-14 days. Do not make more than 4 applications of this product per crop.
Berries, Bushberry subgroup: Blueberry Currant Elderberry Gooseberry Huckleberry Including all cultivars and/or hybrids of these Lingonberry Juneberry Salal	Botryosphaeria Canker Powdery Mildew Septoria Blight Mummyberry Alternaria Fruit Rot Phomopsis Stem Canker Anthracnose Fruit Rot	1-2 tsp.	0	Apply every 7-14 days. Do not make more than 3 applications of this product per crop.

20/13

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
<p>Berries, Caneberry subgroup: Blackberry Bingleberry Boysenberry Dewberry Lowberry Marionberry Olallieberry Youngberry Loganberry Red and black raspberry Including all cultivars and/or hybrids of these</p>	<p>Botryosphaeria Canker Anthracnose Powdery Mildew Leaf Spot Colletotrichum Rot Spur Blight Rosette or Double Blossom of Blackberries</p>	<p>1-2 tsp.</p>	<p>0</p>	<p>Apply every 7-14 days. Do not make more than 6 applications of this product per crop.</p>
<p>Citrus Fruit Calamondin Citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma mandarin Tangerine Including all cultivars and/or hybrids of these</p>	<p>Greasy Spot Melanose Scab Albinism Post Bloom Fruit Drop (PFD) Alternaria Leaf and Fruit Spot</p>	<p>1½-2 tsp.</p>	<p>0</p>	<p>Apply every 7-21 days. Do not make more than 4 applications of this product per crop.</p>
<p>Grapes Including Muscadines</p>	<p>Downy Mildew Phomopsis Cane and Leaf Spot Powdery Mildew Black Rot Suppression Botrytis Bunch Rot</p>	<p>1½-2 tsp.</p>	<p>14</p>	<p>Apply every 10-14 days. Do not make more than 4 applications of this product per crop.</p>
<p>Pistachios</p>	<p>Alternaria Late Blight Botryosphaeria panicle and Shoot Blight Septoria Leaf Spot</p>	<p>1½-2 tsp.</p>	<p>7</p>	<p>Apply every 7-21 days. Do not make more than 3 applications of this product per crop.</p>

21/73

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Stone Fruits Apricot Cherry, sweet Cherry, tart Nectarine Peach Plum Plumcot Prune	Alternaria Spot and Fruit Rot Anthracnose Leaf Rust Powdery Mildew Scab Shot Hole Brown Rot Blossom Blight and Fruit Rot	1½-2 tsp.	0	Apply every 7-21 days. For Brown rot blossom blight begin spray at early bloom and continue through petal fall. Do not make more than 4 applications of this product per crop.
Strawberry	Anthracnose Powdery Mildew Suppression of Botrytis on the Foliage	2 tsp.	0	Apply every 7-10 days. Do not make more than 4 applications of this product per crop.
Tropical Fruit Pawpaw Persimmon	Anthracnose Cercospora Leaf Spot Powdery Mildew Rust	1-2 tsp.	0	Apply every 10-14 days. Do not make more than 4 applications of this product per crop.

22/13

VEGETABLES

General Directions

- 1) Do not apply more than two applications before alternating with a fungicide that has a different mode of action. This product is a group 11 fungicide. Do not alternate with another group 11 fungicide.
- 2) Do not make more than 6 applications of this product per crop (unless specified in use directions).
- 3) Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Asparagus	Stemphyllium purple spot	1½-2 tsp.	100	Apply on a 7-14 day schedule to asparagus foliage after harvest of spears is complete.
Brassica Head and Stem subgroup: Broccoli Chinese broccoli Brussel sprouts Cabbage Chinese cabbage Chinese mustard Cabbage Cauliflower Cavalo broccolo Kohlrabi Including all cultivars and/or hybrids of these	Alternaria Leaf Spot Downy Mildew	1-1¼ tsp.	0	Apply on a 7-14 day schedule.
Brassica Leafy Greens subgroup: Broccoli raab Cabbage Chinese Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Including all cultivars and/or hybrids of these	White Rust Black Spot Cercospora Leaf Spot	1-2 tsp.	0	Apply on a 7-14 day schedule.
	Soil-Borne Diseases Seedling Root Rot, Basal Stem Rot	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.

23/73

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Bulb Vegetables Garlic Leek Onion, bulb Onion, green Welsh onion Shallot	Foliar Diseases Cladosporium Leaf Blotch Purple Blotch Rust White Rot	1-1½ tsp.	0	Apply on a 7-14 day schedule.
	Botrytis Leaf Blight Downy Mildew	1¼-2 tsp.		
	Soil-Borne Diseases Rhizoctonia Damping Off	1/2-7/8 tsp. for 400 feet of row		
Carrots	Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Early Blight Late Blight Powdery Mildew Rust White Mold White Rust	1¼-2 tsp.	0	Apply on a 7-14 day schedule.
	Soil-Borne Diseases Rhizoctonia Root	1/2-7/8 tsp. for 200 feet of row		
Celery	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Early Blight Late Blight Powdery Mildew Septoria Leaf Spot White Rust	1¼-2 tsp.	0	Apply on a 7-14 day schedule
	Soil-Borne Diseases Rhizoctonia Root Rot	1/2-7/8 tsp. for 200 feet of row		
Corn Pop Sweet	Rust	1-1¼ tsp.	7	Apply on a 7-14 day schedule.
	Anthracnose Leaf Blight Eye Spot Gray Leaf Spot Northern Corn Leaf Blight Northern Corn Leaf Spot Southern Corn Leaf Blight	1¼-2 tsp.		

24/73

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Cucurbits Cantaloupe Chayote Chinese-waxgourd Cucumber Gourds Honeydew Melons (bitter melon, balsam apple) Muskmelon Watermelon Pumpkin Squash Zucchini Including cultivars and/or hybrids of these	Anthracnose Belly Rot Downy Mildew Gummy Stem Blight Leaf Spots Myrothecium Canker Powdery Mildew	1½-2 tsp.	1	Make a preventative application on a 5-7 day schedule. For belly rot apply at the 1-3 leaf crop stage with second application made just prior to vine tip over or 10-14 days later, whichever occurs first.
	Soil-Borne Diseases Rhizoctonia root rot	1/2-7/8 tsp. for 200 feet of row		Apply after seeds planting by spraying over the row or directed to the base of transplants or emerged plants.
Leafy Vegetables (except brassica) Amaranth Arugula Cardoon Celery Celtnce Chervil Chrysanthemum-edible Coriander, leaves (Cilantro) Corn salad Cress Dandelion Dock Endive Fennel Lettuce, head and leaf Orach Parsley Purslane Radicchio Rhubarb Spinach Swiss Chard Including cultivars and/or hybrids of these	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Septoria Leaf Spot White Rust	1-2 tsp.	0	Make a preventative application on a 7-14 day schedule. Use lower rate for lettuce and spinach.
	Downy Mildew Powdery Mildew	1½-2 tsp.		
	Soil-Borne Diseases Webb Blight, Bottom Rot, Crater Rot, Root Rot	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.

25/13

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Legume Vegetables, dry and succulent : Bean (<i>Phaseolus spp.</i>) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (<i>Vigna spp.</i>) (includes adzuki bean, asparagus bean, blackeyed pea, cowpea, catjang, Chinese longbean, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava bean) Chickpea (garbanzo bean) Guar Jackbean Lablab bean (hyacinth bean), Pea (<i>Pisum spp.</i>) (includes dwarf pea, edible-pod pea, English pea, garden pea, green pea, field pea, snow pea, sugar snap pea). Pigeon pea Sword bean	Bean Rust Anthracnose Alternaria Leaf Spot Ascochyta Leaf Spot Rust Southern Blight Web Blight Ascochyta Blight Ascochyta Leaf And Pod Spot Alternaria Blight	1 tsp. 1-2 tsp.	0	Make a preventative application on a 7-14 day schedule.
	Soil-Borne Diseases Rhizoctonia Root Rot	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Mint (fresh)	Rust Powdery Mildew	1-2 tsp.	0	Apply on a 7-10 day schedule. Do not make more than 3 sprays/crop.
Pepper Bell Pepper Non-Bell Pepper Sweet Non-Bell Pepper	Powdery Mildew Anthracnose	1-2 tsp.	0	Apply on a 7-14 day schedule.
	Soil-Borne Diseases Rhizoctonia Seedling Rot	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Eggplant Okra				

26/73

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Potatoes	Early Blight Late Blight Black Dot Powdery mildew	1-2 tsp.	14	Begin application when plants are 6 inches high or at prior to first sign of disease. Do not make more than 2 sprays before alternating to a different product with alternate chemistry. Do not make more than 6 applications per season. 7 inches or less.
	Late Blight	2 tsp.		
	Soil-Borne Diseases Black Scurf Silver Scurf	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Tomatoes	Anthracnose Black Mold Buckeye Rot Early Blight Powdery Mildew Septoria Leaf Spot Target Spot	¾ tsp.	0	Do not apply until 21 days after transplanting or 35 days after seeding. Apply on a 7-21 day schedule. For late blight apply on a 5-7 day schedule. Make only one application before alternating to another product with a different chemistry.
	Late Blight	1 tsp.		
Vegetables, leaves of root and tuber, group Beet, garden and sugar Burdock Carrot Cassava, bitter and sweet Celeriac (celery root) Chervil, turnip-rooted Chicory Dasheen (taro) Parsnip Radish Radish, oriental (daikon) Rutabaga Salsify, black Sweet potato Tanier Turnip Yam, true	Foliar Diseases Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Powdery Mildew Rust White Rust	1-2 tsp.	0	Apply every 7-14 days. Use 1 ¼ -2 tsp. rate for Cercospora leaf spot and Powdery Mildew. For Powdery Mildew apply on 5-7 day schedule. Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
	Soil-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	1/2-7/8 for 200 feet of row		

27/73

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Vegetables, root, subgroup Beet, garden and sugar Burdock Carrot Celeriac Chervil, turnip-rooted Chicory Ginseng Horseradish Parsley, turnip-rooted Parsnip Radish Radish, oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Alternaria Leaf Spot Ascochyta Leaf Spot Rust White Rust	1-2 tsp.	0	Apply every 7-14 days.
	Cercospora Leaf Spot Powdery Mildew	1.25 -2 tsp.		For Powdery Mildew apply on 5-7 day schedule.
	Soil-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	1/2-7/8 for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Vegetables, tuberous and corm, subgroup Arracacha Arrowroot Artichoke, Chinese and Jerusalem Canna Cassava, edible, bitter and sweet Chayote (root) Chufa Dasheen (Taro) Ginger Leren Potato Sweet Potato Tanier Tumeric Yam, bean Yam, true	Foliar Diseases Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Powdery Mildew Rust White Rust	1-2 tsp.	14	Apply every 7-14 days. Use 1 ¼ -2 tsp. rate for Cercospora leaf spot and Powdery Mildew. For Powdery Mildew apply on 5-7 day schedule.
	Soil-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	1/2-7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Watercress	Cercospora Leaf Spot	1-2 tsp.	7	Apply on a 7-10 day schedule.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

28/73

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27419
www.syngenta.com

29/73

**Azoxystrobin H&G
Fungicide Concentrate**

Active Ingredient:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)
pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate^{1*} 22.9%

Inert Ingredients: 71.1%

Total: 100.00%

Contains 2.08 lbs. of active ingredient per gallon

¹CAS No. 131860-33-8

*IUPAC

Product of the United Kingdom

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on label.

EPA Reg. No. 100-

EPA Est.

SCP

Net Contents/Net Weight

Appendix 1 - Consolidated List of Label Claims

- Broad spectrum and systemic disease control on fruits, nuts, berries and vines
- Broad spectrum and systemic disease control on fruits, nuts, berries and vines
- Controls brown patch, powdery mildew, anthracnose, rust, blights. Scab, and more
- Prevents and stops major diseases on fruits, nuts, berries and vines
- Concentrate - Mix with water and spray
- For use in pump-up and hose-end sprayers
- Absorbed into the plant - Can't be washed off by rain or sprinklers after spray has dried
- Rainfast shortly after spray has dried
- Not affected by rain after spray has dried
- Systemic action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on fruits, nuts, berries and vines
- Works up to 4 weeks for certain diseases
- Prevents and stops ...
 - Anthracnose Brown Patch
 - Blights Powdery Mildew
 - Rust Leaf Spot
 - and more

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash thoroughly with soap and water after handling.

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.



People and pets should not be allowed in treated areas until spray has dried.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available; for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Alternate Warranty Statement

Notice: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The buyer assumes responsibility for lack of performance or safety if not used according to the instructions on this label.

DIRECTIONS FOR USE

For Outdoor Residential Use Only

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR BEST RESULTS READ AND FOLLOW ALL LABEL DIRECTIONS

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-888-8372, day or night.

Pesticide/Container Disposal

If container is partially filled:

Important: It is best to use all of the product in accordance with label directions.

If it is necessary to dispose of unused product, please follow any applicable state or local guidelines or call your local solid waste agency.

Never place unused product down any indoor or outdoor drain.

If Container is Empty:

Do not reuse the container.
Place container in the trash.

GENERAL INFORMATION

This water-based systemic fungicide effectively prevents and controls major diseases on fruit and nut trees, berries and vines.

USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.
- **DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.**
- If you need more information about fruits, nuts, berries and vines diseases, contact your local County Extension Service.

To control diseases apply product to the plant until it begins to run off of the leaves.

- Allow spray to completely dry before watering.

MIXING PROCEDURES

Shake well before using.

NOTE: $\frac{1}{4}$ fl. oz = $1\frac{1}{2}$ teaspoons
 $\frac{1}{2}$ fl. oz. = 1 Tablespoon = 3 teaspoons
 1 fl. oz. = 2 Tablespoons = 6 teaspoons

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides.

Move indoor houseplants outdoors before treating. Allow to dry before handling or moving indoors.

Pump-Up Sprayer (for all uses)

- Fill with $\frac{1}{2}$ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

Concentrate Plus Water Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).

- From the following table, determine the amount of concentrate required.
- Pour concentrate into sprayer jar to the required "fl. oz." level.

- Then add the required amount of water to the "gals." level.
- Replace sprayer top on jar and shake well.
- Attach sprayer to hose and apply.
- For fruits, nuts, berries and vines, apply to obtain sufficient spray coverage without excessive runoff.

Dial-Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).

- Fill the sprayer jar with only enough product to cover the plants or the area to be treated.
- Set Dial on the "fl. oz./gallon" level specified in the following table.
- Close sprayer, attach to hose and apply.
- For fruits, nuts, berries and vines, apply to obtain sufficient spray coverage without excessive runoff.

**(Optional language for product sold in a HOSE-END SPRAYER package)
HOSE-END SPRAYER: Easy to Use. (use for application to fruits, nuts, berries and vines).**

- Make sure knob control is in the Off position. Shake well and attach to hose.
- Turn water on at faucet, aim nozzle toward lawn or desired plant and turn control lever to ON to begin spraying.
- When finished turn control lever to OFF position. Discharge residual water pressure by turning ON and OFF again.
- For fruits, nuts, berries and vines, apply to obtain sufficient spray coverage without excessive runoff.

APPLICATION INSTRUCTIONS



People and pets should not be allowed in treated areas until spray has dried.

Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

Apply sufficient water to obtain coverage without excessive runoff.

38/13

FRUIT AND NUT TREES, berries and vines: NOT FOR USE ON APPLES, CRAB-APPLES, OR FLOWERING (ORNAMENTAL) CHERRIES

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

General Directions

- 1) Do not apply more than two applications before alternating with a fungicide that has a different mode of action.
- 2) Do not make more than 6 applications of this product per crop unless specified in comments.
- 3) Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Almonds	Alternaria Leaf And Fruit Spot Anthracnose Brown Rot Blossom Blight Leaf Blight Leaf Rust Scab Shothole	1½-2 tsp.	28	Start applications during early bloom. Apply every 7-14 days. Do not make more than 4 applications of this product per year.
Berries, Bushberries Blueberry Currant Elderberry Gooseberry Huckleberry Lingonberry Juneberry Salal	Botryosphaeria Canker Powdery Mildew Septoria Blight Mummyberry Alternaria Fruit Rot Phomopsis Stem Canker Anthracnose Fruit Rot Leafspot Colletotrichum Rot Spur Blight	1-2 tsp.	0	Apply every 7-14 days. Do not make more than 3 applications of this product per crop.
Berries, Caneberry Blackberry Boysenberry Dewberry Lowberry Marionberry Olallieberry Youngberry Loganberry	Colletotrichum Rot Spur Blight Rosette Or Double Blossom Of Blackberries	1-2 tsp.	0	Apply every 7-14 days. Do not make more than 6 applications of this product per crop.

Master Label for Expert Gardener and Homeowner
Home Owner FRUIT AND NUT TREES, berries and vines

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Citrus Fruit Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Tangerine	Greasy Spot Melanose Scab Albinism Post Bloom Fruit Drop (PFD) Alternaria Leaf And Fruit Spot	1½-2 tsp.	0	Apply every 7-21days. Do not make more than 6 applications of this product per crop.
Grapes Including Muscadines	Downy Mildew Phomopsis Cane And Leaf Spot Powdery Mildew Black Rot Suppression Botrytis Bunch Rot	1½-2 tsp.	14	Apply every 10-14 days. Do not make more than 4 applications of this product per crop.
Pistachios	Alternaria Late Blight Botryosphaeria panicle and shoot blight Septoria leaf spot	1½-2 tsp.	7	Apply every 7-21 days. Do not make more than 4 applications of this product per crop.
Stone Fruits Apricot Cherry, sweet Cherry, tart Nectarine Peach Plum Plumcot Prune	Alternaria Spot and Fruit Rot Anthracnose Leaf Rust Powdery Mildew Scab Shot Hole Brown Rot Blossom blight and Fruit rot	1½-2 tsp.	0	Apply every 7-21 days. For Brown rot blossom blight begin spray at early bloom and continue through petal fall. Do not make more than 4 applications of This product per crop.
Strawberry	Anthracnose Powdery mildew Suppression of Botrytis on the foliage	1-2 tsp.	0	Apply every 7-10 days. Do not make more than 4 applications of This product per crop.
Tropical Fruit Pawpaw Persimmon	Anthracnose Cercospora leaf spot Powdery Mildew Rust	1-2 tsp.	0	Apply every 10-14 days. Do not make more than 6 applications of This product per crop.

**Azoxystrobin H&G
Fungicide Concentrate**

Active Ingredient:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)
pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate^{1*} 22.9%

Inert Ingredients: 71.1%

Total: 100.00%

Contains 2.08 lbs. of active ingredient per gallon

¹CAS No. 131860-33-8

*IUPAC

Product of the United Kingdom

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on label.

EPA Reg. No. 100-

EPA Est.

SCP

Net Contents/Net Weight

41/73

Appendix 1 - Consolidated List of Label Claims

- Broad spectrum and systemic disease control on home lawns
- For use on home lawns.
- Controls brown patch, powdery mildew, anthracnose, rust, blights. Scab, and more
- Prevents and stops major diseases on lawns
- Concentrate - Mix with water and spray
- For use in pump-up and hose-end sprayers
- 4 oz. will treat up to 12,000 sq. ft. of lawn
- 8 oz. will treat up to 24,000 sq. ft. of lawn
- 16 oz. will treat up to 48,000 sq. ft. of lawn
- Absorbed into the plant - Can't be washed off by rain or sprinklers after spray has dried
- Rainfast shortly after spray has dried
- Not affected by rain after spray has dried
- Systemic action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on home lawns
- Works up to 4 weeks for certain diseases
- Prevents and stops ...
 - Anthracnose Brown Patch
 - Blights Powdery Mildew
 - Rust Leaf Spot
 - and more

42/73

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash thoroughly with soap and water after handling.

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.



People and pets should not be allowed in treated areas until spray has dried.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available; for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

43/73

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

45/73

Alternate Warranty Statement

Notice: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The buyer assumes responsibility for lack of performance or safety if not used according to the instructions on this label.

DIRECTIONS FOR USE

For Outdoor Residential Use Only

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR BEST RESULTS READ AND FOLLOW ALL LABEL DIRECTIONS

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-888-8372, day or night.

Pesticide/Container Disposal

If container is partially filled:

Important: It is best to use all of the product in accordance with label directions.

If it is necessary to dispose of unused product, please follow any applicable state or local guidelines or call your local solid waste agency.

Never place unused product down any indoor or outdoor drain.

If Container is Empty:

Do not reuse the container.
Place container in the trash.

GENERAL INFORMATION

This water-based systemic fungicide effectively prevents and controls major diseases on lawns.

USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.
- **DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.**
- If you need more information about lawn diseases, contact your local County Extension Service.
- Allow spray to completely dry before watering.

MIXING PROCEDURES

Shake well before using.

NOTE: ¼ fl. oz = 1½ teaspoons
 ½ fl. oz. = 1 Tablespoon = 3 teaspoons
 1 fl. oz. = 2 Tablespoons = 6 teaspoons

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides.

Pump-Up Sprayer

- Fill with ½ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

Concentrate Plus Water Style Hose-End Sprayer.

- From the following table, determine the amount of concentrate required.
- Pour concentrate into sprayer jar to the required "fl. oz." level.
- Then add the required amount of water to the "gals." level.
- Replace sprayer top on jar and shake well.
- Attach sprayer to hose and apply.
- Apply to the measured square foot area to be sprayed.

Dial-Style Hose-End Sprayer.

- Fill the sprayer jar with only enough product to cover the plants or the area to be treated.
- Set Dial on the "fl. oz./gallon" level specified in the following table.
- Close sprayer, attach to hose and apply.
- Apply to the measured square foot area to be sprayed.

(Optional language for product sold in a HOSE-END SPRAYER package)
HOSE-END SPRAYER: Easy to Use.

- Make sure knob control is in the Off position. Shake well and attach to hose.
- Turn water on at faucet, aim nozzle toward lawn or desired plant and turn control lever to ON to begin spraying.
- When finished turn control lever to OFF position. Discharge residual water pressure by turning ON and OFF again.
- Apply to the [xxx] square foot area to be sprayed. [Square feet value to be inserted per specific package size].

APPLICATION INSTRUCTIONS



People and pets should not be allowed in treated areas until spray has dried.

Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

Apply sufficient water to obtain coverage without excessive runoff.

49-73

LAWNS

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

General Directions

- 1) Apply after mowing or allow sprayed area to completely dry before mowing.
- 2) Measure the length and width of area to be treated before application. Length x width = square feet. For example, an area 25 ft. long and 20 ft. wide is 500 sq. ft.
- 3) Begin applications when conditions are favorable for disease infection, prior to disease symptoms.
- 4) Can be safely applied after seeding or at seed germination.

Disease	How Much to Apply (in 1 gal. of water per 250 sq. ft. of lawn)	Comments
Brown Patch	½-1 tsp.	First application should be in May or June before disease is present. If needed, repeat treatment every 14-28 days.
Anthracnose Fusarium Patch Gray Leaf Spot Leaf Spot Rhizoctonia Leaf Spot Necrotic Ring Spot Rusts Powdery Mildew Red Thread	½-1 tsp.	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Apply every 14-28 days.
Take-All Patch	½-1 tsp.	Apply twice in the Fall (Sept. and Oct.) and twice in the Spring (April and May). Southern lawns also need to be treated twice in the summer (June-August).
Snowmold (Gray or Pink)	1½ tsp. 1 tsp.	Make a single application in late fall or Apply twice in the fall prior to snow cover waiting 10 -14 days between each application.

50/73

Disease	How Much to Apply (in 1 gal. of water per 250 sq. ft. of lawn)	Comments
Zoysia Patch	½-1 tsp.	Apply 1-2 applications approximately one month prior to zoysiagrass dormancy (14-28 day interval)

51/73

**Azoxystrobin H&G
Fungicide Concentrate**

Active Ingredient:

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy)
pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate^{1*} 22.9%

Inert Ingredients: 71.1%

Total: 100.00%

Contains 2.08 lbs. of active ingredient per gallon

¹CAS No. 131860-33-8

*IUPAC

Product of the United Kingdom

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on label.

EPA Reg. No. 100-
EPA Est.

SCP

Net Contents/Net Weight

Appendix 1 - Consolidated List of Label Claims

- Broad spectrum and systemic disease control on ornamentals, shrubs and trees
- Controls brown patch, powdery mildew, anthracnose, rust, blights. Scab, and more
- Prevents and stops major diseases on
- Concentrate - Mix with water and spray
- For use in pump-up and hose-end sprayers
- Absorbed into the plant - Can't be washed off by rain or sprinklers after spray has dried
- Rainfast shortly after spray has dried
- Not affected by rain after spray has dried
- Systemic action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on ornamentals, shrubs and trees
- Works up to 4 weeks for certain diseases
- Prevents and stops ...
 - Anthracnose Brown Patch
 - Blights Powdery Mildew
 - Rust Leaf Spot
 - and more

53/73

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash thoroughly with soap and water after handling.

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.



People and pets should not be allowed in treated areas until spray has dried.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available; for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

5473

Alternate Warranty Statement

Notice: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The buyer assumes responsibility for lack of performance or safety if not used according to the instructions on this label.

DIRECTIONS FOR USE

For Outdoor Residential Use Only

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR BEST RESULTS READ AND FOLLOW ALL LABEL DIRECTIONS

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-888-8372, day or night.

Pesticide/Container Disposal

If container is partially filled:

Important: It is best to use all of the product in accordance with label directions.

If it is necessary to dispose of unused product, please follow any applicable state or local guidelines or call your local solid waste agency.

Never place unused product down any indoor or outdoor drain.

If Container is Empty:

Do not reuse the container.
Place container in the trash.

GENERAL INFORMATION

This water-based systemic fungicide effectively prevents and controls major diseases on ornamentals, shrubs and trees.

USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.
- **DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.**
- If you need more information about ornamentals, shrubs and tree diseases, contact your local County Extension Service.
- Apply product to the plant until it begins to run off of the leaves.
- Allow spray to completely dry before watering.

MIXING PROCEDURES

Shake well before using.

NOTE: ¼ fl. oz = 1½ teaspoons
 ½ fl. oz. = 1 Tablespoon = 3 teaspoons
 1 fl. oz. = 2 Tablespoons = 6 teaspoons

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides.

Move indoor houseplants outdoors before treating. Allow to dry before handling or moving indoors.

Pump-Up Sprayer (for all uses)

- Fill with ½ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

Concentrate Plus Water Style Hose-End Sprayer (ornamentals, shrubs and trees).

- From the following table, determine the amount of concentrate required.
- Pour concentrate into sprayer jar to the required "fl. oz." level.
- Then add the required amount of water to the "gals." level.
- Replace sprayer top on jar and shake well.
- Attach sprayer to hose and apply.
- For ornamentals, shrubs and trees apply to obtain sufficient spray coverage without excessive runoff.

Dial-Style Hose-End Sprayer (use for application to ornamentals, shrubs and trees).

- Fill the sprayer jar with only enough product to cover the plants or the area to be treated.
- Set Dial on the "fl. oz./gallon" level specified in the following table.
- Close sprayer, attach to hose and apply.
- For ornamentals, shrubs and trees, apply to obtain sufficient spray coverage without excessive runoff.

**(Optional language for product sold in a HOSE-END SPRAYER package)
HOSE-END SPRAYER: Easy to Use. (use for application to ornamentals, shrubs and trees).**

- Make sure knob control is in the Off position. Shake well and attach to hose.
- Turn water on at faucet, aim nozzle toward lawn or desired plant and turn control lever to ON to begin spraying.
- When finished turn control lever to OFF position. Discharge residual water pressure by turning ON and OFF again.
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

APPLICATION INSTRUCTIONS



People and pets should not be allowed in treated areas until spray has dried.

Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

609/13

Apply sufficient water to obtain coverage without excessive runoff.

6/73

TREES, ORNAMENTALS AND SHRUBS

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

General Directions

1. Do not apply this product to apples, crabapples, or flowering cherry trees due to possible phytotoxicity.
2. Apply sufficient water to obtain coverage without excessive runoff.
3. Do not apply more than two applications before alternating with a fungicide that has a different mode of action.

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Roses	Downy Mildew Leaf Spot Powdery Mildew Rusts	1/4-1/2 tsp.	Thorough coverage needed for best results. Begin application at budbreak. Apply every 7-21 days.
Flowers, Annual and Perennial - Bedding Plants	Downy Mildew Leaf Spots Leaf Blights Powdery Mildew Rusts Scab	1/8-1/4 tsp.	Apply every 7-14 days prior to infection. Do not apply the 3 teaspoon rate on less than 14 day spray intervals.
Annual and Perennial Flowers/Mature Plants	Downy Mildew Leaf Spots Leaf Blights Powdery Mildew Rusts Scab	1/4-1/2 tsp.	Apply before disease begins. Apply every 7-28 days.
Dogwoods and Other Landscape (Ornamental) and Shade Trees	Anthrachnose Conifer Blights Downey Mildew Leaf Spot Powdery Mildew Scab	1/2-1 tsp. 1/8-1/2 tsp.	Apply every 7-28 days. Apply every 7-28 days.

62/73

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose	1/2-1 tsp.	Apply every 7-28 days.
	Downy Mildew	1/8-1/2 tsp.	Apply every 7-28 days.
	Leaf Spots		
	Leaf Blights		
	Powdery Mildew		
	Rusts		
Scab			

63/73

**Azoxystrobin H&G
Fungicide Concentrate**

Active Ingredient:

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy) pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate ^{1*}	22.9%
Inert Ingredients:	71.1%
Total:	100.00%

Contains 2.08 lbs. of active ingredient per gallon

¹CAS No. 131860-33-8

*IUPAC

Product of the United Kingdom

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on label.

EPA Reg. No. 100-
EPA Est.

SCP

Net Contents/Net Weight

Appendix 1 - Consolidated List of Label Claims

- Broad spectrum and systemic disease control on vegetables
- Controls brown patch, powdery mildew, anthracnose, rust, blights. Scab, and more
- Prevents and stops major diseases on vegetables
- Concentrate - Mix with water and spray
- For use in pump-up and hose-end sprayers
- Absorbed into the plant - Can't be washed off by rain or sprinklers after spray has dried
- Rainfast shortly after spray has dried
- Not affected by rain after spray has dried
- Systemic action - Can't be washed off by rain after spray has dried
- Works up to 4 weeks for certain diseases
- Prevents and stops ...
 - Anthracnose Brown Patch
 - Blights Powdery Mildew
 - Rust Leaf Spot
 - and more

65/73

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash thoroughly with soap and water after handling.

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.



People and pets should not be allowed in treated areas until spray has dried.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available; for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

66/73

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

The active ingredient, azoxystrobin, in this product can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

67/73

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

68/73

Alternate Warranty Statement

Notice: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The buyer assumes responsibility for lack of performance or safety if not used according to the instructions on this label.

69/73

DIRECTIONS FOR USE

For Outdoor Residential Use Only

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR BEST RESULTS READ AND FOLLOW ALL LABEL DIRECTIONS

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-888-8372, day or night.

Pesticide/Container Disposal

If container is partially filled:

Important: It is best to use all of the product in accordance with label directions.

If it is necessary to dispose of unused product, please follow any applicable state or local guidelines or call your local solid waste agency.

Never place unused product down any indoor or outdoor drain.

If Container is Empty:

Do not reuse the container.
Place container in the trash.

GENERAL INFORMATION

This water-based systemic fungicide effectively prevents and controls major diseases on vegetables.

USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.
- **DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.**
- If you need more information about fruit, vegetable, lawn and ornamentals diseases, contact your local County Extension Service.
- Apply product to the plant until it begins to run off of the leaves.
- Allow spray to completely dry before watering.

MIXING PROCEDURES

Shake well before using.

NOTE: $\frac{1}{4}$ fl. oz = $1\frac{1}{2}$ teaspoons
 $\frac{1}{2}$ fl. oz. = 1 Tablespoon = 3 teaspoons
 1 fl. oz. = 2 Tablespoons = 6 teaspoons

Food utensils such as measuring spoons and cups should not be used for food purposes after use with pesticides.

Move indoor houseplants outdoors before treating. Allow to dry before handling or moving indoors.

Pump-Up Sprayer

- Fill with $\frac{1}{2}$ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

APPLICATION INSTRUCTIONS



People and pets should not be allowed in treated areas until spray has dried.

Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

Apply sufficient water to obtain coverage without excessive runoff.

7/2/13

VEGETABLES

DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

General Directions

- 1) Do not apply more than two applications before alternating with a fungicide that has a different mode of action. Azoxystrobin H&G is a group 11 fungicide. Do not alternate with another group 11 fungicide.
- 2) Do not make more than 6 applications of this product per crop (unless specified in use directions).
- 3) Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait to Harvest	Comments
Asparagus	Stemphyllium purple spot	1½-2 tsp.	100	Apply on a 7-14 day schedule to asparagus foliage after harvest of spears is complete.
Broccoli, Brussel Sprouts Cabbage, Cauliflower, Kohlrabi	Leaf Spot, Downy Mildew	1-1¼ tsp.	0	Apply on a 7-14 day schedule.
Garlic, Leek, Onion (Bulb and Green)	Purple Blotch, Rust And White Rot Botrytis Leaf Blight Downy Mildew	1-1½ tsp. 1¼-2 tsp.	0	Apply on a 7-14 day schedule.
Beet, Broad (Fava) bean, Carrots, Celery, Eggplant, Horseradish, Lettuce (Head and Leaf), Okra, Parsley, Parsnip, Peas, Pepper, Radish, Rhubarb, Rutabaga, Snap bean, Spinach, Turnip, Wax bean	Anthrachnose, Belly Rot, Downy Mildew, Early Blight, Gummy Stem Blight, Leaf Spot, Late Blight, Powdery Mildew, Rust, Southern Blight, White Mold	1¼-2 tsp.	0	Apply on a 7-14 day schedule.
Corn (Sweet and Pop)	Rust, Leaf Blight, Leaf Blight	1-1¼ tsp.	7	Apply on a 7-14 day schedule.
Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon, Zucchini	Anthrachnose, Belly Rot, Downy Mildew, Gummy Stem Blight, Leaf Spot, Powdery Mildew	1½-2 tsp.	1	Make a preventative application on a 5-7 day schedule. For belly rot apply at the 1-3 leaf stage with second application made 10-14 days later.

73/13

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait to Harvest	Comments
Potatoes, Sweet Potato, Yam	Early Blight, Black Dot, Powdery Mildew	1-2 tsp.	14	Begin application when plants are 6 inches high or at prior to first sign of disease. Do not make more than 2 sprays before alternating to a different product with alternate chemistry. Do not make more than 6 applications per season.
	Late Blight (Potato)	2 tsp.		
Tomatoes	Leaf Spot, Powdery Mildew, Rust	¾ tsp.	0	Do not apply until 21 days after transplanting or 35 days after seeding. Apply on a 7-21 day schedule. For late blight apply on a 5-7 day schedule. Make only one application before alternating to another product with a different chemistry.
	Late Blight	1 tsp.		

Syngenta Crop Protection, Inc.
 Greensboro, North Carolina 27419
www.syngenta.com

AZO H&G DRAFT – nsp 09-30-04

000100-RERO.20041001b.Azoxystrobin H&G.doc
 000100-RERO.20041001b.Azoxystrobin H&G.pdf