

100-1242

2/27/2009

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

**OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES**

Ms. Teresa Cox
Regulatory Affairs
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

FEB 27 2009

Dear Ms. Cox:

Subject: Scholar SC
EPA Registration Number 100-1242
Your submission dated October 29, 2008
OPPIN Decision Number 402549

The labeling referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided you make the following changes:

Page 3 - Conditions of Sale and Limitation of Warranty and Liability:

- In the paragraph beginning "The Directions for Use of this product must be followed carefully." change the sentence "Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Syngenta Crop Protection, Inc. or Seller." to "Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Syngenta Crop Protection, Inc. or Seller."
- Throughout the Conditions of Sale and Limitation of Warranty and Liability section change "to the extent permitted by applicable law" to "to the extent consistent with applicable law."

2/17

Please submit one copy of the final printed labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones on my team by phone at (703) 308-9424 or by e-mail at jones.lisa@epa.gov.

Sincerely,



Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped accepted label

3/17

[MASTER]

Scholar® SC

GROUP 12 FUNGICIDE

Fungicide

| | |
|--------------------|--------|
| Active Ingredient: | |
| Fludioxonil:* | 20.4% |
| Other Ingredients: | 79.6% |
| Total: | 100.0% |

*CAS No. 131341-86-1

Scholar SC is a flowable suspension concentrate

Scholar SC contains 1.92 lbs. ai per gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1242

EPA Est.

Product of
Formulated in

SCP 1242A

Net Contents

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
FEB 27 2009**
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

100-1242

| FIRST AID | |
|---|---|
| If in eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If on skin: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not contaminate water when disposing of equipment wash waters or rinsates.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not formulate this product into other End-use products.

GENERAL INFORMATION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL.

GROUP 12 FUNGICIDE

Scholar SC is a protective fungicide used to aid in the control of several post harvest diseases **in post-harvest treatment facilities**. Scholar SC contains fludioxonil that is in the phenylpyrrole class of chemistry and has a unique mode of action, which leads to increased glycerol synthesis [Fungicide Action Group 12]. Fungal isolates with acquired resistance to Group 12 may eventually dominate the fungal population if Group 12 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by fludioxonil or other Group 12 fungicides. A disease management program that includes alternation or tank mixes between Scholar SC and other labeled fungicides that have a different mode of action may prevent pathogen populations from developing resistance. Sanitation and other cultural practices to minimize disease are also recommended to aid in control as well as to assist in preventing/delaying resistance development.

NOTE: Scholar SC may be degraded by exposure to direct sunlight. Treated fruit should not be stored in direct sunlight.

MIXING PROCEDURES

Vigorously shake the product container before mixing. Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use.

To determine the physical compatibility of Scholar SC with other products, use a jar test as described below.

Jar Compatibility Test: Using a quart jar, add the proportionate amounts of the products to 1 qt. of water or wax/oil emulsion. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

If using Scholar SC in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate may be exceeded and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing. Tank mixtures are permitted only in those states where the tank mix partner is registered.

THE CROP SAFETY OF ALL POTENTIAL TANK MIXES INCLUDING ADDITIVES AND OTHER PESTICIDES ON ALL CROPS HAS NOT BEEN TESTED. BEFORE APPLYING ANY TANK MIXTURE, THE SAFETY TO THE TARGET CROP SHOULD BE CONFIRMED.

Add ½ of the required amount of water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion) to the spray or mixing tank. With the agitator running, open the container and add the Scholar SC to the tank. Continue agitation while adding the remainder of the carrier. Begin application of the solution after the Scholar SC has completely and uniformly dispersed into the mix carrier. Maintain agitation until all of the mixture has been applied.

If tank-mixing, add the desired amount of other products recommended for tank mixture after Scholar SC has completely and uniformly dispersed into the mix carrier. In general, tank mix partners should be added in this order: wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied. Maintain agitation until all of the mixture has been applied.

CROP USE DIRECTIONS

Citrus: Calamondin (*Citrus mitis*, *Citrofortunella mitis*), Citrus citron (*Citrus medica*), Citrus hybrids (*Citrus* spp.) (includes chironja, tangelo, tangor), Grapefruit (*Citrus paradisi*), Kumquat (*Fortunella* spp.), Lemon (*Citrus jambhiri*, *Citrus limon*), Lime (*Citrus aurantiifolia*), Mandarin (tangerine) (*Citrus reticulata*), Orange, sour (*Citrus aurantium*), Orange, sweet (*Citrus sinensis*), Pummelo, (*Citrus grandis*, *Citrus maxima*), Satsuma mandarin (*Citrus unshiu*)

Use Scholar SC as a post-harvest dip, drench, flood, or spray for the control of post-harvest diseases caused by:

- Green or Blue mold (*Penicillium* spp.)
- Diplodia stem-end rot (*Lasioidiplodia theobromae*)
- Phomopsis stem-end rot (*Diaporthe citri*)
- Gray mold (*Botrytis cinerea*)

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|---|-------------------------------------|---|
| In-Line Dip/Drench | Green mold Blue mold Diplodia stem-end rot Gray mold | 33-66 fl. oz./100 gals. | <ul style="list-style-type: none"> • Mix 33-66 fl. oz. of Scholar SC in 100 gals. of an appropriate water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion. • Dip for a minimum of 30 seconds and allow fruit to drain. |
| In-line Aqueous or Fruit Coating Spray application | Green mold Blue mold Diplodia stem-end rot Gray mold | 33-66 fl. oz./250,000 lbs. of fruit | <ul style="list-style-type: none"> • Ensure proper coverage of the fruit. • Mix the fungicide solution in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. • Use T-jet, CDA, or similar application system. |
| Do not make more than two applications to citrus fruit. For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing. | | | |
| <ul style="list-style-type: none"> • Ensure the Scholar SC solution remains in suspension by using agitation. • Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks. | | | |

Kiwi

Use Scholar SC as a post-harvest dip or spray for the control of Botrytis fruit rot in kiwi.

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|--------------------|-------------------------------------|---|
| In-Line Dip/Drench | Botrytis fruit rot | 16-32 fl. oz./100 gals. | <ul style="list-style-type: none"> • Mix 16-32 fl. oz. of Scholar SC in 100 gals. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. *Dip for approximately 30 seconds and allow fruit to drain. |
| In-line Aqueous or Fruit Coating Spray application | Botrytis fruit rot | 16-32 fl. oz./200,000 lbs. of fruit | <ul style="list-style-type: none"> • Ensure proper coverage of the fruit. • Mix the fungicide solution in an appropriate amount of water, wax/emulsion, or aqueous dilution of wax/oil emulsion for the crop being treated. |
| Do not make more than one post-harvest application to the fruit. | | | |
| <ul style="list-style-type: none"> • Ensure the Scholar SC solution remains in suspension by using agitation. • Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks. | | | |

Pome fruit: Apple (*Malus domestica*), Crabapple (*Malus spp.*), Loquat (*Eriobotrya japonica*), Mayhaw (*Crataegus aestivalis*, *C. opaca*, and *C. rufula*), Pear (*Pyrus communis*), Pear, oriental (*Pyrus pyrifolia*), Quince (*Cydonia oblonga*)

Use Scholar SC as a post-harvest dip, drench, flood, or spray for the control of post-harvest diseases caused by:

- Blue mold (*Penicillium expansum*)
- Gray mold (*Botrytis cinerea*)
- Bull's-eye rot (*Pezicula malacorticis*)
- Rhizopus rot (*Rhizopus stolonifer*)
- Bitter rot (*Colletotrichum gloeosporioides*)
- Sphaeropsis rot (*Sphaeropsis pyriputrescens*)
- Phacidiopycnis rot (*Phacidiopycnis piri*)
- Phacidiopycnis rot (*Phacidiopycnis washingtonensis*)
- White rot (*Botryosphaeria dothidea*)

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|--|------------------------------------|--|
| Bin/Truck Drench or In-Line Dip/Drench or Flooder | Blue mold Gray mold | 10-16 fl. oz/100 gals. | <ul style="list-style-type: none"> • Ensure proper coverage of the fruit. • For re-cycling in-line drench or dip treatments, the fungicide solution may be prepared in water. • For in-line drench or dip applications, treat fruit for 15-30 seconds and allow fruit to drain. • Fruit coatings may be applied separately after aqueous fungicide treatments. |
| | Rhizopus rot Bull's-eye rot Bitter rot Sphaeropsis rot Phacidiopycnis rot Speck rot White rot | 16 fl. oz./100 gals. | |
| In-line Aqueous or Fruit Coating Spray application | Blue mold Gray mold Rhizopus rot Bull's-eye rot Bitter rot Sphaeropsis rot Phacidiopycnis rot White rot | 16-32 fl oz /200,000 lbs. of fruit | <ul style="list-style-type: none"> • Ensure proper coverage of the fruit. • Mix the fungicide solution in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. • Use T-jet, CDA, or similar application system. |
| <p>Do not make more than two applications to pome fruit. For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing.</p> <ul style="list-style-type: none"> • Ensure the Scholar SC solution remains in suspension by using agitation. • Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfect high-volume, recycling tanks. | | | |

Pomegranates

Use Scholar SC as a post-harvest dip for the control of Botrytis fruit rot in pomegranates:

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|--------------------|----------------------|---|
| In-Line Dip/Drench | Botrytis fruit rot | 32 fl. oz./100 gals. | <ul style="list-style-type: none"> Mix 32 fl. oz. of Scholar SC in 100 gals. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. *Dip for approximately 30 seconds and allow fruit to drain. Dip solution should be replaced with fresh dip solution after 200,000 pounds of fruit has been treated. |
| Do not make more than one post-harvest application to the fruit. | | | |
| <ul style="list-style-type: none"> Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfect high-volume, recycling tanks. | | | |

Stone Fruit: Apricot (*Prunus armeniaca*), Nectarine (*Prunus persica*), Peach (*Prunus persica*), Plum (*Prunus domestica*, *Prunus spp.*), Plum, Chickasaw (*Prunus angustifolia*), Plum, Damson (*Prunus domestica* spp. *insititia*), Plum, Japanese (*Prunus salicina*), Plumcot (*Prunus armeniaca* × *P. domestica*), Prune (fresh), (*Prunus domestica*, *Prunus spp.*), as well as other cultivars and hybrids of these.

Use Scholar SC as a post-harvest dip or spray for the control of post-harvest diseases caused by:

- Brown rot (*Monilinia* spp.)
- Gray mold (*Botrytis cinerea*)
- Rhizopus rot (*Rhizopus stolonifera*)
- Gilbertella rot (*Gilbertella persicaria*)

11/17

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|---|-------------------------------------|---|
| In-Line Dip/Drench | Brown rot Gray mold Rhizopus rot Gilbertella rot | 16 fl. oz./100 gals. | <ul style="list-style-type: none"> Mix 16 fl. oz. of Scholar SC in 100 gals. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain. Dip solution should be replaced with fresh dip solution after 200,000 pounds of fruit has been treated. |
| In-line Aqueous or Fruit Coating Spray application | Brown rot Gray mold Rhizopus rot Gilbertella rot | 16-32 fl. oz./200,000 lbs. of fruit | <ul style="list-style-type: none"> Ensure proper coverage of the fruit. Mix 16-32 fl. oz. of Scholar SC in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-Jet, CDA, or similar application system. For maximum efficacy, use low volume concentrate application systems for treatment of plums. |
| Do not make more than one post-harvest application to the fruit. | | | |
| <ul style="list-style-type: none"> Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfect high-volume, recycling tanks. | | | |

Cherries: Cherry, sweet (*Prunus avium*), Cherry, tart (*Prunus cerasus*), as well as other cultivars and hybrids of these.

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|---|------------------------------------|--|
| In-line Aqueous or flooders application High-Volume (dilute-spray) application | Brown rot Gray mold Rhizopus rot Gilbertella rot | 16-32 fl. oz./50,000 lbs. of fruit | <ul style="list-style-type: none"> Mix 16 fl. oz. of Scholar SC in 50-100 gals. or 32 fl. oz. of Scholar SC in 100 gals. of an appropriate water, wax/emulsion, or aqueous dilution of a wax/oil emulsion. Use flooders, T-jet, or similar application system. |
| Do not make more than one post-harvest application to the fruit. | | | |
| <ul style="list-style-type: none"> Ensure the Scholar SC solution remains in suspension by using agitation. | | | |

True Yam

Use Scholar SC as a post-harvest dip for the control of certain post-harvest rots caused by *Penicillium* and *Fusarium* species.

12/17

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|---|------------------------|--|
| Post Harvest Dip Application | Brown rot Gray mold Rhizopus rot Gilbertella rot | 16-32 fl. oz./100 gals | <ul style="list-style-type: none"> Mix 16-32 fl. oz. of Scholar SC in 100 gals. of an appropriate water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain. |
| Do not make more than one post-harvest application to the tubers. | | | |
| <ul style="list-style-type: none"> Ensure the Scholar SC solution remains in suspension by using agitation. | | | |

NOTE: Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC may be degraded by exposure to direct sunlight. Treated fruit should not be stored in direct sunlight.

Sweet Potato

Use Scholar SC as a post-harvest dip and low volume application for the control of post-harvest rots caused by *Rhizopus stolonifer*.

| Application Method | Disease | Rate (fl. oz.) | Remarks |
|--|--------------|---|---|
| In-Line Dip/Drench | Rhizopus rot | 16-32 fl. oz./100 gals. | <ul style="list-style-type: none"> Mix 16-32 fl. oz. of Scholar SC in 100 gals. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain. Add 8 fl. oz. of Scholar SC to 100 gals. of treating suspension after 500 bushels are treated. After each 1000 bushels treated, drain and flush the tank. Refill with fresh dip suspension. |
| In-line Aqueous or Fruit Coating Spray application | Rhizopus rot | 16 fl. oz./200,000 lbs. of sweet potatoes | <ul style="list-style-type: none"> Ensure proper coverage of the fruit. Mix 16 fl. oz. of Scholar SC in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-Jet, CDA, or similar application system. |
| Do not make more than one post-harvest application to the sweet potatoes. | | | |
| <ul style="list-style-type: none"> Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks. | | | |

NOTE: Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC may be degraded by exposure to direct sunlight. Treated sweet potatoes should not be stored in direct sunlight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Do not store near heat or open flame. Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For Bulk or Minibulk Containers:

Container Disposal: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning.

Container Precautions: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

Refill only with Scholar SC. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than Scholar SC will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

14/17

Scholar®, the Syngenta logo and the CP FRAME  are trademarks of a Syngenta Group Company.

©200X Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1242A

15/17

[BASE/BACK LABEL]

Scholar® SC

GROUP 12 FUNGICIDE

Fungicide

| | |
|--------------------|--------|
| Active Ingredient: | |
| Fludioxonil:* | 20.4% |
| Other Ingredients: | 79.6% |
| Total: | 100.0% |

*CAS No. 131341-86-1

Scholar SC is a flowable suspension concentrate

Scholar SC contains 1.92 lbs. ai per gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in attached booklet.

EPA Reg. No. 100-1242

EPA Est.

Product of
Formulated in

1 gallon
Net Contents

SCP 1242A

| FIRST AID | |
|---|---|
| If in eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If on skin: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not contaminate water when disposing of equipment wash waters or rinsates.

17/
17

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Do not store near heat or open flame. Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Scholar® and the Syngenta logo are trademarks of a Syngenta Group Company.

©200X Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1242A

Scholar SC AMEND-F 10-08 - bb - 2-25-09

000100-01242.20080110F.SCHOLARSC-JAN08-AMEND.pdf