



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

MAR 16 2011

Mr. Pat Dinnen Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, NC 27419

Subject: Removal of Expired Patent Numbers

Dear Mr. Dinnen:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated March 3, 2011 for:

EPA Reg. No 100-1247 EPA Reg. No. 100-1248

Cruiser Maxx Potato Insecticide and Fungicide

The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that the label change(s) requested falls within the scope of PRN-98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov.

Sincerely,

Dani Daniel

Registration Division (7504P)

Insecticide/Rodenticide Branch

March 3, 2011

4. Typed Name

Pat Dinnen

3

NOTIFICATION
JAN 1 6 2011

Certification with Respect to Label Integrity

Version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration # Date Submitted Electronic file name to EPA		
100-1248	3/3/11	000100-01248.20110303.PATENT_NOTIF_MAR2011.pdf

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Pat clinnen	March 3, 2011
Signature	Date
Pat Dinnen Name (typed)	
Label Group Leader	
Title	



NOTIFICATION

JAN 1 7 2011

(booklet)

Note: Do not use, sell or distribute this product within, or into, Nassau County or Suffolk County, New York.

CruiserMaxx™ Potato Insecticide and Fungicide

A seed treatment product for protection against listed insects and diseases in potato tubers.

Active Ingredients:		
Thiamethoxam*	28.00%	
Fludioxonil**	7.00%	
Other Ingredients:	65.00%	
Total:	100.00%	

^{*}CAS No. 153719-23-4 **CAS No. 131341-86-1

One gallon of CruiserMaxx Potato Insecticide and Fungicide contains 2.85 lbs. thiamethoxam, and 0.73 lbs. of fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.


EPA Reg. No. 100-1248 EPA Est. XXXXX

SCP 1248A-M(6-1-09) [1 gal., 2.5 gal.]

SCP 1248A-ML(6-1-09) [15 gal., 120 gal.]

1 gallon 2.5 gallons 15 gallons ___ gallons

Net Contents



	FIRST AID
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	

For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call

1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling, and before a cating, drinking, chewing gum or using tobacco.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride [PVC] ≥ 14 mils or Viton® ≥ 14 mils, section category C)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restrictedentry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves (such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride [PVC] ≥ 14 mils, or Viton® ≥ 14 mils, section category C)
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred, bruised, or damaged seed or seed pieces, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that

planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

Follow recommended Syngenta minimum germination guidelines and seed carryover recommendations when treating seed lots.

USE INFORMATION

CruiserMaxx Potato Insecticide and Fungicide seed treatment contains the active ingredients of thiamethoxam insecticide and fludioxonil fungicide.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neoticotinoid class of chemistry. Thiamethoxam protects against certain chewing and sucking insects through contact and ingestion. These insects include: green peach aphids, Colorado potato beetles, flea beetles, leafhoppers, leaf miners, psyllids and whiteflies. When rate ranges are given, use higher rates when insect pressure is expected to be high.

Fludioxonil protects against damage from certain soil-borne and seed-borne diseases of crop plants. Fludioxonil is active against *Fusarium* dry rot seed decay, seed-borne *Rhizoctonia* that causes stem canker and tuber black scurf and seed-borne *Helminthosporium solani*, the causal agent of silver scurf diseases on potato tubers.

CruiserMaxx Potato Insecticide and Fungicide does not control bacterial disease or diseases present within the seed.

CruiserMaxx Potato Insecticide and Fungicide is to be used as an integral part of a potato pest management strategy. This strategy includes the use of high quality certified seed, proper crop rotation, insect population thresholds, appropriate control measures, optimal harvest time for tubers and proper handling of tubers without bruising. Consult your local agricultural extension agent for more detailed information on insect management practices.

The expected length of protection against the labeled pests is dependent upon the rate used, accurate application of the products to ensure the seed tubers receive the target rate of the active ingredients. Consult your local Syngenta representative or dealer for information relative to your area.

SEED BAG LABEL REQUIREMENTS

Federal law requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and fludioxonil fungicide.
- Do not use for feed, food, or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with CruiserMaxx Potato Insecticide and Fungicide:

- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- In the event of a crop failure or harvest of a crop grown from CruiserMaxx Potato Insecticide and Fungicide treated seed, the field may be replanted immediately to barley, canola, corn, cotton, cucurbit vegetables, fruiting vegetables, legume vegetables (including soybeans), mint (peppermint and spearmint), oil seed crops (rapeseed, Indian rapeseed, Indian mustard seed, field mustard seed, black mustard seed, flax seed, safflower seed, crambe seed and borage seed), root vegetables (except sugar beets), sorghum, strawberry, sunflowers, tobacco, tuberous and corm vegetables and wheat. For any other crop, the minimum plantback interval is 120 days from the date CruiserMaxx Potato Insecticide and Fungicide treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- Do not allow children, pets, or livestock to have access to treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Do not use at a rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season.
- Treated seed must be planted into the soil at a depth greater than 1 inch.

 Do not make any soil or foliar application of products containing thiamethoxam to potatoes grown from seed treated with CruiserMaxx Potato Insecticide and Fungicide.

MIXING PROCEDURES

Important: Always re-circulate CruiserMaxx Potato Insecticide and Fungicide thoroughly before using.

Apply CruiserMaxx Potato Insecticide and Fungicide seed treatment using only Syngenta approved equipment that is designed to apply liquid seed treatments to potatoes. Follow the equipment manufacturers' instructions for set-up and calibration.

CruiserMaxx Potato Insecticide and Fungicide will require dilution prior to atomization and application to potatoes (see equipment use instructions).

Under certain disease conditions, additional amounts (see **CruiserMaxx Potato Insecticide and Fungicide Rate Table**) of Maxim 4FS fungicide may be mixed with the CruiserMaxx Potato Insecticide and Fungicide seed treatment and applied through a calibrated liquid seed treater.

Thoroughly mix the specified amount of CruiserMaxx Potato Insecticide and Fungicide and any additional Maxim 4FS into the required of water for the dilution rate required.

Other tank mix partners may be used with CruiserMaxx Potato Insecticide and Fungicide; however, the user must consider the use rate, formulation, seed and crop safety factors and compatibility of each product to be mixed when determining total application volume.

The total quantity of water and product volume must be adjusted based upon the amount of seed moved under the hood. It is mandatory that the equipment be calibrated to deliver a maximum of 4 fluid ounces of mixture per 100 pounds of seed consistently. Applying excess moisture may predispose the seed to rotting, resulting in poor emergence and stand.

APPLICATION PROCEDURES

Note: If inert dust (fir bark, talc, etc.) or a dust-based fungicide is to be applied, apply the CruiserMaxx Potato Insecticide and Fungicide seed treatment before applying the dust.

Registered dust based fungicides can be applied as a supplemental treatment after the CruiserMaxx Potato Insecticide and Fungicide application. Follow label instructions for these products and ensure that the maximum allowable rates for an active ingredient are not exceeded.

Apply CruiserMaxx Potato Insecticide and Fungicide seed treatment only in well ventilated areas.

Ensure that spray nozzles are properly hooded and shielded to prevent spray from moving off target.

Apply the mixture as a fine spray over the cut or whole seed tubers.

CROP USE DIRECTIONS

CRUISERMAXX POTATO INSECTICIDE AND FUNGICIDE RATE TABLE

Seeding Rate 100 lbs./A	Fluid Ounces of CruiserMaxx Potato Insecticide and Fungicide per 100 lbs. ¹ of Potato Seed Tubers	Volume of Maxim 4FS Needed to Attain 2.5 Grams a.i. (Fludioxonil) per 100 kg.
Less than 22	0.22 – 0.27 fl. oz.	0.03 – 0.04 fl. oz.
22 – 25	0.19 – 0.23 fl. oz.	0.04 – 0.05 fl. oz
26 – 30	0.19 fl. oz.	0.05 fl. oz.

¹ Rates shown are for standard 2.5 ounce seed pieces sizes.

Note: for all other seed sizes or seeding rate make sure you do not apply more than a total or 0.125 lbs. active ingredient (thiamethoxam)/A.

Treated Seed Storage

If the treated tubers need to be stored or held for a few days, make sure that there is adequate cool air (60 degrees Fahrenheit) movement through the pile of cut seed potatoes at a relative humidity of 85 – 90%. Cut and treated seed should not be piled above 6 feet in height. Best results are obtained if potatoes are planted immediately after treatment with CruiserMaxx Potato Insecticide and Fungicide. When transporting cut and treated seed, make sure the seed is covered.

ROTATIONAL RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from CruiserMaxx Potato Insecticide and Fungicide treated seed, the field may be replanted immediately to barley, canola, corn, cotton, cucurbit vegetables, fruiting vegetables, legume vegetables (including soybeans), mint (peppermint and spearmint), oil seed crops (rapeseed, Indian rapeseed, Indian mustard seed, field mustard seed, black mustard seed, flax seed, safflower seed, crambe seed and borage seed), root vegetables (except sugar beets), sorghum, strawberry, sunflowers, tobacco, tuberous and corm vegetables and wheat. For any other crop, the minimum plantback interval is 120 days from the date

CruiserMaxx Potato Insecticide and Fungicide treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (<5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling (> 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times.

Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CruiserMaxx™, Cruiser®, Maxim® and the Syngenta logo are trademarks of a Syngenta Group Company

Viton® is a registered trademark of E.I. DuPont de Nemours and Company

©2009 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1248A-M(6-1-09) [1 gal., 2.5 gal.] SCP 1248A-M(6-1-09) [15 gal., 120 gal.]

CRUMaxxPotato1248A-M(6-1-09) PATENT NOTIF MAR2011 CLEAN – pl – 3/2/11 **000100-01248.20110303.PATENT_NOTIF_MAR2011.pdf**

(Non-detachable Container Label – 1 gal., 2.5 gal.)

Note: Do not use, sell or distribute this product within, or into, Nassau County or Suffolk County, New York.

CruiserMaxx™ Potato Insecticide and Fungicide

A seed treatment product for protection against listed insects and diseases in potato tubers.

Active Ingredients:

Thiamethoxam*	28.00%
	7.00%
Other Ingredients:	65.00%
Total:	100.00%

^{*}CAS No. 153719-23-4

One gallon of CruiserMaxx Potato Insecticide and Fungicide contains 2.85 lbs. thiamethoxam, and 0.73 lbs. of fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1248

EPA Est. XXXXX

SCP 1248A-M(6-1-09) (1 gal., 2.5 gal.)

1 gallon

2.5 gallons

Net Contents

^{**}CAS No. 131341-86-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum or using tobacco.

	FIRST AID
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

Environmental Hazards

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CruiserMaxx™, the Syngenta logo and the CP FRAME are trademarks of a Syngenta Group Company

©2009 Syngenta

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1248A-M(6-1-09) [1 gal., 2.5 gal.]

CRUMaxxPotato1248A-M(6-1-09) PATENT NOTIF MAR2011 CLEAN - pI - 3/2/11 000100-01248.20110303.PATENT_NOTIF_MAR2011.pdf

(Non-detachable Container Label -15 gal. and 120 gal.)

Note: Do not use, sell or distribute this product within, or into, Nassau County or Suffolk County, New York.

CruiserMaxx™ Potato Insecticide and Fungicide

A seed treatment product for protection against listed insects and diseases in potato tubers

Active Ingredients:

Thiamethoxam*	28.00%
Fludioxonil**	7.00%
Other Ingredients:	65.00%
Total:	100.00%

^{*}CAS No. 153719-23-4

One gallon of CruiserMaxx Potato Insecticide and Fungicide contains 2.85 lbs. thiamethoxam, and 0.73 lbs. of fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1248

EPA Est. XXXXX

SCP 1248A-M(6-1-09) [15 gal.]

SCP 1248A-M(6-1-09) [120 gal.]

Product ID XXXXX

15 gallons

120 gallons

Net Contents

^{**}CAS No. 131341-86-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

control center or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	FIRST AID		
Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	If swallowed	 treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious 	
 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) 		 Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment 	
 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) 	If inhaled	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further 	
or doctor, or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	If in eyes	 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment 	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	Have the product container or label with you when calling a poison control center		
For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)			
Call 1-800-888-8372			

Environmental Hazards

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2)Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (> 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CruiserMaxx™, and the Syngenta logo are trademarks of a Syngenta Group Company

©2009Syngenta

Manufactured for : Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1248A-M(6-1-09) [15 gal.] SCP 1248A-M(6-1-09) [120 gal.]

 $\label{eq:crumax} \mbox{CRUMaxxPotato1248A-M(6-1-09) PATENT NOTIF MAR2011 CLEAN - pl - 3/2/11 \mbox{\bf 000100-01248.20110303.PATENT_NOTIF_MAR2011.pdf}$