



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 15, 2018

Adora Clark
Federal Team Lead, Fungicides
Syngenta Crop Protection, LLC
PO Box 18300
Greensboro, NC 27419

Subject: PRIA Label Amendment – New uses of difenoconazole on cranberry, guava, and papaya; expansion from grape use to crop subgroup 13-07F; and conversion of foliar uses on crop subgroups 5A and 5B into crop groups 5-16, 4-16B, and kohlrabi (previously on labels under 5A).
Product Name: Inspire
EPA Registration Number: 100-1262; Petition 6E8484
Application Date: 6/6/2016
Decision Number: 518690, 518692; 518702, 518703

Dear Ms. Clark:

The application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable under FIFRA Section 3(c)(7)(B), subject to the following conditions:

1. You must submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.
2. You are required to comply with the data requirements described in the DCI identified below:
 - a. Difenoconazole GDCI-128847-1602 at <https://www.regulations.gov/document?D=EPA-HQ-OPP-2015-0401-0016>

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:
<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product

if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). If you have any questions, please contact Lindsay Roe by phone at 703-347-0506, or via email at roe.lindsay@epa.gov.

Sincerely,



Cynthia Giles-Parker, Chief
Fungicide Branch
Registration Division (7505P)

Enclosure -stamped “accepted” labels

ACCEPTED

02/15/2018

SUPPLEMENTAL LABELING

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 100-1262

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP

DIFENOCONAZOLE GROUP 3 FUNGICIDE

Inspire®

Fungicide

This supplemental label expires on 02/16/2021 and must not be used or distributed after this date.

Active Ingredient:	
Difenoconazole*:	23.2%
Other Ingredients**:	76.8%
Total:	100.0%

*CAS No. 119446-68-3

**Contains petroleum distillates

Inspire is formulated as an emulsifiable concentrate (EC) containing 2.08 lb ai/gal of difenoconazole.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use on container label.
See First Aid statement on container label.

EPA Reg. No. 100-1262

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Inspire as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Cranberry	Bitter rot <i>(Colletotrichum gloeosporioides)</i> Blotch rot <i>(Phylospora vacciniae)</i> Cottonball <i>(Monilinia oxycocci)</i> Leaf rust <i>(Pucciniastrum vaccinii)</i> Ripe rot <i>(Coleophoma empetri)</i>	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 21 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 3) Do not allow release of irrigation or flood water for at least 7 days after the last application.
- 4) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 5) Do not treat fields used for aquaculture of fish or crustacean.
- 6) Do not drain water from treated fields into ponds used for aquaculture of fish or crustacean.
- 7) Do not use water drained from treated field to irrigate other crops.
- 8) Do not apply Inspire within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Guava	Alternaria Fruit Rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum gloeosporioides)</i> Suppression Rust <i>(Puccinia psidii)</i>	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied the day of harvest. (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Papaya	Blossom blight and fruit rot (<i>Colletotrichum gloeosporioides</i>) Alternaria fruit spot (<i>A. alternata</i>) Powdery Mildew (<i>Erysiphe</i> spp.) Brown Spot (<i>Corynespora cassicola</i>)	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied the day of harvest. (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Watercress	Cercospora leafspot (<i>Cercospora</i> spp.)	7.0 (0.114)	<p>For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application.</p> <p>Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply directly to water and do not allow water in a treated field for at least 24 hours.
- 2) Do not apply more than 28 fl oz/A/year of Inspire.
- 3) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 4) Do not apply Inspire within 30 days of harvest (30-day PHI).

©20XX Syngenta

Inspire® and the Syngenta logo are trademarks of a Syngenta Group Company

[MASTER LABEL]

DIFENOCONAZOLE **GROUP 3** **FUNGICIDE**

Inspire®

Fungicide

Active Ingredient:	
Difenoconazole*:	23.2%
Other Ingredients**:	76.8%
Total:	100.0%

*CAS No. 119446-68-3

**Contains petroleum distillates

Inspire is formulated as an emulsifiable concentrate (EC) containing 2.08 lb ai/gal of difenoconazole.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.
See First Aid statement inside booklet or on container label.

EPA Reg. No. 100-1262

EPA Est.

Net Contents



FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN	
Contains petroleum distillates - vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and all other handlers must wear:

- Chemical-resistant gloves such as barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or Viton® \geq 14 mils
- Protective eyewear
- Long-sleeved shirt and long pants
- Socks and shoes

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic **estuarine/marine** organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this

product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product may contaminate water through drift or spray in wind. This product has a potential for runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Chemical-resistant gloves such as barrier laminate, butyl rubber greater than or equal to 14 mils, nitrile rubber greater than 14 mils, or Viton greater than 14 mils
- Protective eyewear
- Coveralls
- Socks and shoes

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

Inspire is a broad spectrum fungicide with systemic and curative properties recommended for the control of many important plant diseases. Inspire may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Inspire has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. Under high disease pressure, it is recommended to use the highest rate and shortest interval.

Rotational Crops: Please see the table below for the crop rotational restrictions.

Rotational Crop	Planting Time From Last Inspire Application
Artichoke, Globe	0 days
Bean and Pea, Dried Shelled Subgroup 6C	
Berry, Bushberry Subgroup 13-07B	
Berry, Low Growing Subgroup 13-07G	
Brassica (Cole) Head and Stem vegetables Group 5-16	
Brassica (Cole) Leafy greens Subgroup 4-16B	
Bulb vegetables, bulb onion Subgroup 3-07A and green onion Subgroup 3-07B	
Carrots	
Chickpeas	
Citrus fruit Crop Group 10-10	
Cotton Subgroup 20C	
Cucurbit vegetables Crop Group 9	
Fruit, small, vine climbing Subgroup 13-07F, except fuzzy kiwifruit	
Fruiting vegetables Crop Group 8-10	
Ginseng	
Guava	
Kohlrabi	
Papaya	
Pome fruit Crop Group 11-10	
Potatoes	
Rapeseed Subgroup 20A	
Rice	
Soybeans	
Stone fruit Crop Group 12-12	
Strawberry	
Sugar beets	
Tomatoes and tomatillos	
Tree nuts Crop Group 14-12	
Tuberous and corm vegetable subgroup 1C	
Watercress	
Wild rice	
Cereals (wheat, barley, triticale, oats and rye) Root and Tuber Vegetables, Crop Group 1 (except Carrot, Sugar Beet, and Tuberous Corm Vegetable Subgroup 1C)	30 days
All other crops intended for food and feed	60 days

Crop Tolerance: Plant tolerance has been found acceptable for all crops on the label; however, not all possible tank-mix combinations have been tested under all conditions.

When possible, it is recommended to test the combinations on a small portion of the crop to ensure a phytotoxic response will not occur as a result of application.

Spray Drift Management: To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER. More information on managing spray drift can be found on the Syngenta Crop Protection website (<http://www.syngenta-us.com/practicing-stewardship/responsible-pesticide-application>).

IPM: Inspire should be integrated into an overall disease and pest management strategy (IPM) whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. Inspire may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
----------------	-------	----------	-----------

For resistance management, Inspire contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Inspire and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Do not apply to plants grown for transplanting purposes.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Inspire or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.

- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Syngenta Crop Protection at 1-866-796-4368. You can also contact your pesticide distributor or university extension specialist to report resistance.

SPRAY EQUIPMENT

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Use the same size nozzles uniformly spaced across the boom.
- Calibrate sprayer before use.
- Use screens to protect the pump and to prevent nozzles from clogging.
- On suction side of pump use screens that are *16-mesh or coarser*.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - (1) Maintain sufficient pressure at the nozzle tip to give the required flow rate and droplet size to provide acceptable coverage of the target crop.
 - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

MIXING INSTRUCTIONS

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Inspire Alone (no tank mix):

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.

- With the agitator running, add Inspire to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Inspire has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Inspire + Tank Mixtures: Inspire is usually compatible with tank-mix partners. To determine the physical compatibility of Inspire with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Inspire to the spray tank.
- Allow Inspire to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.
- Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product must not be mixed with any product which prohibits such mixing.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture

APPLICATION PROCEDURES

Ground Application:

- Apply in a minimum of 10 gal of water per acre, unless specified otherwise.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- Apply in a minimum of 5 gal of water per acre, unless specified otherwise. For pome fruit apply in a minimum of 10 gal of water per acre.

- Do not apply under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.
- Do not apply directly to humans or animals.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125-0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Note: Do not inject Inspire at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part Inspire. Inspire is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating Inspire through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply $\frac{1}{8}$ - $\frac{1}{2}$ inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying Inspire through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Inspire required to treat the area covered by the irrigation system.
- Add the required amount of Inspire and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the Inspire solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the Inspire solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying Inspire through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of Inspire required to treat the area covered by the irrigation system.

- Add the required amount of Inspire into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Inspire solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC USE DIRECTIONS

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Almonds	Alternaria leafspot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum. acutatum</i>) Blossom Blight Brown rot (<i>Monilinia</i> spp.) Brown rot/hull rot (<i>Monilinia</i> spp.) Powdery mildew (<i>Podosphaera tridactyla</i> , <i>Sphaerotheca pannosa</i>) Scab (<i>Venturia carpophilia</i>) Shot hole (<i>Wilsonmyces carpophilus</i>)	5 - 7 (0.08 - 0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>If monitoring or history indicates the presence of Alternaria, apply 7 fl oz/A of Inspire in the late spring (mid-April to beginning of May) and then repeat the treatment 2-3 weeks later.</p> <p>For blossom blight, begin applications at early bloom and continue through petal fall.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) Do not apply more than 28 fl oz/A/year of Inspire. 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) Do not apply within 14 days of harvest (14-day PHI). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Artichoke, Globe	Ramularia leaf spot Ramularia bud spot (<i>R. cynarae</i>)	7 (0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For ground applications, apply in 50-200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply Inspire within 3 days of harvest (3-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Bean and Pea, Dried Shelled (except soybean) Subgroup 6C</p> <p>To be grown for bean, dried seed only.</p> <p>Phaseolus Vigna Pisum Lupinus</p> <p>See complete list below</p> <p>See specific Directions for soybeans and chickpea</p>	<p>Anthranose (<i>Colletotrichum lindemuthianum</i>) Alternaria leaf spot (<i>A. alternata</i>) Alternaria blight (<i>Alternaria</i> spp.) Ascochyta leaf and pod spot (<i>Ascochyta</i> spp.) Ascochyta blight (<i>Mycosphaerella pinodes</i>) Cercospora leaf spot (<i>Cercospora cruenta</i>)</p>	<p>7 (0.114)</p>	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Complete List of Bean and Pea, dried shelled (except soybean) – Subgroup 6C: Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean; guar; lablab bean; lentil; pea (*Pisum*) (includes field pea); pigeon pea

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 14 fl oz/A/year of Inspire (0.23 lb ai/A/year) for pea vines and hay.
- 3) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 4) Do not feed or harvest cowpeas forage and hay.
- 5) Do not apply Inspire within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Berry Bushberry Subgroup 13-07B Blueberry	Powdery mildew <i>(Microsphaera alni)</i> Anthracnose <i>(Colletotrichum spp.)</i> Septoria leaf spot <i>(S. albopunctata)</i> Alternaria leaf spot <i>(A. tenuissima)</i> Leaf rust <i>(Pucciniastrum vaccinii)</i> Monilinia blight and Mummyberry blight <i>(M. vaccinii-corymbosis)</i>	7 (0.114)	Begin applications prior to disease onset when conditions are conducive for disease. For Monilinia and mummyberry, apply at or near flower bud swell and again at leaf bud swelling. For other diseases, apply during early bloom. Apply Inspire on a 7- to 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.

Complete List of Bushberry Subgroup: Aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry; native currant; salal; sea buckthorn; cultivars, varieties, and/or hybrids of these.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply Inspire within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Berry, Low Growing Subgroup 13-07G Strawberry See additional crops below. See separate instructions for cranberry.	Anthracnose (<i>Colletotrichum acutatum</i>) Leaf spot (<i>Cercospora fragariae</i>) Powdery mildew (<i>Sphaerotheca macularis</i>) Leaf rust (<i>Phragmidium potentillae</i>)	5.5-7 (0.09-0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Additional Low Growing Berries: Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry, Strawberry including all cultivars and/or hybrids of these.			
Specific Use Restrictions: <ol style="list-style-type: none"> 1) Do not apply more than 28 fl oz/A/year of Inspire. 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) Inspire may be applied up to the day of harvest (0-day PHI). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Brassica Head and Stem Vegetable Crop group 5-16</p> <p>Broccoli Brussels sprouts Cabbage Cauliflower</p> <p>See additional crops below.</p> <p>Including all cultivars and/or hybrids of these.</p>	<p>Alternaria diseases (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum higginsianum</i>) Cercospora leafspot (<i>C. brassicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)</p>	<p>5 - 7 (0.08 - 0.114)</p>	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			

Complete list of Brassica Head and Stem Vegetables Crop group 5-16 including: Broccoli; Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; cultivars, varieties, and hybrids of these commodities

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 1 day of harvest (1-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Brassica Leafy Greens Subgroup 4-16B Arugula Chinese cabbage Bok Choy Collards Kale Mustard greens Turnip greens See additional crops below	Alternaria diseases (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum higginsianum</i>) Cercospora leafspot (<i>C. brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)	5-7 (0.08-0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
See separate instructions for watercress.	Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.		
Complete list of Brassica Leafy Greens Vegetable subgroup 4-16B including: Arugula; broccoli raab; broccoli, Chinese; cabbage, abyssinian; cabbage, seakale; cabbage, Chinese, bok choy; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.			
Specific Use Restrictions: 1) Do not apply more than 28 fl oz/A/year of Inspire. 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) Do not apply within 1 day of harvest (1-day PHI).			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Bulb Vegetables Bulb Onion Subgroup 3-07A Onions, bulb Garlic Shallot Green Onion Subgroup 3-07B Onions, green Leek Welsh onion tops	Cercospora leafspot <i>(C. duddiae)</i> Leaf blotch <i>(Cladosporium allii-cepae)</i> Powdery Mildew <i>(Leveillula taurica)</i> Purple blotch <i>(Alternaria porri)</i> Stemphyllium leaf blight <i>(S. vesicarium)</i> Cladosporium leaf blotch <i>(C. allii)</i> Rust <i>(Puccinia allii)</i>	5 - 7 (0.08 - 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Bulb onion subgroup 3-07A: Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.</p>			
<p>Green onion subgroup 3-07B: Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) For green onions, do not apply more than 21 fl oz/A/year of Inspire. 2) For dry bulb onions, do not apply more than 28 fl oz/A/year of Inspire. 3) Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products to green onions. 4) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products to dry bulb onions. 5) Do not apply within 7 days of harvest (7-day PHI). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Carrots	<p>Cercospora leaf spot (<i>Cercospora carotae</i>)</p> <p>Alternaria leaf blight (<i>Alternaria dauci</i>)</p> <p>Powdery mildew (<i>Erysiphe</i> spp.)</p>	5.5-7 (0.09-0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Chickpea	Alternaria blight (<i>A. alternata</i>) Ascochyta blight (<i>A. rabiei</i>) Powdery mildew (<i>Leveillula taurica</i>) Rust (<i>Uromyces ciceris-arietini</i>)	5.5-7 (0.09-0.114)	Begin applications prior to disease onset when conditions are conducive for disease. The minimum retreatment interval is 14 days. Make Inspire applications at least 14 days apart, making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Citrus Fruit Crop Group 10- 10</p> <p>Grapefruit Lemon Lime Orange (sour and sweet) Tangerine</p> <p>Including all cultivars and/or hybrids of these</p> <p>See additional crops in this subgroup below.</p>	<p>Greasy spot (<i>Mycosphaerella citri</i>) Melanose (<i>Diaporthe citri</i>) Scab (<i>Elsinoe fawcettii</i>) Sweet orange scab (<i>Elsinoe australis</i>) Albinism (<i>Alternaria alternata pv citri</i>) Post bloom fruit drop (PFD) (<i>Colletotrichum acutatum</i>) Alternaria leaf and fruit spot (<i>Alternaria citri</i>) Diplodia stem-end rot (<i>Diplodia natalensis</i>) Phomopsis stem- end rot (<i>Phomopsis citrii</i>) Anthracnose (<i>Colletotrichum spp.</i>) Black spot (<i>Guignardia citricarpa</i>)</p>	<p>5 - 7.7 (0.08 - 0.125)</p>	<p>Inspire applications should begin prior to disease development and continue throughout the season on 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			

Citrus Fruit subgroup: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citrus citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangor; trifoliolate orange; unqi fruit; cultivars, varieties, and/or hybrids of these.

Specific Use Restrictions:

- 1) Do not apply more than 28.3 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.5 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied up to the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Cotton (Subgroup 20C)	<p>Alternaria leafspot (<i>Alternaria</i> spp.)</p> <p>Areolate mildew (<i>Ramularia gossypii</i>)</p> <p>Leafspots and blights (<i>Alternaria</i> spp., <i>Ascochyta gossypii</i>, <i>Cercospora</i> spp., <i>Stemphyllium</i> spp.)</p> <p>Target spot (<i>Corynespora</i> <i>cassicola</i>)</p>	5.0 - 7.0 (0.08 - 0.114)	<p>For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates.</p> <p>For foliar disease control, the first application should be targeted approximately at pin-head square to first bloom or when conditions are conducive for disease development. For best control of target spot, adjust the GPA to ensure coverage of upper and lower leaves. Subsequent applications may be made on a 14-21 day interval.</p> <p>Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 21 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply Inspire within 45 days of harvest (45-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Cranberry	Bitter rot <i>(Colletotrichum gloeosporioides)</i> Blotch rot <i>(Physalospora vaccinia)</i> Cottonball <i>(Monilinia oxycocci)</i> Leaf rust <i>(Pucciniastrum vaccinii)</i> Ripe rot <i>(Coleophoma empetri)</i>	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 21 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 3) Do not allow release of irrigation or flood water for at least 7 days after the last application.
- 4) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 5) Do not treat fields used for aquaculture of fish or crustacean.
- 6) Do not drain water from treated fields into ponds used for aquaculture of fish or crustacean.
- 7) Do not use water drained from treated field to irrigate other crops.
- 8) Do not apply Inspire within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Filberts (Hazelnuts)	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	5 - 7 (0.08 - 0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Fruiting Vegetables Crop Group 8-10 Eggplant Groundcherry Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo Tomato See additional crops in this subgroup below.	Early blight <i>(Alternaria solani)</i> Black mold <i>(A. alternata)</i> Gray leafspot <i>(Stemphylium botryosum)</i> Powdery mildew <i>(Leveillula taurica)</i> Septoria leafspot <i>(S. lycopersici)</i> Target spot <i>(Corynespora cassiicola)</i> Anthracnose <i>(Colletotrichum spp.)</i> Leaf mold <i>(Fulvia fulva)</i>	4-7 (0.065-0.114)	Begin applications at first sign of disease or when conditions are conducive for disease development. Apply Inspire on a 7- to 14-day schedule. Inspire can be used in blocking program using a maximum of two consecutive applications before rotating to fungicides with another mode of action that are registered for these diseases. If disease pressure is high, use the highest rate and shortest interval. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Fruiting Vegetables subgroup: African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; pepper, bell; pepper, nonbell; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.			
Specific Use Restrictions: 1) Do not apply more than 28 fl oz/A/year of Inspire. 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) Inspire may be applied up to the day of harvest (0-day PHI).			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Ginseng	Alternaria blight (<i>A. panax</i>) Powdery mildew (<i>Erysiphe</i> spp.)	7 (0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. Use a minimum of 15 gal/A for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied up to the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Grapes (except Concord, Concord Seedless, and Thomcord. See Precaution under Remarks)</p> <p>(Fruit, small, vine climbing, except fuzzy kiwifruit – subgroup 13-07F)</p> <p>See additional crops in this subgroup below.</p>	<p>Alternaria rot (<i>A. alternata</i>) Powdery mildew (<i>Uncinula necator</i>) Rotbrenner (<i>Pseudopezicula tracheiphila</i>) Septoria leaf spot (<i>S. ampelina</i>) Black Rot (<i>Guignardia bidwellii</i>) Angular leafspot (<i>Mycosphearella angulata</i>) Anthracnose (<i>Elsinoe ampelina</i>) Leaf Blight (<i>Pseudocercospora vitis</i>)</p>	<p>5 - 7 (0.08 - 0.114)</p>	<p>For powdery mildew, begin at bud break and apply on a 10- to 21-day interval, making no more than 2 sequential applications before alternating to a fungicide with a different mode of action.</p> <p>For Black rot - begin when shoot length is 1-3 inches and continue on a 10-day interval making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>For all other diseases, begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 10- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p> <p>PRECAUTION: On <i>V. labrusca</i>, <i>V. labrusca</i> hybrids and other non-vinifera hybrids where sensitivity is not known, the use of Inspire by itself or in tank mixtures with materials that may increase uptake (adjuvants, foliar fertilizers) may result in leaf burning or other phytotoxic effects.</p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>

Complete list of small fruit vine climbing, except fuzzy kiwifruit, subgroup 13-07F: Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Guava	Alternaria Fruit Rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum gloeosporioides)</i> Suppression Rust <i>(Puccinia psidii)</i>	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied the day of harvest. (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Kohlrabi	Alternaria diseases (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum higginsianum</i>) Cercospora leafspot (<i>C. brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)	5-7 (0.08-0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 1 day of harvest (1-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Papaya	Blossom blight and fruit rot <i>(Colletotrichum gloeosporioides)</i> Alternaria fruit spot <i>(A. alternata)</i> Powdery Mildew <i>(Erysiphe spp.)</i> Brown Spot <i>(Corynespora cassicola)</i>	7 (0.114)	For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied the day of harvest. (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Pecans	Pecan Scab <i>(Cladosporium caryigenum)</i> Downy Spot <i>(Mycosphaerella caryigena)</i> Liver Spot <i>(Gnomonia caryae)</i> Vein Spot <i>(Gnomonia nerviseda)</i> Zonate Leaf Spot <i>(Grovesinia pyramidalis)</i> Powdery Mildew <i>(Microsphaera penicillata)</i>	5 - 7 (0.08 - 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Pistachios	Panicle and Shoot Blight (<i>Botryosphaeria dothidea</i>) Alternaria late blight (<i>Alternaria</i> spp.) Septoria leaf spot (<i>S. pistaciarum</i>)	5 - 7 (0.08 - 0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>For <i>Botryosphaeria</i>, begin applications when green leaf tissue becomes visible and continue on a 14- to 21-day interval. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Pome Fruit Crop Group 11-10 Apple Crabapple Loquat Mayhaw Pear Pear, Oriental Quince</p> <p>See additional crops in this subgroup below.</p>	<p>Alternaria blotch (<i>Alternaria</i> spp.) Brooks fruit spot (<i>Mycosphaerella pomi</i>) Cedar apple rust (<i>Gymnosporangium juniperi-virginianae</i>) Flyspeck (<i>Zygophiala jamacaicensis</i> (Formerly known as <i>Schizothyrium pomi</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) Quince rust (<i>Gymnosporangium spp.</i>) Scab (<i>Venturia</i> spp.) Sooty blotch (<i>Gloeodes pomigena</i>)</p>	<p>4 (0.065)</p>	<p>Apple Scab - Protective Spray Schedule: Apply every 7-10 days starting at ¼-½ inch green tip or when environmental conditions become conducive for scab. Continue through petal fall until the threat of primary scab is complete. For improved fruit scab control, combine Inspire with a protectant fungicide beginning at bloom.</p> <p>Apple Scab - Curative Spray Schedule: Use a forecasting system beginning at green tip. Apply within 48 hours of the onset of an infection period. Apply a follow-up spray within 7 days. For improved fruit scab control, combine Inspire with a protectant fungicide beginning at bloom.</p> <p>Powdery Mildew: Begin applications of Inspire at tight cluster, and continue on a 7- to 10-day schedule. Follow Inspire with other fungicides as needed.</p> <p>Cedar-Apple Rust, Quince Rust, and Brooks Fruit Spot: Begin applications preventively. Apply Inspire alone or in combination with a protectant fungicide on a 7- to 10-day schedule through the second cover-spray.</p> <p>Sooty Blotch, Flyspeck: For early season control, begin applications preventively. Apply Inspire alone or in combination with a protectant fungicide on a 7- to 10-day schedule through the second cover spray. Follow with other fungicides as needed.</p>
<p>Resistance Management: To help prevent resistance, make no more than 2 consecutive applications with Inspire or another Group 3 fungicide or a Group 3 fungicide mixture before alternating to a different mode of action.</p>			
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Use a minimum of 40 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			
<p>Pome Fruit subgroup: Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) Do not apply more than 20 fl oz/A/year of Inspire. 2) Do not apply more than 0.33 lb ai/A/year of difenoconazole-containing products. 3) Do not apply Inspire within 14 days of harvest (14-day PHI). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Potatoes Vegetable, Tuberous and corm subgroup 1C For other crops in this subgroup, please see below</p>	<p>Black dot (<i>Colletotrichum coccodes</i>) Brown spot (<i>Alternaria alternata</i>) Early blight (<i>Alternaria solani</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>) Septoria leafspot (<i>Septoria</i> spp.)</p>	<p>4-7 (0.065- 0.114)</p>	<p>Begin applications at first sign of disease or when conditions are conducive for disease development. Apply Inspire on a 7- to 14-day schedule. Inspire can be used in blocking program using a maximum of 2 consecutive applications before rotating to fungicides with another mode of action that are registered for these diseases.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Vegetables, tuberous and corm, subgroup: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (Taro), Ginger, Leren, Tanier, Tumeric, and Yam (bean and true).</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) Do not apply more than 28 fl oz/A/year of Inspire. 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) Do not apply within 14 days of harvest (14-day PHI). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Rapeseed Subgroup 20A Including cultivars, varieties, and/or hybrids of these and others in this group (see below).	Alternaria black spot <i>(Alternaria brassicae)</i> Black leg/Phoma <i>(Leptosphaeria maculans)</i> Cercospora leaf spot <i>(C. brassicicola)</i> Head rot <i>(Rhizoctonia solani)</i> Leaf spot and pod rot <i>(Alternaria alternata)</i> Powdery mildew <i>(Erysiphe polygoni)</i>	4 – 7 (0.065 – 0.114)	For Phoma control, apply during the rosette stage between 2nd true leaf and bolting. For Alternaria, make an application at the end of flowering/early pod set. For other foliar diseases, apply at first sign of disease. For head rot, apply at 50% flowering. If disease pressure is high, use the highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, air, or chemigation. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Complete List of Rapeseed Subgroup 20A: Borage, crambe, cuphea, echium, flax seed, gold of pleasure, hare's ear mustard, lesquerella, lunaria, meadowfoam, milkweed, mustard seed, oil radish, poppy seed, rapeseed, sesame, sweet rocket and cultivars and/or hybrids of these.

Specific Use Restrictions:

- 1) Do not apply more than one application of Inspire per year.
- 2) Do not apply more than 7 fl oz/A/year of Inspire.
- 3) Do not apply more than 0.114 lb ai/A/year of difenoconazole-containing products.
- 4) Do not apply within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Rice	Sheath Blight (<i>Rhizoctonia solani</i>) Aggregate Sheath Spot (<i>Rhizoctonia oryzae-sativae</i>) Black Sheath Rot (<i>Gaeumannomyces graminis var. graminis</i>) Sheath Spot (<i>Rhizoctonia oryzae</i>) Brown Leaf spot (<i>Cochliobolus miyabeanus</i>) Narrow Brown Leaf spot (<i>Cercospora oryzae</i>) Suppression only: False smut (<i>Ustilaginoidea virens</i>) Kernel Smut (<i>Neovossia barclayana</i>) Stem Rot (<i>Sclerotium oryzae</i>)	7.5 (0.122)	Apply 7.5 fl oz when disease is less than 4 inches above water line usually between panicle differentiation (PD) +5 days to PD +10 days or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied. The minimum re-treatment interval is 14 days.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not allow release of irrigation or flood water for at least 7 days after the last application.
- 2) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 3) Do not apply more than 15 fl oz/A/year of Inspire.
- 4) Do not apply more than 0.244 lb ai/A/year of difenoconazole-containing products.
- 5) Do not apply Inspire within 28 days of harvest (28-day PHI).
- 6) Do not treat fields used for aquaculture of fish or crustacean.
- 7) Do not drain water from treated rice fields into ponds used for aquaculture of fish or crustacean.
- 8) Do not use water drained from treated field to irrigate other crops.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Soybean	Anthracnose (<i>Colletotrichum truncatum</i>) Alternaria leaf spot (<i>Alternaria</i> spp.) Brown spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot (<i>C. kikuchii</i>) Frog-eye leaf spot (<i>Cercospora sojina</i>) Pod and stem blight (<i>Diaporthe phaseolorum</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	5.5-7 (0.09-0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. For aerial applications, use a minimum of 2 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Specific Use Restrictions:

- 1) Do not apply more than 13.5 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.22 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).
- 4) Do not feed soybean hay, forage and silage.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
<p>Stone Fruit Crop Group 12-12</p> <p>Apricots Cherries, Sweet Cherries, Tart Nectarines Peaches Plums Plumcot Prunes</p> <p>And cultivars and/or hybrids of these.</p>	<p>Brown rot blossom blight and fruit rot (<i>Monilinia fructicola</i>, <i>M. laxa</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>)</p> <p>Alternaria spot and fruit rot (<i>A. alternata</i>)</p> <p>Anthracnose (<i>Colletotrichum</i> spp.)</p> <p>Leaf rust (<i>Tranzschelia discolor</i>)</p> <p>Powdery mildew (<i>Sphaerotheca pannosa</i>, <i>Podosphaera clandestina</i>)</p> <p>Shot hole (<i>Wilsonomyces carpophilus</i>)</p>	<p>5.5-7 (0.09-0.114)</p>	<p>For Brown rot blossom blight, begin applications at early bloom and continue through petal fall. For Brown rot on fruit, apply as needed a maximum of two sprays during the preharvest period up to the day of harvest (minimum of a 7-day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered non-Group 3 fungicide.</p> <p>For all other diseases, follow the Brown rot blossom blight schedule. Make additional applications on a 10- to 14-day interval from the end of petal fall to harvest.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground or aerial application. Stone fruit diseases are most effectively controlled by ground applications. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>

Stone Fruit Crop Group: Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Inspire may be applied up to the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Sugar Beets	Cercospora leafspot (<i>C. beticola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)	4-7 (0.065-0.114)	<p>Apply Inspire on a 7- to 28-day schedule, alternating with a non-triazole fungicide (non-Group 3) that is registered on sugar beets for these diseases.</p> <p><i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.</p> <p><i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.</p> <p><i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 7 days of harvest (7-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Tree Nuts Crop Group 14-12 Beechnut Brazil Nut Butternut Cashew Chestnut Macadamia Walnut, Black Walnut, English See specific Directions for Almonds Filberts Pecans Pistachios	Foliar Diseases	5 - 7 (0.08 - 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate. <i>Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval. <i>Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.
Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by either ground, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.			

Tree Nuts Crop Group: African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these.

Specific Use Restrictions:

- 1) Do not apply more than 28 fl oz/A/year of Inspire.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply within 14 days of harvest (14-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Watercress	Cercospora leafspot (<i>Cercospora</i> spp.)	7.0 (0.114)	<p>For best activity, apply Inspire prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application.</p> <p>Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply directly to water and do not allow water in a treated field for at least 24 hours.
- 2) Do not apply more than 28 fl oz/A/year of Inspire.
- 3) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 4) Do not apply Inspire within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Remarks
Wild Rice	Helminthosporium leaf blight Brown Spot (<i>Bipolaris</i> spp.)	7.5 (0.122)	Apply 7.5 fl oz/A at both booting and heading. Minimum re-treatment interval is 14 days.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Inspire can be applied by ground or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not allow release of irrigation or flood water for at least 7 days after the last application.
- 2) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 3) Do not apply more than 15 fl oz/A/year of Inspire.
- 4) Do not apply more than 0.244 lb ai/A/year of difenoconazole-containing products.
- 5) Do not apply Inspire within 28 days of harvest (28-day PHI).
- 6) Do not use water drained from treated field to irrigate other crops.
- 7) Do not treat fields used for aquaculture of fish or crustacean.
- 8) Do not drain water from treated rice fields into ponds used for aquaculture of fish or crustacean.
- 9) Do not use water drained from treated field to irrigate other crops.

Inspire Conversion Table

Fl oz product/A	Lb ai/A
2	0.0325
3	0.05
4	0.065
5	0.08
5.5	0.09
6	0.10
7	0.114
7.5	0.122
7.7	0.125
8	0.13

Ornamentals – For outdoor application to established ornamentals in landscaped areas, to include recreational parks, institutional sites, golf courses and residential properties by commercial and non-professional (e.g., homeowner) applicators.

Use the calibrated measuring [device] [syringe] supplied with the product to accurately measure the designated amount of product to be added to each gallon of water.

Note: This use for application to ornamentals does not include crops grown for production, harvest and sale in commercial greenhouse or nursery production sites.

Ornamental	Disease	Rate Ml/1 Gal (1 Gal Treats 1000 Sq Ft)	Spray interval days	Timing of application
Carnation	Rust	4.8	7-10	Begin applications when flowers are half opened or when conditions are conducive for disease development. Apply Inspire on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Gladiolus	Rust	4.8	7-14	Begin applications at the 2- to 3-leaf stage or when conditions are conducive for disease development. Apply Inspire on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Iris	Rust	1.4-2.7	7-21	Begin applications at the first sign of disease or when conditions are conducive for disease development. Apply Inspire on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Roses	Black spot Rust Powdery mildew	4.8	7-14	Begin applications at the first sign of disease or when conditions are conducive for disease development. Apply Inspire on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.

Restrictions:

- For ornamentals, **do not** apply more than 4.8 ml/1000 sq ft of Inspire (0.115 lb ai/A) in a single application.
 - For ornamentals, **do not** apply more than 19.2 ml/1000 sq ft of Inspire (0.46 lb ai/A) per year.
-

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in the original container in a cool place.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative of the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Inspire® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

Viton® trademark of E. I. du Pont de Nemours and Company

©201X Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300