

100-1294

11/20/2008

1/19



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1294

Date of Issuance:

NOV 20 2008

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Cruiser® 70WS Insecticide

Name and Address of Registrant (include ZIP Code):

Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Continued on pages 2 and 3

Signature of Approving Official:

Venus Eagle
Venus Eagle, Product Manager

Insecticide-Rodenticide Branch, Registration Division (7505P)

Date:

NOV 20 2008

3. As a condition of registration, submit within 18 months of the date on this registration notice data depicting residues of thiamethoxam and CGA-322704 in plants grown from treated seed and planted at the maximum per-acre thiamethoxam loading rate (0.266 lb a.i./A). At a minimum, conduct three trials each of carrots, leaf lettuce, and mustard greens in Regions 3 (one trial) and 10 (two trials).
4. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Reg. No. so it reads, "EPA Reg. No. 100-1294."
 - b. On pages 1 and 14, change the word "certain" to "listed" so the statements reads, "A seed treatment product for control of **listed** insects..."
 - c. On page 3, under the PPE section, revise the third section heading so it reads, "**All other Workers and Handlers must wear: (Workers involved in bagging treated seed or sewing seed bags).**"
 - d. On page 4, under the Engineering Controls section, revise the text so it reads:

This product must be packaged in water soluble bags. **Handlers handling this product while it is enclosed in intact water-soluble packets may elect to not to wear the respirator.**

IMPORTANT: When the respirator is not worn because intact water-soluble packets are in use, handlers must still be provided and have immediately available a respirator for use in an emergency (such as the water soluble bag breaking prior to mixing).
 - e. On page 7, under the Mixing Procedures section, revise the first sentence so it reads: "Cruiser 70WS is a wettable powder **in water soluble bags**, formulated especially for seed treatment uses."
 - f. On page 8, under the Seed Bag Label Requirements section, add the following statement to the first paragraph: "**User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.**"
 - g. On page 8, under the Seed Bag Label Requirements section, add the following statements under the second paragraph: "**Do not use at a rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season**" and "**Contains x milligrams thiamthoxam per seed.**"
 - h. On page 9, under the Instructions for the Cruiser 70WS Applicator paragraph, delete the phrase "When using water soluble bags of Cruiser 70WS for delivery of the product..." Begin the paragraph with the sentence "Adjust the volume..."

3
19

- i. On page 9, under the Instructions for the Cruiser 70WS Applicator paragraph, revise the last sentence so it reads, "...in the treated seed container/**seed bag label.**" Move this heading and first paragraph so it appears at the very beginning of the Crop Use Directions section. These instructions are applicable to all crops.
 - j. On page 12, revise the heading "Root Vegetables: Carrots Only" so it reads, "Carrots."
 - k. On pages 13 and 16, under the Container Disposal section, delete references to plastic containers. For consistency with other products and per the recommendations of the Label Review Manual, it is suggested that the Storage and Disposal section be placed in a box of solid line to increase prominence.
5. For future submissions, make the following changes to your Confidential Statement of Formulation (EPA Form 8570-4):
- a. Box 4: Revise the EPA Reg. No. so it reads, "EPA Reg. No. 100-1294."

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Julie Chao at 703-308-8735 or chao.julie@epa.gov.

Venus Eagle, Product Manager
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

4
198

(Master Label)

Cruiser® 70WS
Insecticide

A seed treatment product for control of certain insects on Carrots, Leafy Vegetables and Brassica (Cole) Leafy Vegetables.

Active Ingredient:	
Thiamethoxam ¹	70.0%
Other Ingredients:	30.0%
Total:	100.0%

¹CAS No. 153719-23-4

Cruiser 70WS contains 0.7 pounds of Thiamethoxam per pound of formulated product.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-XXXXX

EPA Est. XXXX

Product of xxxx
Formulated in xxxx

SCP xxxxA-M(draft-Cruiser 70WS)

Net Weight

ACCEPTED
With COMMENTS
In EPA Letter Dated:
NOV 20 2008

Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
pesticide Registered under EPA Reg. No:

100-1294

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

Personal Protection Equipment (PPE)

Workers involved with Treating the Seed must wear:

(Workers involved with treating the seed are responsible for loading and treating raw seed, applying coating material, liquids and binders to form the seed pellets)

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material - Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks
- Respirator protection (i.e., single use dust mask, quarter or half mask respirator)

Multiple Task Workers must wear:

(Multiple task workers perform multiple tasks in one day such as mixing, product application, bagging/filling seed containers, bag sewing, and clean-up)

- Chemical-resistant coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material - Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks
- Respirator protection (i.e., single use dust mask, quarter or half mask respirator)

All Other Applicators and Other Handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material - Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks

User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

This product must be packaged in water soluble bags.

User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling.
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment washwater.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use in hopper box, planter box, slurry box, or other farmer-applied applications. Apply Cruiser 70WS in commercial seed treatment facilities only. For lettuce seed (head, leaf and romaine): Apply Cruiser 70WS to pelleted seed using conventional coating pans (i.e., batch pans) only. Do not use rotary coaters.

Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Cruiser 70WS (thiamethoxam).

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality, crop or variety, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

USE INFORMATION

Cruiser 70WS is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Cruiser 70WS provides protection against certain chewing and sucking insects following contact and ingestion. Specifically, Cruiser 70WS is shown to provide early season protection against aphids, cabbage maggots, certain seed and root maggots, flea beetles, leaf miners, whiteflies and wireworm.

For protection against certain diseases Cruiser 70WS may be applied along with labelled fungicides. Consult professional applicators or university extension service for approved fungicides.

Cruiser 70WS must be applied in conjunction with polymers, pelleting materials and seed coating materials that are tested to be safe on leafy vegetable, *Brassica* vegetable, or carrot seed. The end product that combines Cruiser 70WS with labelled fungicides must be tested for seed safety to determine any detrimental effects on seed germination or plant stand establishment. Pre-test the germination of a small sample of seed lot with Cruiser 70WS prior to large scale commercial application.

Resistance Management

Cruiser 70WS contains a Group 4A insecticide (thiamethoxam, belonging to the neonicotinoid class of chemistry). Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Cruiser 70WS or other Group 4A insecticides.

In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season, foliar applications of Group 4A insecticides should not be made following a seed or soil applied Group 4A insecticide.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action as long as the involved products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

MIXING PROCEDURES

Cruiser 70WS is a wettable powder formulated especially for seed treatment uses. Cruiser 70WS must be applied through an approved pelleting film coating process or a standard delivery system that is deemed safe to the seed. Cruiser 70WS must be applied to pelleted lettuce seed (head, mature leaf and romaine) only. Uneven or incomplete seed coverage may not give the desired level of insect control. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment.

Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. Ensure seed is dry before bagging.

Add 1/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Cruiser 70WS into the tank all at once. Continue agitation while adding the remainder of the water. Allow the packets of Cruiser 70WS to dissolve and the product to completely disperse into the

mix water. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

Each foil envelope of Cruiser 70WS contains water-soluble bags of Cruiser 70WS wettable powder. The foil envelope must be opened for use. Because water-soluble bags readily dissolve in the presence of moisture, do not allow them to get wet before adding to the applicator tank. Wear dry gloves when handling the water-soluble bags. Handle with care as excessive handling may cause rupture of the water-soluble bag. Unused water-soluble bags should be stored in the original foil envelope, folded over to minimize exposure to moisture.

SEED BAG LABEL REQUIREMENTS

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with thiamethoxam insecticide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs."

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with Cruiser® 70WS (thiamethoxam):

- Store away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and waterproof gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Left over treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Cruiser 70WS (thiamethoxam).

CROP USE DIRECTIONS

Leafy Vegetables (except *Brassica*) Crop Group

Crops in this group are: Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (roquette); Cardoon; Celery; Celtuce; Chervil; Chinese celery; Edible-leaved chrysanthemum; Garland chrysanthemum; Corn salad; Cress (garden and upland); Dandelion; Dock (sorrel); Endive (escarole); Fennel, Florence (finocchio); Head lettuce; Leaf lettuce; Orach; Parsley; Purslane (garden and winter); Radicchio (red chicory); Rhubarb; Spinach; New Zealand spinach; Vine spinach (Malabar spinach, Indian spinach); Swiss chard

Cruiser 70WS seed treatment provides early season protection against injury from aphids, flea beetles, leaf miners, and whiteflies for certain leafy vegetables (except *Brassica*).

Pelleted Lettuce Seed (Head, Mature Leaf and Romaine Lettuce)

The application rate of Cruiser 70WS is dependent on the expected seeding rate per acre. Refer to the table listed under **Instructions for the Seed Planter** later in this label to determine how much Cruiser 70WS to apply to each seed.

Instructions for the Cruiser 70WS Applicator

When using water-soluble bags of Cruiser 70WS for delivery of the product, adjust the volume (or count) of seed for each batch to the exact amount of active ingredient available in each water-soluble bag. Once Cruiser 70WS is applied and the treated seed is packaged, indicate the amount (number of seeds) and the Cruiser 70WS rate in milligrams thiamethoxam per seed on the treated seed container.

For head, romaine and leaf lettuce, Cruiser 70WS must be applied in conjunction with polymers, pelleting materials and seed coating materials that are tested to be safe on the seed. Cruiser 70WS is applied during the process of pelleting lettuce seed and therefore all precautionary measures mentioned in the label must be followed. For lettuce seed (head, mature leaf, and romaine), apply Cruiser 70WS to pelleted seed using conventional coating pans (i.e., batch pans) only. Do not use rotary coaters. Consult with Syngenta Crop Protection for advice on application. The end product that combines Cruiser 70WS with labelled fungicides must be tested for seed safety to determine any detrimental effects on seed germination or plant stand establishment. Pre-test the germination of a small sample of seed lot with Cruiser 70WS prior to large scale commercial application.

$$\frac{13}{19}$$

Instructions for the Lettuce Seed Planter

Determine the seeding rate of the lettuce seed per acre and select the appropriate treated seed container showing the seeds per acre to be planted and the corresponding Cruiser 70 WS rate from the table below.

Seeds per Acre	Milligrams Active Ingredient (thiamethoxam) per Seed
35,000-50,000	1.2
60,000	1.0
70,000	0.9
80,000	0.8
90,000	0.7
100,000	0.6
110,000	0.54
120,000	0.5
130,000	0.46
140,000	0.43
150,000	0.4
160,000	0.38
170,000	0.35
180,000	0.33
190,000	0.32
200,000	0.3

The total thiamethoxam active ingredient load must not exceed 120.66 grams per acre, per season.

For protection against certain diseases, Cruiser 70 WS may be applied along with labelled fungicides. Consult professional applicators or university extension service for approved fungicides.

Spinach

Cruiser 70WS may reduce the injury from aphids, flea beetles, leaf miners, whiteflies for spinach. Cruiser 70WS will not provide complete protection against the damage caused by these insects.

Suggested Seeding Rates	Milligrams (thiamethoxam) per Seed
Number of Seeds per Acre	
Up to 1,000,000	0.12

Baby leafy lettuce and mesclan mixes

Cruiser 70WS may reduce the injury from aphids, flea beetles, leaf miners, whiteflies for baby leafy lettuce and mesclan mixes. Cruiser 70WS will not provide complete protection against the damage caused by these insects.

Suggested Seeding Rates.	Milligrams (thiamethoxam) per Seed
Number of Seeds per Acre	
Up to 2,000,000	0.06
Up to 3,000,000	0.04

Note to the applicator: Once Cruiser 70WS is applied and the treated seed is packaged, indicate the amount (number of seeds) and the Cruiser 70WS rate in milligrams thiamethoxam per seed on the treated seed container.

Refer to university extension guidelines or reliable local recommendations for seeding rates of individual crops.

Restrictions:

Do not use a Cruiser 70WS rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season.

Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Cruiser 70WS (thiamethoxam).

Brassica (Cole) Leafy Vegetables

Crops in this group are: Broccoli; Broccoli raab (rapini), Brussels sprouts; Cabbage; Chinese broccoli (gai lon); Chinese cabbage (bok choy); Chinese cabbage (napa); Chinese mustard cabbage (gai choy); Cauliflower; Cavalo broccolo; Collards; Kale; Kohlrabi; Mizuna; Mustard greens; Mustard spinach; Rape greens

To provide early season protection of seedlings against injury from aphids, cabbage maggots, certain seed and root maggots, flea beetles, whiteflies and wireworm, apply Cruiser 70WS at 0.05 to 0.10 mg. per seed.

Refer to university extension guidelines or reliable local recommendations for seeding rates of individual crops.

Restrictions:

Do not use a Cruiser 70WS rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season.

Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Cruiser 70WS (thiamethoxam).

Root Vegetables: Carrots only

To provide early season protection of seedlings against injury from aphids, beetles, certain seed and root maggots, flea beetles, whiteflies and wireworm, apply Cruiser 70WS at 0.03 to 0.05 mg. per seed.

Refer to university extension guidelines or reliable local recommendations for seeding rates of individual crops.

Restrictions:

Do not use a Cruiser 70WS rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season.

Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Cruiser 70WS (thiamethoxam).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Keep water-soluble bags in original foil package and handle carefully when stored at temperatures less than 50°F to avoid breakage of bags.


Pesticide Disposal

Wastes resulting from the use of this product maybe disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Container Disposal (bags, plastic containers, bulk and mini-bulk and foil pouches/carton that accompany water-soluble bags as long as water-soluble bag is unbroken)

Non-refillable container. Do not use or refill this container. Offer for recycling if available.

Cruiser®, the Syngenta logo and the CP FRAME  are trademarks of a Syngenta Group Company
U.S. Patent No. xxxxx
©2008 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 942A-M(draft-Cruiser 70WS)

Cruiser 70WS draft-clean pm-B-11-19-08

17
19

(Non-detachable container label for booklet)

Cruiser® 70WS
Insecticide

A seed treatment product for control of certain insects on Carrots, Leafy Vegetables and *Brassica* (Cole) Leafy Vegetables.

Active Ingredient:	
Thiamethoxam ¹	70.0%
<hr/>	
Other Ingredients:	30.0%
<hr/>	
Total:	100.0%

¹CAS No. 153719-23-4

Cruiser 70WS contains 0.7 pounds of Thiamethoxam per pound of formulated product.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-XXXXX

EPA Est. XXXX

SCP xxxxA-M(draft-Cruiser 70WS)

Net Weight

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Keep water-soluble bags in original foil package and handle carefully when stored at temperatures less than 50°F to avoid breakage of bags.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Container Disposal (bags, plastic containers, bulk and mini-bulk and foil pouches/carton that accompany water-soluble bags as long as water-soluble bag is unbroken)

Non-refillable container. Do not use or refill this container. Offer for recycling if available.

Cruiser®, and the Syngenta logo are trademarks of a Syngenta Group Company
U.S. Patent No.
©2008 Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 942A-M(draft Cruiser 70WS)

Cruiser 70WS draft-clean-pm-B-11-19-08