111 12 200

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460



OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

November 12, 2009

Patrick McCain Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, NC 27419-8300

Dear Mr. McCain:

Subject: Label Notification per Pesticide Registration Notice 98-10

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10, dated November 5, 2009, for the product "CruiserMaxx[™] Cereals Insecticide with Fungicides," EPA Registration Number 100-1305.

The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10 Section II-B. The information submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please contact me directly at 703-308-8735 or email me at chao.julie@epa.gov.

Sincerely,

Jane Alao

Julie A. Chao Insecticide-Rodenticide Branch Registration Division (7505P) Office of Pesticide Programs

	<u>, un rereite serere con</u> ina. /	<u></u>	·····		6	1002
SEPA	Environmental Washing	J States Protection A ton, DC 20460	lgency	Ame X Oth	er, it	OPP Identifier Number
		Application for	or Pesticide - Se	ction I		
1. Company/Product N 100-1305	umber		2. EPA Pro Venus Eagle	duct Mana	iger 3.	Proposed Classification
4. Company/Product (N CruiserMaxx [™] Cer	ame) eals Insecticide with	Fungicides	PM# PM 01 IRB		×	None Restricted
Syngenta Crop P. O. Box 1830)	ode)	my product is sin	Review. In nilar or identi	accordance with cal in correction	FIFRA Section 3(c)(3) (b)(i),
Greensboro, NO	if this is a new address		EPA Reg. No. Product Name	- <u></u>	NUV	2 2009
			Section - II			
Amendment - Ex	plain below. response to Agency letter of	dated		Final printed Agency lette "Me Too" Ap	r dated	bels in response to NOTIFICATION
x Notification - Exp				Other - Expla		NOV 1 2 2009
Syngenta Crop Prote Reg. No. 100-1305) highlighted) are attac	nd I may be subject to enfor ction Inc., is submitting to add a disease as all hed. is SCP 1305A-M(Notif	g a Notificatio owed by PR N	n for CruiserMax	k™ Cereal	s Insecticide v	
			ection – III			
	ct Will Be Packaged In:					
Child-Resistant Packaging Yes* No	Unit Packaging		Water Soluble Pac	kaging	2. Type of (Container Metal Plastic
*Certification must be submitted	lf "Yes" Unit Packaging wg		lf "Yes" Unit Packaging wgt.	No. per container		Glass Paper Other (Specify)
3. Location of Net Cont	-	4. Size(s) R	Retail Container	Ę	On Lal	
Label	_ Container			L	On Lai	beling accompanying product
6. Manner in Which Lab	el is Affixed to Product	Lithogi Paper Stencil	glued	Other		
		S	ection – IV			
	olete items directly below fo			acted, if nece		
Name Patrick McCain		R	^{itle} Regulatory Produc Syngenta Crop Pro		r 336-6	one No. (Include Area Code) 032-7317
l acknowledge that a both under applicable	ments I have made on this ny knowingly false or mislea i law.	ading statement r	chments thereto are t may be punishable by		conment or	6. Date Application Rieduived (Stamped)
2. Signature	ude alo le	Sr. Ro	itle egulatory Product enta Crop Protect			
4. Typed Name Patrick McCain		5. D Nove	ate mber 5, 2009		((((
EPA Form 8570-1 (R	ev. 8-94) Previous edi	tions are obso	olete.			

1.1.1

υ ς αυμείς

000000

666

0000

0000

0.0

00000

00000

Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, NC 27419-8300



November 5, 2009

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

ATT: Ms. Linda Arrington

SUBJECT: CRUISERMAXX™ CEREALS INSECTICIDE WITH FUNGICIDES EPA REG. NO. 100-1305 NOTIFICATION TO ADD A DISEASE PER PR NOTICE 98-10

Dear Ms. Arrington,

Syngenta Crop Protection Inc., is submitting a Notification for CruiserMaxx[™] Cereals Insecticide with Fungicides (EPA Reg. No. 100-1305) to add a disease as allowed by PR Notice 98-10. Two copies of labeling (one with changes highlighted) are attached.

The label copy code is SCP 1305A-M(Notif).

To complete this Notification, EPA form 8570-1 is enclosed.

We trust that you will find this submission in order. However, should you have any questions, please contact me at 336.632-7317 or via e-mail at patrick.mccain@syngenta.com.

Sincerely,

la Cin

Patrick McCain Sr. Regulatory Product Manager Syngenta Crop Protection, Inc.

Enclosures:

2 copies of labeling (1 highlighted) EPA Form 8570-1

cc: Venus Eagle PM Team (01) IRB

CruiserMaxx™ Cereals

Insecticide with Fungicides

A seed treatment product for protection against damage from certain insects, seedborne diseases, and seedling diseases of cereals

•
2.80%
0.56%
3.36%
93.28%
100.00%

*CAS No. 153719-23-4 **CAS No. 70630-17-0 and CAS No. 69516-34-3 ***CAS No. 119446-68-3

One gallon of CruiserMaxx Cereals contains 0.26 pounds thiamethoxam, 0.05 pounds mefenoxam and 0.31 pounds difenoconazole.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Si usted no entienda la etiqueta busque a alguien para se la explique en detalle. (If you do not understand the label find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1305 EPA Est. XXXXX

SCP 1305A-M(Notif)

NOTIFICATION

NOV 1 2 2009

Net Contents

. . . i c

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. If swallowed • Call a poison control center or doctor immediately for treatment advice. If swallowed • Call a poison control center or doctor immediately for treatment advice. If swallowed • Call a poison control center or doctor immediately for treatment advice. If on skin or clothing • Take off contaminated clothing. If on skin or clothing • Take off contaminated clothing. If inhaled • Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-tomouth, if possible. • Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call		FIRST AID
Iteratment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. If inhaled • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor for further treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	lf in eyes	 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment
 clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) 	If swallowed	 treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious
 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to- mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) 		 Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment
or doctor or going for treatment. HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)		 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to- mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)		
For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)	or doctor or going	
1-800-888-8372		ur Medical Emergency Assistance (Human or Animal) al Emergency Assistance (Spill, Leak, Fire or Accident) Call

,

2

50f26

•.:

.

•••

(((() ()

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material Category A (e.g., barrier laminate, butyl rubber ≥14 mils, natural rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyethylene, polyvinyl chloride [PVC] ≥14 mils or viton ≥14 mils)
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

11111

Engineering Control Statements

For commercial seedtreatment applications only, applicators must use a closed system. When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Exposed treated seeds may be hazardous to wildlife.

Groundwater Advisory

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of the SYNGENTA.

. . . .

5

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restrictedentry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material Category A (e.g., barrier laminate, butyl rubber ≥14 mils, natural rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyethylene, polyvinyl chloride [PVC] ≥14 mils or viton ≥14 mils)
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES. Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a small portion of the seed before committing the total seed lot to a selected seed treatment.

7

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with CruiserMaxx Cereals.

GENERAL INFORMATION

CruiserMaxx Cereals is a seed treatment product containing the active ingredients: thiamethoxam (insecticide) and difenoconazole and mefenoxam (fungicides). CruiserMaxx Cereals protects against damage from certain early season insects and also protects against certain seed- and soil-borne diseases of cereal crops.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neoticotinoid class of chemistry. Thiamethoxam protects against certain chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against Pythium.

Difenoconazole fungicide is active against several seed, soil and foliar diseases, as listed in the table in the **Crop Use Directions** section.

MIXING PROCEDURES

Important: Always re-circulate CruiserMaxx Cereals thoroughly before using.

CruiserMaxx Cereals is a ready-to-use water-based formulation for use in commercial seed treatment facilities utilizing closed-system application techniques. In addition, CruiserMaxx Cereals may also be applied by on-site/on-farm applications. Apply CruiserMaxx Cereals as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect and disease control. If necessary, thoroughly mix the recommended amount of CruiserMaxx Cereals into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

The typical density of CruiserMaxx Cereals is 9.2 pounds per gallon. Concut the manufacturer of the application equipment you plan to employ for suitability for this capplication and for instructions on operation and calibration of the equipment. Follow, the manufacturer application instructions for the seed treatment equipment being used.

Under certain disease conditions, additional amounts of fungicides may be required. When needed, apply additional Apron® XL LS according to the **CROP USE DIRECTIONS**. Other tank mix partners may be used with CruiserMaxx Cereals; however, the user must consider the use rate, formulation, seed and crop safety factors and compatibility of each product to be mixed when determining the total application volume.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Ideal storage temperature is above freezing and below 86°F (30 °C). Repeated freezing and thawing of CruiserMaxx Cereals will not affect the physical integrity of the product. If the product should freeze, bring the product back to room temperature and ensure the contents are mixed well prior to application.

Allow seed to dry before bagging.

Follow planter manufacturer recommendations for use of talc or other hopper box additives at planting. Seed should be completely dry before adding to planter.

Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of the treated seed as food for man or feed for animals. Refer to 21 CRF, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 180.1001.

CruiserMaxx Cereals contains an EPA approved dye/colorant that imparts an unnatural color to the seed as required by the Federal Seed Act.

SEED BAG LABEL REQUIREMENTS

Federal law requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and difenoconazole and mefenoxam fungicides.
- Do not use treated seed for feed, food, or oil purposes.
- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.

- Dispose of seed packaging or containers in accordance with local requirements.
- For seed treated with CruiserMaxx Cereals, do not graze or feed livestock on treated areas for 45 days after planting.

CROP USE PRECAUTIONS

Resistance Management

CruiserMaxx Cereals contains thiamethoxam insecticide which belongs to the neonicotinoid class of chemistry, which is not known to be cross-resistant to any other class of insecticide. However, insect pests are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. The following label restrictions must be followed to reduce the chance of resistance development:

• Do not apply a neonicotinoid insecticide to any crop which has been grown from CruiserMaxx Cereals treated seed.

CruiserMaxx Cereals contains the fungicide mefenoxam, a systemic fungicide having a specific mode of action against fungal pathogens.

Note: The fungicide metenoxam contained in this product could be subject to development of insensitive strains of fungi or may be ineffective against naturally occurring strains of fungi. Development of insensitivity or natural tolerance cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive or tolerant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible insensitive or tolerant strains of fungi which may occur.

Consult your local pest control advisor or extension office for additional methods for preventing resistance development. Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal and insect pests on this label.

CROP USE DIRECTIONS

USE RATES

For protection against certain seed and soil-borne diseases and insects on small-grain cereals, apply 5.0 fluid ounces of CruiserMaxx Cereals per 100 lb. of seed to the crops listed in the following table.

10

(

ſ

130926

Crop	Pests	
	Insects	
Barley	Wireworms ¹	
•	Aphids ²	
	Diseases Controlled	
	General Seed Rots ³	
<u>د</u>	Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borr	ne Fusarium
	and soil-borne Pythium ⁴	le l'acanam
	Early Season Rhizoctonia Root Rot	
	Covered and False Loose Smut	
	Diseases Suppressed ⁵	
	Common Root Rot (Cochliobolus spp.)	
	Fusarium Crown and Foot Rot	
	Take-All	
	Insects	•
Triticale	Wireworms ¹	
	Aphids ²	·
	Diseases Controlled	
	General Seed Rots ³	
	Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borr	ne Fusarium
	and soil-borne Pythium ⁴	
	Early Season Rhizoctonia Root Rot	
	Loose Smut	
	Diseases Suppressed ⁵	
	Common Root Rot (Cochliobolus spp.)	
	Fusarium Crown and Foot Rot	
	Take-All	•
	Insects	
Winter Wheat	Wireworms ¹	
white wheat	Aphids ²	
	Hessian Fly ²	
	Diseases Controlled	
	General Seed Rots ³	
		no Eucorium
	Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borr	le rusanum
	and soil-borne Pythium ⁴	
	Common and Dwarf Bunt ⁶	
	Loose Smut	
	Diseases Suppressed ⁵	
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.)	
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season <i>Rhizoctonia</i> Root Rot	
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot	
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season <i>Rhizoctonia</i> Root Rot	
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot	
Spring Wheat	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All	
Spring Wheat	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects	
Spring Wheat	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ²	, , , , , , , , , , , , , , , , , , ,
Spring Wheat	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) <u>Early Season Rhizoctonia Root Rot</u> Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ²	(, , () () () ()
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled	(((
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³	τις τις τις τ
	Diseases Suppressed ⁵ Common Root Rot (Cochliobolus spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-bohr	τις τις τις τ
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and s¢il-born and soil-borne Pythium ⁴	e Fusarium
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot	te Fusarium
	Diseases Suppressed ⁵ Common Root Rot (Cochliobolus spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot Common Bunt ⁶	e Fusarium
	Diseases Suppressed ⁵ Common Root Rot (<i>Cochliobolus</i> spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot	e Fusarium
	Diseases Suppressed ⁵ Common Root Rot (Cochliobolus spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot Common Bunt ⁶	e Fusarium
	Diseases Suppressed ⁵ Common Root Rot (Cochliobolus spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot Common Bunt ⁶	د د (
	Diseases Suppressed ⁵ Common Root Rot (Cochliobolus spp.) Early Season Rhizoctonia Root Rot Fusarium Crown and Foot Rot Take-All Insects Wireworms ¹ Aphids ² Hessian Fly ² Diseases Controlled General Seed Rots ³ Seedling Blight, Root Rot, and Damping-Off caused by seed- and scil-born and soil-borne Pythium ⁴ Early Season Rhizoctonia Root Rot Common Bunt ⁶	د E Fusarium (۱۹۹۵) (۱۹۹) (۱۹۹۵) (۱۹۹) (۱۹) (۱) (۱) (۱) (۱) (۱) (۱) (۱) (۱

Сгор	Pests
	Diseases Suppressed ⁵
	Common Root Rot (Cochliobolus spp.)
	Fusarium Crown and Foot Rot
	Take-All

¹ CruiserMaxx Cereals provides suppression of wireworm activity; however, if the wireworm pressure is high, additional protection may be required. Under these circumstances, mix 5.0 fluid ounces of CruiserMaxx Cereals with 0.25 fluid ounces of Cruiser[®] 5FS. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

² For protection against aphids and Hessian fly, an additional amount of Cruiser 5FS must be mixed with CruiserMaxx Cereals. Mix 5.0 fluid ounce of CruiserMaxx Cereals with 0.48 to 1.0 fluid ounce of Cruiser 5FS with appropriate quantities of water and apply to 100 pounds of seed. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

³ Protection against general seed rots include those caused by saprophytic organisms such as <u>Rhizoctonia</u> Fusarium, Pythium, Penicillium and Aspergillus.

⁴ For additional *Pythium* protection, add 0.0425 fluid ounces of Apron XL[®] LS per 100 pounds of seed.

⁵ Suppression means effective protection at a level that could provide commercial benefit.

⁶ Provides protection against both seed- and soil-borne bunts (common, dwarf).

Post-Treatment Storage Protection

When treated according to the directions for post-planting protection against listed pests indicated above, CruiserMaxx Cereals will also provide protection during post treatment storage of the seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If seed to be treated has existing infestations of stored grain insects, it is recommended that the seed be fumigated with labelled products, prior to treating with CruiserMaxx Cereals and bagging.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [Bulk/Mini-Bulk]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least once complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repetit this procedure two more times. Then offer for recycling if available or puncture and the container of the container or times.

dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Apron XL[®], Cruiser®, CruiserMaxx[™] Cereals, the Syngenta logo and the CP FRAME

are trademarks of a Syngenta Group Company

U.S. Patents 5,852,012 and 6,022,871 and other pending US patent applications. ©20098 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, North Carolina 27419-8300 www.syngenta-us.com

SCP 1305A-M(Notif)

CruiserMaxx Cereals 1305A-M(Notif)marked-lg-11-5-09)

160524

(Non-detachable Container Label - 1 gal, 2.5 gal, 15 gal.)

CruiserMaxx[™] Cereals

Insecticide with Fungicides

A seed treatment product for protection against damage from certain insects, seedborne diseases, and seedling diseases of cereals

Active Ingredients:	
Thiamethoxam*	2.80%
Mefenoxam**	0.56%
Difenoconazole***	
Other Ingredients:	93.28%
Total:	100.00%

*CAS No. 153719-23-4

CAS No. 70630-17-0 and CAS No. 69516-34-3 *CAS No. 119446-68-3

One gallon of CruiserMaxx Cereals contains 0.26 pounds thiamethoxam, 0.05 pounds mefenoxam and 0.31 pounds difenoconazole.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Si usted no entienda la etiqueta busque a alguien para se la explique en detalle. (If you do not understand the label find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1305		ι <u>ι</u> ετεςτη ι
EPA Est. XXXXX SCP 1305A-M(Notif)	. ((()) () () () () () () () () () () () () () () (י ((((הרול (הרול (הרול (הרול)
Net Contents		

PRECAUTIONARY STATEMENTS

	FIRST AID	
lf in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
lf on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 	
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to- mouth, if possible. Call a poison control center or doctor for further treatment advice. 	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.		
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372		

Hazards to Humans and Domestic Animals

CAUTION

. .

.

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor (Wash (Content with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Exposed treated seeds may be hazardous to wildlife.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [Bulk/Mini-Bulk]

Non-refillable container. Do not reuse or refill this container. Offer for recycliny if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least once complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container or onto its other end and tip it back and forth several times. Empty the circulate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and

> **ι** (ι ι ι(ι)

dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CruiserMaxx[™] and the Syngenta logo are trademarks of a Syngenta Group Company U.S. Patents 5,852,012 and 6,022,871 and other pending US patent applications.

©20098 Syngenta

Manufactured for: Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, North Carolina 27419-8300 www.syngenta-us.com

SCP 1305A-M(Notif)

CruiserMaxx Cereals 1305A-M(Notif)marked-Ig-11-5-09

200f210

ειιι ι ιειιι (Non-detachable Container Label - Bulk)

CruiserMaxx[™] Cereals

Insecticide with Fungicides

A seed treatment product for protection against damage from certain insects, seedborne diseases, and seedling diseases of cereals

Active Ingredients:	
Thiamethoxam*	
Mefenoxam**	0.56%
Difenoconazole***	3.36%
Other Ingredients:	93.28%
Total:	100.00%

*CAS No. 153719-23-4 **CAS No. 70630-17-0 and CAS No. 69516-34-3 ***CAS No. 119446-68-3

One gallon of CruiserMaxx Cereals contains 0.26 pounds thiamethoxam, 0.05 pounds mefenoxam and 0.31 pounds difenoconazole.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Si usted no entienda la etiqueta busque a alguien para se la explique en detalle. (If you do not understand the label find someone to explain it to you in detail.)

.....

· · · · ·

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1305

EPA Est. XXXXX

SCP 1305A-M(Notif)

Net Contents

PRECAUTIONARY STATEMENTS

	FIRST AID
lf in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
lf on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
lf inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to- mouth, if possible. Call a poison control center or doctor for further treatment advice.
Have the product of or doctor or going f	ontainer or label with you when calling a poison control center or treatment.
	HOT LINE NUMBER ur Medical Emergency Assistance (Human or Animal) al Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372
Hazards to Huma	ns and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing vapore Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not contaminate water when disposing of equipment wash water.

Exposed treated seeds may be hazardous to wildlife.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or har dling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES; INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT; SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

20

1302710

1 () 1 () 1 () SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment of a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least once complete revolution, for 30 ceconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into

.....

application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CruiserMaxx[™], the Syngenta logo and the CP FRAME *I* are trademarks of a Syngenta Group Company U.S. Patents 5,852,012 and 6,022,871 and other pending US patent applications.

©20098 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, Inc. P.O. Box 18300 Greensboro, North Carolina 27419-8300 <u>www.syngenta.us.com</u>

SCP xxxxA-M(Notif)

CruiserMaxx Cereals 1305A-M(Notif)marked-Ig-11-5-09)

2005-20

ιιι

eco

Chron

. :

4-16-08 – Approved with comments.

7-2-09 – Approved – notification for PR Notice 2007-4 conainer handling 11-5-09 Notification to add *Rhizoctonia Root Rot*