

100-1318

09/19/2008

1/22



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Timothy Wilson
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419

SEP 19 2008

Dear Mr. Wilson:

Subject: Labeling Amendment; Revised Rotational Crop Section
Durivo
EPA Registration No. 100-1318
Submission Date: September 3, 2008

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this label, please contact me at (703) 306-0415.

Sincerely yours,

A handwritten signature in black ink, appearing to read "Kable Bo Davis".

Kable Bo Davis
Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure- Stamped Labeling

(Booklet)

DURIVO™

GROUP	4A	INSECTICIDE
GROUP	28	INSECTICIDE

For control of listed insect pests infesting various crops

Active Ingredient:

Thiamethoxam¹ 17.5%

Chlorantraniliprole² 8.8%

Other Ingredients: 73.7%

Total: 100.0%

¹CAS No. 153719-23-4

²CAS No. 500008-45-7

Durivo is a soluble concentrate containing 1.67 lb. of thiamethoxam and 0.835 lb. chlorantraniliprole per gallon.

KEEP OUT OF REACH OF CHILDREN.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1318

EPA Est. XXXXX

SCP 1318A-M(draft)

Net Contents



PRECAUTIONARY STATEMENTS

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) or
Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call

1-800-888-8372

Have the product container or label with you when calling a poison control center
or doctor, or going for treatment.

Hazards to Humans and Domestic Animals

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves – Category C such as barrier laminate; butyl rubber ≥ 14 mils; nitrile rubber ≥ 14 mils; neoprene rubber ≥ 14 mils; polyvinyl chloride (PVC) ≥ 14 mils; or Viton[®] ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates, oysters and shrimp. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

- **Surface Water Advisory**

This product may contaminate water through runoff. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (see manual at the following Internet address:

<http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>)

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

- **Ground Water Advisory**

This product has properties and characteristics associated with chemicals detected in ground water. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. TO THE EXTENT PERMITTED BY APPLICABLE LAW, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, Category C, such as barrier laminate; butyl rubber \geq 14 mils; nitrile rubber \geq 14 mils; neoprene rubber \geq 14 mils; polyvinyl chloride (PVC) \geq 14 mils; or Viton[®] \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

INFORMATION

Durivo is a broad spectrum, systemic insecticide that controls key lepidopteran, sucking and chewing insect pests in vegetables. When Durivo is applied through drip (trickle) chemigation, it is readily taken up by the roots of germinating seedlings or transplants and is rapidly translocated throughout the plant. Because of its systemic activity, Durivo provides excellent residual control of key insect pests in vegetables.

For best performance, always follow these directions:

- Durivo should be applied only via drip (trickle) chemigation which allows the insecticide to be absorbed by plant roots. Although Durivo is rapidly taken up by plant roots and rapidly moves throughout the plant, the use of sufficient water volume will ensure that the chemical contacts the roots, resulting in optimal uptake and performance.
- Because Durivo is applied through drip (trickle) chemigation and controls labeled insects that ingest plant parts and fluids, it should have minimal impact on beneficial arthropods, and its use is compatible with integrated pest management programs.
- In addition to control of key insect pests on this label, Durivo may aid in the suppression of other pests. Suppression can mean either inconsistent control (good to poor), or consistent control at a level below that generally considered acceptable for commercial control.
- Durivo has a wide margin of plant safety when used in accordance with this label.

RESTRICTIONS

Do not treat plants grown for transplanting. Not for use in nurseries, plant propagation houses, or greenhouses by commercial transplant producers on plants being grown for transplanting.

- Use this product only in commercial and farm settings.
- Not for use on ornamental plants or plants being grown for ornamental purposes.
- Not for use in home plantings.

RESISTANCE MANAGEMENT

Some insect pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

Durivo contains a Group 4A insecticide (thiamethoxam, belonging to the neonicotinoid class of chemistry) and a Group 28 insecticide (chlorantraniliprole, belonging to the anthranilic diamide class of chemistry). Insect biotypes with acquired or inherent resistance to Group 4A or Group 28 insecticides may eventually dominate the insect population if Group 4A or Group 28 insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Durivo or other Group 4A or Group 28 insecticides.

In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season:

- 1) Only a single, drip (trickle) chemigation application of Durivo should be made per crop season.
- 2) Foliar applications of Group 4A insecticides should not be made following a long residual, soil application of Durivo, or other seed or soil applied Group 4A insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action Group as long as the mixture products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

APPLICATION PROCEDURES

RESTRICTIONS

Durivo must only be applied via drip (trickle) chemigation.

Do not apply Durivo as a broadcast foliar spray with ground or aerial equipment.

Drip (trickle) Chemigation

Applications of Durivo alone or in combination with other pesticides registered for application through drip irrigation systems may be applied in irrigation water at rates recommended on this label. This product may be applied only through drip type irrigation systems. Do not apply Durivo through any other type of irrigation system.

DIRECTIONS FOR RECOMMENDED IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Dilute Durivo in the solution tank at a ratio of at least 15 parts of water to one part of Durivo. Inject Durivo solution at a ratio of 50:1 or greater. Injecting a larger volume of a more dilute solution will usually allow a more accurate calibration of the metering equipment. Meter the insecticide into the irrigation water during the irrigation cycle.

Using Water from Public Water Systems: DO NOT APPLY DURIVO THROUGH ANY IRRIGATION SYSTEM **PHYSICALLY CONNECTED** TO A PUBLIC WATER SYSTEM. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Durivo may be applied through irrigation systems, which may be **supplied** by a public water system **only if** the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements.

Operating Instructions FOR DRIP IRRIGATION SYSTEMS

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Application Instructions

Durivo must be applied on the schedule specified in the **Crop Use Directions**, not according to the drip irrigation schedule. The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through drip type irrigation equipment. Users should check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemicals through irrigation equipment.

1. Each run of the irrigation system must be calibrated separately to determine the time it takes water to move through the system and to make sure all emitters in the system are putting out the same amount of water.
2. Only pressure injection or venturi equipment is recommended.
3. Determine the area to be treated in each irrigation run.
4. Measure the output of each of the emitters or drip tubes closest to and farthest from the injection point.

5. For calibration, substitute a concentrated detergent (such as Wisk) for the Durivo mixture in the injector (solution) tank. It is important to use the same volume of soap solution as the planned volume of Durivo solution when calibrating the system. The detergent will bubble as it leaves the emitters. The time period over which the bubbles occur should be checked for both the closest and farthest emitters. If these times are not within 2 minutes of each other, adjust the dilution ratio and/or the injection rate.

Step-by-Step Calibration and Application Instructions

1. Before starting to calibrate, operate the system until all of the emitters are putting out at equal flow rates, or until the system is operating at full pressure.
2. Make up an indicator solution of detergent or fertilizer, using the same rate of indicator as the planned volume of Durivo to be used in the mix.
3. Set the injector to apply the indicator solution at the injection rate to be used in the actual Durivo application.
4. Attach a 12-inch length of flexible tubing over the emitter closest to the injection point, another 12-inch length over the emitter farthest away. Both emitters should be monitored to determine the time intervals that the indicator solutions are observed.
5. Begin injecting the indicator solution. Direct the flow from the flexible tubes into a small container. Begin timing when the indicator solution is first detected. Stop timing when the indicator solutions are no longer detected in the container.
6. If the period of detection of the indicator solution between the near and far emitter is within 2 minutes, comparable coverage will be obtained. If they are not, make adjustments by increasing the dilution ratio, using more water per part of Durivo, or adjust the injector to a slower flow rate.
7. Once the system is calibrated, dilute the needed amount of Durivo with water and any other tank mix partners in the injection tank at a minimum dilution of 15 parts water to 1 part Durivo. Follow the directions for mixing and equipment set up in the **Mixing Procedures** section of this label for complete details.
8. Do not begin to inject Durivo into the system until all emitters are producing equal flow rates, or until the system is at full pressure.
9. Inject the Durivo solution into the system in the middle of the irrigation set in 1/2-1 inch of irrigation water.

MIXING PROCEDURES

Prepare no more injector tank mixture than is needed for the immediate operation. Thoroughly clean drip irrigation equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the drip irrigation operation. Do not let the injector tank mixture stand overnight in the injector tank. Flush the drip irrigation equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

Durivo Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Durivo to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Durivo has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Durivo + Tank Mixtures: Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, tank mix partners should be added in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables such as Durivo, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

Note: When using Durivo in tank mixtures, all products in water-soluble packaging should be added to the tank before any other tank mix partner, including Durivo. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using Durivo in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility: Durivo is compatible with most commonly used pesticides and nutritional sprays. However, since it is not possible to test all possible mixtures, the user should pre-test to insure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with Durivo. To determine the physical compatibility of Durivo with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables,

13/
12

and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the injector tank.

The crop safety of all potential tank mixes on all crops has not been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target crop should be confirmed.

M/22

CROP USE DIRECTIONS

Crop	Pest	Rate Per Acre Per Application
Brassica (Cole) Leafy Vegetables		
<p>Head & Stem Brassica Broccoli Broccoli, Chinese Brussels sprouts Cabbage Cabbage, Chinese (napa) Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi</p> <p>Leafy Brassica Greens Broccoli raab Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens</p>	<p>Aphids Beet Armyworm Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Southern Cabbageworm Thrips Whiteflies Yellowstriped Armyworm</p>	<p>10.0 – 13.0 fl. oz./A</p>

- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Methods:** Apply specified dosage in sufficient water volume to ensure uniform application and incorporation into the root zone using drip (trickle) chemigation.
- **Pre-Harvest Interval (PHI):** ~~30 Days~~
- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.

Refer to Resistance Management section.

15/100

Crop	Pest	Rate Per Acre Per Application
Cucurbit Vegetables Chayote Chinese waxgourd Citron melon Cucumber Edible gourd Gherkin <i>Momordica</i> species Muskmelon Pumpkin Squash: summer and winter Watermelon	Aphids Cucumber Beetles ¹ Flea Beetles Leafhoppers Leafminers ¹ Melonworm Pickleworm Thrips Whiteflies	10.0 – 13.0 fl. oz./A

- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to insure uniform application and incorporation into the root zone using drip (trickle) chemigation.
- **Pre-Harvest Interval (PHI):** 30 Days
- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.
- **Pest Control:** ¹Suppression.

Refer to Resistance Management section.

16/22

Crop	Pest	Rate Per Acre Per Application
Fruiting Vegetables Tomato Peppers: bell, chili, cooking, pimento, and sweet Eggplant Groundcherry Pepino Tomatillo	Aphids Beet Armyworm Colorado Potato Beetle Fall Armyworm Flea Beetles Hornworm species Leafhoppers Leafminers ¹ Loopers Potato Psyllid Southern Armyworm Thrips Tomato Fruitworm Tomato Pinworm Whiteflies Yellowstriped Armyworm	10.0 – 13.0 fl. oz./A

- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to insure uniform application and incorporation into the root zone using drip (trickle) chemigation.
- **Pre-Harvest Interval (PHI):** 30 Days
- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. ai of chlorantraniliprole containing products per acre per growing season.
- **Pest Control:** ¹Suppression

Refer to Resistance Management section.

17/22

Crop	Pest	Rate Per Acre Per Application
Leafy Vegetables Amaranth Arugula Cardoon Celery Celtuce Chinese celery Chrysanthemum: edible- leaved & garland Corn Salad Cress: Garden & upland Dandelion Dock Endive Fennel Lettuce: Head & Leaf Orach Parsley Purslane: Garden & Winter Radicchio Rhubarb Spinach: New Zealand & Vine Swiss chard	Aphids Beet Armyworm Cabbage Looper Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Leafhoppers Leafminers ¹ Whiteflies	10.0 – 13.0 fl. oz./A

- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to ensure uniform application and incorporation into the root zone using drip (trickle) chemigation.
- **Pre-Harvest Interval (PHI):** 30 Days
- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.
- **Pest Control:** ¹Suppression

Refer to Resistance Management section.

18/22

Durivo Conversion Chart for Drip Chemigation Application

	20"	30"	34"	36"	38"	40"	46"	60"	72"	78"	80"	84"	Row Spacing
	26,136	17,424	15,374	14,520	13,756	13,068	11,363	8,712	7,260	6,702	6,534	6,223	Linear Ft./A.
Rate (oz./A)	Rate in ounces. of product per 1,000 linear feet for specified row spacing and rate per acre												lb. a.i./A
10.0	0.38	0.57	0.65	0.69	0.73	0.77	0.88	1.15	1.38	1.49	1.53	1.61	0.195
11.0	0.42	0.63	0.72	0.76	0.80	0.84	0.97	1.26	1.52	1.64	1.68	1.77	0.215
12.0	0.46	0.69	0.78	0.83	0.87	0.92	1.06	1.38	1.65	1.79	1.84	1.93	0.234
13.0	0.50	0.75	0.85	0.90	0.95	0.99	1.14	1.49	1.79	1.94	1.99	2.09	0.254

ROTATIONAL RESTRICTIONS

Treated areas may be replanted immediately following harvest or as soon as practical following the last application with any of the crops listed on this label (*Brassica* leafy vegetables, cucurbit vegetables, fruiting vegetables, and leafy vegetables) or to potatoes or cotton. Members of the Root and Tuber vegetables (Crop Group 1) except sugarbeets may be planted back immediately. The following crops may be planted 30 days following the last application of Durivo: Legume vegetables (Crop Group 6), soybean, barley (ID, ND, OR, SD, and WA only) wheat, sorghum and strawberry. Any cover crop planted for erosion control or soil improvement may be planted as soon as practical following the last application. However, the cover crop may not be grazed or harvested for food or feed. For all other rotational crops intended for food or feed, a 12 month plant back interval must be observed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of unused pesticide, mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during

19/22

cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Container Disposal

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Residue Removal

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

DURIVO, the Syngenta logo, and the CP FRAME , are trademarks of a Syngenta Group Company

©2008 Syngenta

Viton® is a registered trademark of E.I. DuPont de Nemours and Company

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300
www.syngenta-us.com

SCP 1318A-M(draft)

DURIVO 1318A-M AMEND-B SEPT2008 CL – pd – 09/04/08
000100-01318.20080903B.DURIVO_ROTROP_AMEND_SEPT2008-CL.pdf

20/22

(Non-detachable container)

DURIVO™

GROUP	4A	INSECTICIDE
GROUP	28	INSECTICIDE

For control of listed insect pests infesting various crops

Active Ingredient:

Thiamethoxam¹ 17.5%
 Chlorantraniliprole² 8.8%

Other Ingredients: 73.7%

Total: 100.0%

¹CAS No. 153719-23-4

²CAS No. 500008-45-7

Durivo is a soluble concentrate containing 1.67 lb. of Thiamethoxam and 0.835 lb. Chlorantraniliprole per gallon.

KEEP OUT OF REACH OF CHILDREN.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard

EPA Reg. No. 100-1318

EPA Est. XXXXX

SCP 1318A-M(draft)

Net Contents

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) or
Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call

1-800-888-8372

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates, oysters and shrimp. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

~~Pesticide wastes are toxic. Improper disposal of unused pesticide, mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.~~

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Container Disposal

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Residue Removal

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Refer to Surface Water Advisory, Ground Water Advisory, Physical or Chemical Hazards and Chemigation sections in booklet.

DURIVO and the Syngenta logo are trademarks of a Syngenta Group Company
©2008 Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300
www.syngenta-us.com

SCP 1318A-M(draft)

DURIVO 1318A-M AMEND-B SEPT2008 CL - pd - 09/04/08
000100-01318.20080903B.DURIVO_ROTROP_AMEND_SEPT2008-CL.pdf
