

100-1347

07/18/2012

1/24



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Ms Adora Clark Ph D  
Syngenta Corp Protection LLC  
P O BOX 18300  
Greensboro NC 27419 8300

JUL 18 2012

Subject            Product Name Concert II  
                         EPA Reg No 100 1347  
                         Submission date 6/18/12  
                         Amendment Addition of the recently required California DPR Prop 65 Statement  
                         Decision Number 466637

Dear Registrant

The amendment referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act as amended is acceptable

One copy of the label stamped Accepted is enclosed for your records This label supersedes all labels previously accepted for this product Please submit one copy of the final printed label before the product is released for shipment

If you have questions concerning this letter please call Banza Djapao at 703 305 7269 or via email at djapao banza@epa gov or you may call me at 703 308 9443

Sincerely

Tony Kish  
Product Manager 22  
Fungicide Branch  
Registration Division (7504P)

**Concert® II**

**Fungicide**

Broad spectrum fungicide for control of plant diseases

*Active Ingredient*

Propiconazole (CAS No 60207 90 1)	2.9%
Chlorothalonil (CAS No 1897 45 6)	38.5%
<i>Other Ingredients</i>	58.6%
<b>Total</b>	<b>100.0%</b>

Concert II contains 0.3 lb a.i. propiconazole and 4.0 lb a.i. chlorothalonil per gallon

**KEEP OUT OF REACH OF CHILDREN**

**DANGER/PELIGRO**


Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

See additional precautionary statements and directions for use inside booklet

EPA Reg No 100-1347

EPA Est 50534 TX 001

2.5 gallons  
Net Contents



<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 20 minutes</li> <li>• Remove contact lenses if present after the first 5 minutes then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15 20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
<p>Probable mucosal damage may contraindicate the use of gastric lavage Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment</p>	
<b>HOT LINE NUMBER</b>	
<p>For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call <b>1 800 888 8372</b></p>	

---

**PRECAUTIONARY STATEMENTS**

---

**Hazards to Humans and Domestic Animals****DANGER/PELIGRO**

Corrosive Causes irreversible eye damage Do not get in eyes or on clothing Wear protective eye wear (goggles face shield or safety glasses) Harmful if swallowed Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

**Personal Protective Equipment (PPE)****Mixers loaders applicators and all other handlers must wear**

- Long sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material - Category A (e.g. barrier laminate butyl rubber nitrile rubber neoprene rubber natural rubber polyethylene polyvinyl chloride (PVC) or Viton®)
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate DO NOT reuse them Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

**Engineering Control Statements**

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

**User Safety Recommendations****Users should**

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

**Environmental Hazards**

Propiconazole is toxic to fish and chlorothalonil is toxic to aquatic invertebrates and wildlife Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff may be hazardous to aquatic organisms in neighboring area Do not contaminate water when disposing of equipment washwater or rinsate

Chlorothalonil is known to leach through soil into groundwater under certain conditions as a result of label use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination

Chlorothalonil can contaminate surface water through spray drift Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters frequently flooded areas areas over laying extremely shallow groundwater areas with infield canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips and areas over laying tile drainage systems that drain to surface water

Attention This product contains a chemical known to the State of California to cause cancer

---

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

---

**NOTICE** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Crop injury ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather or crop conditions presence of other materials or other influencing factors in the use of the product which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller To the extent consistent with applicable law Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions To the extent consistent with applicable law (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not

6/24

reasonably foreseeable to or beyond the control of Seller or SYNGENTA and (2) Buyer and User assume the risk of any such use TO THE EXTENT CONSISTENT WITH APPLICABLE LAW SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL

To the extent consistent with applicable law in no event shall SYNGENTA be liable for any incidental consequential or special damages resulting from the use or handling of this product **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT**

SYNGENTA and Seller offer this product and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL**

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

7/24

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material: Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC), or Viton®)
- Shoes plus socks
- Protective eyewear

**Special Eye Irritation Provisions:** This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6 days entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.
- (2) Workers must be informed in a manner they can understand:
  - that residues in the treated area may be highly irritating to their eyes
  - that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes
  - that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water
  - how to operate the eyeflush container

8/24

### **NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

**DO NOT enter or allow others to enter area until sprays have dried**

---

### **GENERAL INFORMATION**

Concert II is a combination of systemic and contact fungicides that provides broad spectrum control of many important plant diseases

#### **Integrated Pest (Disease) Management**

Concert II is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases Concert II is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM) which include the use of disease resistant crop varieties cultural practices pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides

#### **Resistance Management**

Concert II is effective for strategic use in programs that attempt to minimize disease resistance to fungicides Some other fungicides which are at risk from disease resistance exhibit a single site mode of fungicidal action Concert II with a multi site mode of action may be used to delay or prevent the development of resistance to single site fungicides Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of Concert II in programs which seek to minimize the occurrence of disease resistance to other fungicides

---

### **GENERAL USE PRECAUTIONS**

**Do not use in greenhouses**

**Do not use on home lawns and turf sites associated with apartment buildings daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie , elementary middle and high schools), campgrounds, churches and theme parks**



9/24

## Spray Drift Management

This product must not be applied within 25 feet of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body

Avoiding spray drift at the application site is the responsibility of the applicator

## Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas)

---

## APPLICATION INSTRUCTIONS

---

Apply Concert II at rates and timings as described in this label

- **DO NOT** apply by air
- **DO NOT** apply through any type of irrigation system
- Apply only by ground or hand held application equipment

## Spray Equipment

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to (1) maintain at least 35-40 psi at nozzles and (2) provide sufficient **continuous** agitation in tank to keep mixture in suspension. This requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

Concert II is a suspoemulsion concentrate. It is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be *16 mesh or coarser*. Do not place a screen in the recirculation line. Use 50 mesh or coarser screens between the pump and boom and where required at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state and local agricultural recommendations.

10/24

## **Spraying/Mixing**

Prepare no more spray mixture than is required for the immediate operation. Do not allow the spray mixture to sit overnight. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**Alone** Add  $\frac{1}{2}$   $\frac{2}{3}$  of the required amount of water to the spray or mixing tank. With the agitator running, add Concert II to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after Concert II has completely dispersed into the water. Maintain continuous agitation until all of the mixture has been sprayed.

**Concert II + Tank Mixtures** Consult your dealer or SYNGENTA sales representative for information concerning compatibility with other tank mix partners. NOTE: When using Concert II in mixture, any product in water-soluble packaging should be added to the tank prior to adding Concert II. Allow water-soluble packaging to completely dissolve and the mixture product to completely disperse before adding Concert II to the tank.

Add  $\frac{1}{2}$   $\frac{2}{3}$  of the required amount of water to the spray or mixing tank. Again, if tank mixing with any product packaged in water-soluble packets, add this product first with the agitator running. Allow the material to completely dissolve and disperse in the mix water. In general, additional tank mix partners should be added in the following order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable concentrations. Always allow each tank mix partner to become fully dispersed before adding the next product. Maintain agitation while adding the rest of the water while adding the other tank mix partners and throughout the mixing and spraying operation.

**Important** Do not combine Concert II with DiPel<sup>®</sup>, Latron B 1956<sup>®</sup>, or Latron AG 98<sup>®</sup> as phytotoxicity may result.

11/24

---

**TURF USE DIRECTIONS**

---

**Turf Restrictions**

- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields athletic fields located on or next to schools (ie elementary, middle and high schools) campgrounds churches and theme parks
- Sod farm turf treated with Concert II prior to harvest must be mechanically cut rolled and harvested
- Do not graze animals on treated areas Do not feed clippings from treated areas to livestock or poultry
- For rates **up to and including** 5.5 fl oz /1 000 sq ft (15 pts /A)
  - The minimum re treatment interval is 7 days
- For rates **greater than** 5.5 fl oz /1 000 sq ft (15 pts /A)
  - The minimum re treatment interval is 14 days
  - Limit of two applications per year on Tees and Greens
  - Limit of one application per year on All Other Turf
- Maximum single application rate of Concert II 8.5 fl oz /1 000 sq ft (23 pts /A)
- Maximum amount of Concert II per growing season

	Fl oz / 1,000 sq ft	Pts /A	Lbs a i /A Propiconazole	Lbs a i /A Chlorothalonil
Greens	53	146	5.5	73
Tees	38	104	3.9	52
Sod Farm	9.5	26	1.0	13
All Other Turf	19	52	2.0	26

- Do not use on sod farms at application rates greater than 13 lbs of chlorothalonil per acre per year
- Do not use Concert II on fine fescue turf due to the potential for phytotoxicity or turfgrass injury
- Do not water for 24 hours after treatment

- Bermudagrass can be sensitive to Concert II Do not exceed 6 fl oz /1 000 sq ft every 30 days on any variety of bermudagrass In FL do not apply Concert II to bermudagrass golf course greens when temperatures exceed 90 F

## **TURFGRASS AND DICHONDRA DISEASE CONTROL**

- 1 USE CONCERT II IN A PREVENTATIVE DISEASE CONTROL PROGRAM
- 2 Apply in sufficient water to ensure thorough coverage (30 to 40 gallons of water per acre)
- 3 Apply by ground or hand held applicators only
- 4 Apply after mowing **OR** allow sprayed area to completely dry before mowing
- 5 For control of foliar diseases allow sprayed area to completely dry before irrigation
- 6 For control of soil borne diseases Concert II can be watered in after application
- 7 Under conditions optimum for high disease pressure use the higher rate and the shorter interval
- 8 For optimum turf quality and disease control use Concert II in conjunction with turf management practices that promote good plant health and optimum disease control
- 9 Evaluate spray additives prior to use Label directions are based on data obtained with no additives
- 10 Before use of any fungicide proper diagnosis of the organism causing the disease is important Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures

**Turfgrass Specific Diseases, Rates, and Application Timing**

Disease	Fl oz per 1 000 Sq Ft	Pints per Acre	Application Interval/ Timing	Instructions
<b>Anthracnose</b> ( <i>Colletotrichum graminicola</i> )	4 5 5 5	12 5 15	7 14 days	Apply when conditions are favorable for disease development. When disease pressure is high use higher rates of Concert II and shorter intervals. For broad spectrum control tank mix with a registered contact fungicide at the label rate.
	5 5 8 5	15 23	14 28 days	
<b>Brown Patch</b> ( <i>Rhizoctonia solani</i> )	3 5 5	8 2 15	7 14 days	Begin applications in May or June before disease is present.  Under conditions of high temperatures and high humidity use the higher rates of Concert II and shorter intervals.
	5 5 8 5	15 23	14 28 days	
<b>Copper Spot</b> ( <i>Gloeocercospora sorghi</i> )	5 5 8 5	15 23	14 days	Apply when conditions are favorable for disease development.
<b>Dollar Spot</b> ( <i>Sclerotinia homoeocarpa</i> )	1 5 3	4 8 2	7 10 days	Apply when conditions are favorable for disease development.
	3 5 5	8 2 15	14 21 days	
	5 5 8 5	15 23	14 28 days	
<b>Fusarium Patch</b> ( <i>Microdochium nivale</i> )	8 5	23	14 28 days Fall Early Spring	Apply when conditions are favorable for disease development. A second application may be required.
<b>Gray Leaf Spot</b> ( <i>Pyricularia grisea</i> )	3 5 5	8 2 15	7 14 days	Apply when conditions are favorable for disease development.
	5 5 8 5	15 23	14 21 days	
<b>Melting Out Leaf Spot</b> ( <i>Bipolaris</i> spp ) ( <i>Drechslera</i> spp )	3 5 5	8 2 15	7 14 days	Apply when conditions are favorable for disease development.
	5 5 8 5	15 23	14 21 days	
<b>Powdery Mildew</b> ( <i>Erysiphe graminis</i> )	4 5 8 5	12 5 23	14 28 days	Apply when conditions are favorable for disease development. If disease is present use 8.5 fl. oz. of Concert II per 1 000 sq ft.
<b>Red Thread</b> ( <i>Laetisaria fuciformis</i> ) <b>Pink Patch</b> ( <i>Limonomyces roseipellis</i> )	3 5 5	8 2 15	7 14 days	Apply when conditions are favorable for disease development.
	5 5 8 5	15 23	14 21 days	
<b>Rust</b> ( <i>Puccinia graminis</i> )	4 5 8 5	12 5 23	14 28 days	Apply when conditions are favorable for disease development. If disease is present use 8.5 fl. oz. of Concert II per 1 000 sq ft.

14/24

Disease	Fl oz per 1 000 Sq Ft	Pints per Acre	Application Interval/ Timing	Instructions		
<b>Snowmold</b>	5.5	15	14 days	Apply in the late fall before snow cover in enough carrier to obtain adequate coverage (2.8 gallons per 100 sq ft). Do not apply on top of snow. For optimum disease control tank mix with the label rate of another snow mold registered fungicide label rates. The 5.5 fl oz rate can be repeated once at 14 days. Make only one application at rates above 6 fl oz /1 000 sq ft. Tees and greens if snow cover is intermittent or lacking during the winter you may make a second application.		
<b>Gray</b> ( <i>Typhula</i> spp.) <b>Pink</b> ( <i>Microdochium nivale</i> )	8.5	23	28 days Late Fall			
<b>Stem Rust (Bluegrass)</b> ( <i>Puccinia graminis</i> )	3.5	8.2	15 days	Apply in early fall or when conditions are favorable for disease development.		
	5.5	8.5	15		23	14 days
<b>Stripe Smut</b> ( <i>Ustilago striiformis</i> ) ( <i>Urocystis agropyri</i> )	4.5	8.5	12.5	23	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
<b>Summer Patch</b> <b>Poa Patch</b> ( <i>Magnaporthe poae</i> )	8.5	23	14 days	Apply Concert II beginning in April. Use the 8.5 fl oz /1 000 sq ft rate on a 14 day schedule.		
<b>Take All Patch</b> ( <i>Gaeumannomyces graminis</i> )	8.5	23	14 days Spring and Fall	Apply Concert II to reduce the severity of take all patch. Make 1-2 fall applications in September and October or when night temperatures drop to 55 F and 1-2 spring applications in April and May depending on local recommendations.		

**Dichondra Specific Disease, Rate and Application Timing**

Disease	Fl oz per 1 000 Sq Ft	Pints per Acre	Application Interval/ Timing	Instructions			
<b>Dichondra Rust</b> ( <i>Puccinia dichondrae</i> )	5.5	8.5	15	23	14	21 days	Apply when conditions are favorable for disease development.
<b>Dichondra Leaf Spot</b>	4.5	5.5	12.5	15			

15/24

**Other Pests Rate and Application Timing**

Disease	Fl oz per 1 000 Sq Ft	Pints per Acre	Application Interval/ Timing	Instructions
Algae	3 5 5	8 2 15	7 14 days	Apply preventively when conditions are favorable for development
	5 5 8 5	15 23	14 28 days	

**Concert II Rate Conversion Table**

Fl oz per 1,000 sq ft	Pints/A	Lbs a i /A Propiconazole	Lbs a i /A Chlorothalonil	Treated Acres/ Gal Product
1 5	4	0 15	2	2
3	8 2	0 3	4 25	1
4 5	12 5	0 47	6 25	0 7
5 5	15	0 56	7 5	0 5
7	19	0 7	9 5	0 4
8 5	23	0 86	11 5	0 35

**ESTABLISHMENT OF COOL SEASON TURFGRASS**

Concert II provides control of many diseases of turf and its primary use is as a fungicide for use against the diseases listed on this label. As an additional benefit Concert II will improve the rate of establishment when it is applied to cool season grass seedlings or sod.

**New Seedlings** Apply 4 5 fl oz /1 000 sq ft at the 2 to 3 leaf stage of growth for faster root development and top growth.

**Sod** Apply 4 5 fl oz /1 000 sq ft 2 6 weeks before cutting for increased sod knitting and faster establishment after laying.

**DISEASE CONTROL IN NURSERIES (FIELD) AND LANDSCAPE PLANTINGS**

- 1 USE CONCERT II IN A PREVENTATIVE DISEASE CONTROL PROGRAM. To determine the use directions for controlling a disease on an ornamental plant species, select the plant species in Table 1. The number following the plant species refers to the disease(s) controlled in Table 2. Find the disease in Table 2. The letter following the disease refers to the application regime in Table 3.
- 2 Allow spray to dry before overhead irrigation is applied.

- 3 Optimum benefit of Concert II is obtained when used in conjunction with sound disease management practices

### Restrictions

- Do not use in greenhouses
- Do not use mist blowers high pressure spray equipment cold fog or other fogging application equipment when making applications of Concert II
- For use on non-bearing fruits and nuts only (no harvestable fruits within 12 months)
- Refer to Concert II Rate Conversion Table for maximum gallons per acre

### General Recommendations

Concert II may be used for control of diseases of ornamental plant species (see Tables 1 2 and 3)

Apply by ground or hand held applicators only For general disease control in landscapes apply 26 35 fl oz /100 gals water every 21 days For best control begin Concert II applications before disease development

### Plant Safety

- Before using Concert II on ornamentals test Concert II on a small scale basis To determine if conditions may result in damage apply Concert II to a small number of plants under the growing conditions and timing that you would apply Concert II and observe for two weeks before treating the entire crop
- Knock Out® and Double Delight roses can be sensitive to Concert II applications resulting in damage to foliage under certain growing conditions
- Do not apply Concert II to African violets begonias Boston fern geraniums variegated or green Pittosporum or Schefflera
- Applications of Concert II may result in injury to buds blooms or tender new growth Avoid applications during bloom period on plants where flower injury is unacceptable
- Addition of surfactants wetters spreader sticker or other pesticides or fertilizers should be tested on a small scale basis



**Table 1**

<b>Ornamental Plant Species</b>	<b>Disease*</b>
<b>Herbaceous Ornamental</b>	
Calendula	4a
Carnation	5e
Chrysanthemum	2a
Delphinium	4a
Gomphrena	3a
Iris	5d
Marigold	3a
Monarda	4c
Phlox	4c
Sweet William ( <i>Dianthus barbatus</i> )	3i
Zinnia	4c
<b>Woody Ornamental</b>	
Amelanchier	4d
Ash	4c
Azalea	2c 4b
Crabapple	3b 3i 4c 5a
Crape Myrtle	4a
Dogwood	3f 4c
Hawthorn	5a
Lilac	4c
Oaks	3k
Poplars	5b
Pyracantha	3j
Red Tip Photinia	3g
Rhododendron	2c
Roses (Outdoor Use Only)	3e 5b
A maximum of 9.0 gallons of Concert II may be applied/acre/crop/year Do not apply more than 36.0 lbs a.i. chlorothalonil per acre per year Do not apply more than 7.00 lbs a.i. propiconazole per acre per year	
<b>Pines</b>	
Pines	1a 1b
A maximum of 4.1 gallons of Concert II may be applied/acre/crop/year Do not apply more than 16.5 lbs a.i. chlorothalonil per acre per year Do not apply more than 7.00 lbs a.i. propiconazole per acre per year	
<b>Non-bearing Fruits and Nuts (Nurseries (Field) and Landscape Plantings)</b>	
Apple	3l 4d 5a
Apricot	2b
Cherry	2b 3c
Nectarine	2b
Peach	2b
Plum	2b
Walnut	3h
A maximum of 3.8 gallons of Concert II may be applied/acre/crop/year Do not apply more than 15.5 lbs a.i. chlorothalonil per acre per year Do not apply more than 7.0 lbs a.i. propiconazole per acre per year	

Numbers refer to diseases controlled in Table 2

18/24

**NOTICE TO USER** Plant tolerance to Concert II has been found to be acceptable for the specific genera and species listed on this label. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to Concert II. Neither the Manufacturer nor the Seller has determined whether or not Concert II can be used safely on ornamental plants not specified on this label. The professional user should determine if Concert II can be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use.

**Table 2**

Disease/Pathogen		Application Regime*
<b>1</b>	<b>Conifer Blights</b>	
1a	<i>Sirococcus strobilinus</i> (Tip Blight)	C
1b	<i>Sphaeropsis sapinea</i> (Diplodia Tip Blight)	B
<b>2</b>	<b>Flower Blight</b>	
2a	<i>Ascochyta chrysanthemi</i> (Ray Blight)	B
2b	<i>Monilinia</i> spp	A
2c	<i>Ovulinia</i> spp	B
<b>3</b>	<b>Leaf Blights/Spots</b>	
3a	<i>Alternaria</i> spp	B
3b	<i>Cladosporium</i> spp (Scab)	B
3c	<i>Coccomyces hiemalis</i>	A
3d	<i>Collectrichum</i> spp	B
3e	<i>Diplocarpon rosae</i> (Blackspot)	B
3f	<i>Discula</i> spp (Anthracnose)	A
3g	<i>Fabraea maculata</i> (syn <i>Entomosporium maculata</i> )	B
3h	<i>Gnomonia leptostyla</i> (Anthracnose)	B
3i	<i>Heterosporium echinulatum</i>	B
3j	<i>Spilocaea pyracanthae</i>	B
3k	<i>Tubakia dryina</i>	C
3l	<i>Venturia inaequalis</i> (Scab)	A
<b>4</b>	<b>Powdery Mildew</b>	
4a	<i>Erysiphe</i> spp	B
4b	<i>Microsphaera</i> spp	B
4c	<i>Oidium</i> spp	B
4d	<i>Podosphaera</i> spp	B
<b>5</b>	<b>Rust</b>	
5a	<i>Gymnosporangium juniper virginianae</i>	A
5b	<i>Melampsora occidentalis</i>	C
5c	<i>Phragmidium</i> spp	B
5d	<i>Puccinia</i> spp	B
5e	<i>Uromyces dianthi</i>	B

Letters refer to application regimes in Table 3

19/24

**Table 3**

Application Regime	Directions
A	Mix 9 17 fl oz (0 6 1 1 pts ) of Concert II in 100 gals of water and apply as a full coverage spray to the point of drip Apply every 14 21 days during the period of primary infection For dogwoods apply the 9 17 fl oz rate every 14 days or apply 35 fl oz of Concert II every 28 days
B	Mix 22 35 fl oz (1 4 2 2 pts ) of Concert II in 100 gals of water and apply as a full coverage spray to the point of drip Apply as needed beginning when conditions are favorable for disease development For blackspot apply with a registered contact fungicide labeled for blackspot For Calendula apply every 30 days For Diplodia tip blight make 3 applications every 14 days prior to major period of infection For juniper Phomopsis blight make first application as soon as junipers start to grow and repeat the applications every 14 21 days during periods of active growth
C	Mix 69 fl oz (4 3 pts ) of Concert II in 100 gals of water and apply as a full coverage spray to the point of drip Apply every 14 28 days beginning when conditions are favorable for disease development Do not make more than 3 applications For Douglas fir needle rust apply once in May For tip blight initial application is in mid late winter apply 3 times every 60 days

**Concert II Rate Conversion Table**

Fl oz / 100 Gallons	Pints/ 100 Gallons	Lbs a.i. Propiconazole/ 100 Gallons	Lbs a.i. Chlorothalonil/ 100 Gallons	Maximum Gallons/ A
9	0 6	0 02	0 28	600
17	1 1	0 04	0 53	600
22	1 4	0 05	0 69	450
35	2 2	0 08	1 1	280
69	4 3	0 16	2 2	140

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**Pesticide Storage**

Store in a cool dry place

**Pesticide Disposal**

Pesticide wastes are toxic Improper disposal of unused pesticide spray mixture or rinse water is a violation of Federal law If these wastes cannot be used according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods

20/24

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

**Container Disposal**

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple-rinse container (or equivalent) promptly after emptying. Triple-rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER**

---

Concert® II, the Syngenta logo and the CP FRAME  are trademarks of a Syngenta Group Company.

DiPel® is a registered trademark of Valent BioSciences Corporation.

Knock Out® is a registered trademark of CP, Inc.

Latron B 1956® and Latron AG 98® are registered trademarks of Dow AgroSciences LLC.

Viton® is a trademark of E. I. duPont de Nemours and Company.

©20\_\_ Syngenta

For non-emergency (e.g., current product information) call  
Syngenta Crop Protection at 1-800-334-9481

Manufactured for  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

2/24

BASE LABEL

**Concert® II**

**Fungicide**

Broad spectrum fungicide for control of plant diseases

Active Ingredient

Propiconazole (CAS No 60207 90 1) 2.9%

Chlorothalonil (CAS No 1897 45 6) 38.5%

Other Ingredients 58.6%

Total 100.0%

Concert II contains 0.3 lb a.i. propiconazole and 4.0 lb a.i. chlorothalonil per gallon

**KEEP OUT OF REACH OF CHILDREN**

**DANGER/PELIGRO**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

See additional precautionary statements and directions for use inside booklet

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. Refer to supplemental labeling under Agricultural Use Requirements in the Directions for Use section for information about this standard.

EPA Reg No 100 1347

EPA Est 50534 TX 001

2.5 gallons  
Net Contents

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>• Remove contact lenses if present after the first 5 minutes then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15 20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth-to-mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
<p>Probable mucosal damage may contraindicate the use of gastric lavage Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment</p>	
<b>HOT LINE NUMBER</b>	
<p>For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call <b>1 800 888 8372</b></p>	

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**DANGER/PELIGRO**

Corrosive Causes irreversible eye damage Do not get in eyes or on clothing Wear protective eye wear (goggles face shield or safety glasses) Harmful if swallowed Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

**Environmental Hazards**

Propiconazole is toxic to fish and chlorothalonil is toxic to aquatic invertebrates and wildlife Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff may be hazardous to aquatic organisms in neighboring area Do not contaminate water when disposing of equipment washwater or rinsate

Chlorothalonil is known to leach through soil into groundwater under certain conditions as a result of label use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination

Chlorothalonil can contaminate surface water through spray drift Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters frequently flooded areas areas over laying extremely shallow groundwater areas with infield canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips and areas over laying tile drainage systems that drain to surface water

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**Pesticide Storage**

Store in a cool dry place

**Pesticide Disposal**

Pesticide wastes are toxic. Improper disposal of unused pesticide spray mixture or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1 800 888 8372 day or night.

**Container Disposal**

Non-refillable container: Do not reuse or refill this container. Offer for recycling if available. Triple-rinse container (or equivalent) promptly after emptying. Triple-rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER**

---

Concert® II and the Syngenta logo are trademarks of a Syngenta Group Company

©20\_\_ Syngenta

Manufactured for  
Syngenta Crop Protection, LLC  
P O Box 18300  
Greensboro, North Carolina 27419 8300

CONCERT II 1347 MAS 0509 AMEND 0612 CLEAN – sw – 7/18/12  
000100 01347 20120615 CONCERT\_II AMEND 0612 CLEAN PDF

