

100-1384

8/3/2012

1/30

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

AUG - 3 2012

Patrick McCain
Syngenta Crop Protection, Inc
P O Box 18300
Greensboro, NC 27419

Dear Mr McCain

Subject Amended label to add legume group
Product Name Four-way VAP™
EPA Registration No 100-1384
EPA Decision No 467643

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable. A stamped copy of the label is enclosed for your records. Please submit two copies of your final printed labeling before you release the product for shipment. Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions, please contact Julie Chao at (703) 308-8735 or chao.julie@epa.gov

Regards,

A handwritten signature in black ink that reads "Venus Eagle".

Venus Eagle, Product Manager (01)
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

Four-way VAP™

GROUP	4A	INSECTICIDE
GROUP	4 11 12	FUNGICIDES

Insecticide with Three Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on legume vegetables (including soybeans)

Active Ingredients:

Thiamethoxam*	22.60%
Mefenoxam**	1.70%
Fludioxonil***	1.12%
Azoxystrobin****	0.90%
Other Ingredients:	73.68%
Total:	100.00%

*CAS No. 153719-23-4

**CAS No. 70630-17-0 and CAS No. 69516-34-3

***CAS No. 131341-86-1

****CAS No. 131860-33-8

One gallon of Four-way VAP contains 2.13 lb. thiamethoxam, 0.16 lb. mefenoxam, 0.11 lb. fludioxonil and 0.08 lb. azoxystrobin.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1384

EPA Est. XXXXX

SCP 1384A-M(draft-b legumes)

Net Contents

ACCEPTED
AUG - 3 2012
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 100-1384

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible • Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <i>[1-800-888-8372] [Alternate phone number]</i>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals**CAUTION**

Causes moderate eye irritation Harmful if swallowed Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet Avoid contact with eyes, skin or clothing

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material – Category C (e.g., barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride [PVC] ≥14 mils or Viton® ≥14 mils)
- Shoes plus socks

Follow manufacturer’s instructions for cleaning/maintaining PPE If no such instructions exist for washables, use detergent and hot water Keep and wash PPE separately from other laundry

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Recommendations

Users should

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment wash water.

Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops. Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Azoxystrobin in this product can be persistent for several months or longer. Azoxystrobin degradation products and thiamethoxam have properties and characteristics similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use. Use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training, decontamination, notification, and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours Exception If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is

- Coveralls
- Chemical-resistant gloves made of any waterproof material – Category C (e g , barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride [PVC] ≥14 mils or Viton® ≥14 mils)
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with Four-way VAP.

USE INFORMATION

Four-way VAP is a seed treatment product containing the active ingredients thiamethoxam (insecticide) and azoxystrobin, fludioxonil and mefenoxam fungicides. Four-way VAP protects against damage from listed early season insects, soil-borne and seed-borne diseases of crop plants.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam protects against listed chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against *Pythium*, *Phytophthora* and systemic downy mildew.

Fludioxonil fungicide is active against *Fusarium*, *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Azoxystrobin is active against *Rhizoctonia* and some seed-borne diseases.

MIXING PROCEDURES

Important Always re-circulate Four-way VAP thoroughly before using.

Apply Four-way VAP as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Four-way VAP into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. Four-way VAP is compatible with several liquid inoculant products. Consult the maker of the

9/30

inoculant product and a Syngenta Crop Protection representative for directions before applying Four-way VAP with inoculants

Under certain disease conditions, additional amounts of fungicides may be required. When needed, apply additional Apron XL® according to the **CROP USE DIRECTIONS**. Other tank mix partners may be used with Four-way VAP, however, the user must consider the use rate, formulation, seed and crop safety factors and compatibility of each product to be mixed when determining the total application volume.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. The minimum slurry volume to achieve adequate coverage is 4.0 fluid ounces per 100 pounds of seed. More diluent may be required to obtain complete coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging.

The typical density of Four-way VAP is 9.43 pounds per gallon. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

Four-way VAP contains an EPA approved dye/colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155 (c).

USE RESTRICTIONS

- Store away from feeds and foodstuffs
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Forage may not be grazed until 30 days after planting.
- Do not allow children, pets, or livestock to have access to treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.

- Dispose of all excess treated seed Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements
 - Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used for agronomic practice
- Do not use at a rate that will result in more than 0.266 lb thiamethoxam per acre (120.66 grams a /A) per year
- With the exception of soybeans do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Four way VAP For soybeans do not apply a neonicotinoid insecticide within 45 days of planting seed treated with Four way VAP

ROTATIONAL RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from Four way VAP treated seed the field may be replanted immediately to alfalfa *Brassica* (cole) leafy vegetables cereal grains (including barley buckwheat corn pearl millet proso millet oats popcorn rice (dry seeded) rye sorghum teosinte triticale wheat and wild rice) canola cotton cucurbit vegetables dry bulb onions fruiting vegetables leafy vegetables legume vegetables (including soybeans) mint (peppermint and spearmint) oil seed crops (rapeseed Indian rapeseed Indian mustard seed field mustard seed black mustard seed flax seed safflower seed crambe seed and borage seed) root vegetables strawberry sunflowers tobacco and tuberous and corm vegetables For any other crop the minimum plant back interval is 120 days from the date Four way VAP treated seed was planted A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval however the crop may not be grazed or harvested for food or feed

SEED BAG LABEL REQUIREMENTS

Federal law requires that bags containing treated seeds shall be labeled with the following statements

- This seed has been treated with thiamethoxam insecticide and azoxystrobin fludioxonil and mefenoxam fungicides
- Do not use for feed food or oil purposes
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act

In addition the U S Environmental Protection Agency requires the following statements on bags containing seeds treated with Four way VAP

- Ground Water Advisory
Thiamethoxam has properties and characteristics associated with chemicals detected in ground water This chemical may leach into the ground water if used in areas where soils are permeable particularly where the water table is shallow Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination
Azoxystrobin in this product can be persistent for several months or longer Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use Use of azoxystrobin in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination
- Pollinator Precautions Thiamethoxam is highly toxic to bees and effects are possible as a result of exposure to translocated residues in blooming crops
- Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used for agronomic practice
- Store away from feeds and foodstuffs
- Wear long sleeved shirt long pants and chemical resistant gloves when handling treated seed
- Treated seeds exposed on soil surface may be hazardous to wildlife Cover or collect treated seeds spilled during loading
- Do not contaminate water bodies when disposing of planting equipment wash waters
- Dispose of seed packaging in accordance with local requirements
- In the event of a crop failure or harvest of a crop grown from Four way VAP treated seed the field may be replanted immediately to alfalfa *Brassica* (cole) leafy vegetables cereal grains (including barley buckwheat corn pearl millet proso millet oats popcorn rice (dry seeded) rye sorghum teosinte triticale wheat and wild rice) canola cotton cucurbit vegetables dry bulb onions fruiting vegetables leafy vegetables legume vegetables (including soybeans) mint (peppermint and spearmint) oil seed crops (rapeseed Indian rapeseed Indian

12/30

mustard seed field mustard seed black mustard seed flax seed safflower seed
crambe seed and borage seed) root vegetables strawberry sunflowers
tobacco and tuberous and corm vegetables For any other crop the minimum
plant back interval is 120 days from the date Four way VAP treated seed was
planted A cover crop other than the crops listed above that is planted for
erosion control or soil improvement may be planted sooner than the 120 day
interval however the crop may not be grazed or harvested for food or feed

- Forage may not be grazed until 30 days after planting
- Do not allow children pets or livestock to have access to treated seed
- Treated seed must be planted into the soil at a depth greater than 1 inch
- Dispose of all excess treated seed Leftover treated seed may be doublesown
around the headland or buried away from water sources in accordance with local
requirements
- Do not use at a rate that will result in more than 0.266 lb thiamethoxam per acre
(120.66 grams a /A) per year
- This seed has been treated with ___mg ai thiamethoxam per seed
- With the exception of soybeans do not make any soil or foliar application of
products containing thiamethoxam to crops grown from seed treated with Four
way VAP For soybeans do not apply a neonicotinoid insecticide within 45 days
of planting seed treated with Four way VAP

CROP USE PRECAUTIONS

Resistance Management

Four way VAP contains thiamethoxam a Group 4A insecticide fludioxonil a Group 12 fungicide mefenoxam a Group 4 fungicide and azoxystrobin a Group 11 fungicide

Some insect pests are known to develop resistance to products after repeated use Because resistance development cannot be predicted the use of this product should conform to sound resistance management strategies established for the crop and use area Syngenta encourages responsible product stewardship to ensure effective long term control of the insects on this label

If resistance to this product develops in your area this product or other products with a similar mode of action may not provide adequate control If poor performance cannot be attributed to improper application or extreme weather conditions a resistant strain of insect may be present If you experience difficulty with control and resistance is a reasonable cause immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area

Four way VAP contains a Group 4A insecticide (thiamethoxam belonging to the neonicotinoid class of chemistry) Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species This may result in partial or total loss of control of those species by Four way VAP or other Group 4A insecticides

In order to maintain susceptibility to this class of chemistry

- Avoid using Group 4A insecticides exclusively for season long control of insect species with more than one generation per crop season
- For insect species with successive or overlapping generations apply Four way VAP or other Group 4A insecticides using a treatment window approach A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern Within the treatment window depending on the length of residual activity there may either be single or consecutive applications (seed treatment soil foliar unless otherwise stated in the Directions for Use) of the Group 4A insecticides Do not exceed the maximum Four way VAP allowed per growing season
- Following a treatment window of Group 4A insecticides rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 4A insecticides
- A treatment window rotation along with other IPM practices for the crop and use area is considered an effective strategy for preventing or delaying a pest s ability to develop resistance to this class of chemistry
- If resistance is suspected do not reapply Four way VAP or any other Group 4A insecticides

Other Insect Resistance Management (IRM) practices include

- Incorporating IPM techniques into your insect control program
- Monitoring treated insect populations for loss of field efficacy
- Using tank mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied

For additional information on Insect Resistance Management

- Contact your local extension specialist certified crop advisor and/or product manufacturer for additional insect resistance management recommendations
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>

Four way VAP contains the fungicide mefenoxam a systemic fungicide having a specific mode of action against fungal pathogens

Note The fungicide mefenoxam contained in this product could be subject to development of insensitive strains of fungi or may be ineffective against naturally occurring strains of fungi. Development of insensitivity or natural tolerance cannot be predicted. Therefore [Syngenta][company name] cannot assume liability for crop damage resulting from insensitive or tolerant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible insensitive or tolerant strains of fungi which may occur.

CROP USE DIRECTIONS

When applied according to the **FOUR WAY VAP RATE TABLE** Four way VAP provides early season protection against injury by aphids bean leaf beetle grape colaspis leaf miners leaf hoppers Mexican bean beetle seed corn maggot threecornered alfalfa hopper thrips white grubs and wireworm
Four way VAP provides protection against damping off and seed borne rots due to *Pythium Phytophthora Fusarium Rhizoctonia* species and early season *Phytophthora* root rot Four way VAP also suppresses seed borne *Sclerotinia* and *Phomopsis* species

Legume Vegetable Group

- Bean (All *Lupinus* species)** (includes grain sweet white white sweet lupin)
- Bean (All *Phaseolus* species)** (includes black bean cranberry bean field bean great Northern bean kidney bean lima bean navy bean pinto bean runner bean small red bean snap bean tepary bean wax bean yellow bean)
- Bean (All *Vigna* species)** (includes adzuki bean asparagus bean blackeyed pea catjang Chinese longbean cowpea Crowder pea moth bean mung bean rice bean southern pea urd bean yardlong bean)
- Broad bean (fava bean)**
- Chickpea (garbanzo bean)**
- Guar**
- Jackbean**
- Lablab bean (hyacinth bean)**
- Lentil**
- Pigeon pea**
- Soybean** (including soybean vegetable)
- Sword bean**

When to add additional Apron XL®

If target fields have a history of high *Phytophthora* pressure add additional Apron XL as directed in the rate table and the Apron XL label The additional Apron XL may reduce compatibility with some rhizobia inoculants Consult with the maker of rhizobia inoculants before adding the additional Apron XL

FOUR WAY VAP RATE TABLE LEGUME VEGETABLES

Crop	Rate of Four way VAP		Additional Apron XL	
	fl oz per 100 lb seed	grams ai per 100 kg seed	fl oz per 100 lb seed	grams ai per 100 kg seed
Bean(All Lupinus species) including grain sweet white white sweet lupin Bean(All Phaseolus species including) black bean cranberry bean field bean great Northern bean kidney bean lima bean navy bean pinto bean runnerbean snap bean small red bean tepary bean wax bean yellow bean (continued)	3.0 fl oz per 100 lb seed	Thiamethoxam 50 Azoxystrobin 2 Mefenoxam 3.75 Fludioxnil 2.5	0.16 0.48 fl oz	3.75 11.25 gm

NOTE

- Under cool conditions at planting azoxystrobin has been documented in some geographies to delay emergence of certain varieties within the Legume Vegetable Group
- Due to the large number of varieties within the Legume Vegetable Group some varieties have not been evaluated for seed safety. If the Four way VAP seed safety of a particular variety is unknown a seed safety germination study should be conducted prior to applying Four way VAP
- When treated according to the directions for post planting protection against listed pests Four way VAP will also provide protection during post treatment storage of the seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*) Rice Weevil (*Sitophilus oryzae*) Red Flour Beetle (*Tribolium castaneum*) and Lesser Grain Borer (*Rhizopertha dominica*)
- If seed to be treated has existing infestations of stored grain insects it is recommended that the seed be fumigated prior to treating with Four way VAP and bagging

FOUR-WAY VAP RATE TABLE-LEGUME VEGETABLES* (continued)

Crop	Rate of Four-way VAP		Additional Apron XL	
	fl oz per 100 lb seed	grams ai per 100 kg seed	fl oz per 100 lb seed	grams ai per 100 kg seed
Bean(All <i>Vigna</i> species)including: adzuki bean asparagus bean blackeyed pea catjang Chinese longbean cowpea Crowder pea moth bean mung bean rice bean southern pea urd bean yardlong bean Broad bean (fava bean) Chickpea (garbanzo bean) Guar Jackbean Lalab bean (hyacinth bean) Lentil Pigeon pea Sword bean	3.0 fl oz per 100 lb seed	Thiamethoxam 50 Azoxystrobin 2 Mefenoxam 3.75 Fludioxinil 2.5	0.16 - 0.48 fl oz	3.75 - 11.25 gm

NOTE:

- Under cool conditions at planting, azoxystrobin has been documented in some geographies to delay emergence of certain varieties within the Legume Vegetable Group.
- Due to the large number of varieties within the Legume Vegetable Group, some varieties have not been evaluated for seed safety. If the Four-way VAP seed safety of a particular variety is unknown, a seed safety germination study should be conducted prior to applying Four-way VAP.
- When treated according to the directions for post-planting protection against listed pests, Four-way VAP will also provide protection during post treatment storage of the seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).
- If seed to be treated has existing infestations of stored grain insects, it is recommended that the seed be fumigated prior to treating with Four-way VAP and bagging.

FOUR WAY VAP RATE TABLE SOYBEAN

Crop	Rate of Four way VAP				Additional Apron XL	
	fl oz per 100 lb seed or fl oz per 140 000 seeds	grams ai per 100 kg seed	mg ai per seed	ml per 1 000 seeds	fl oz per 100 lb seed or gm ai per 100 kg	mg ai per seed or fl oz per 1 000 seeds or fl oz per 140 000 seeds
Soybean including soybean vegetable	3.0 fl oz per 100 lb seed or 1.40 fl oz per 140 000 seeds	Thiamethoxam 50 Azoxystrobin 2 Mefenoxam 3.75 Fludioxinil 2.5	Thiamethoxam 0.0762 Azoxystrobin 0.003 Mefenoxam 0.0057 Fludioxinil 0.0039 Total = 0.0858	0.2957	0.16048 fl oz per 100 lb seed or 3.751125 gm ai per 100 kg seed	0.005700170 mg ai per seed or 0.0005300016 fl oz per 1000 seeds or 0.07460224 fl oz per 140 000 seeds

The mg ai per seed ml Four way VAP per 1 000 seeds and fl oz Four way VAP per 140 000 seeds rates are based on 3 000 seeds per pound

NOTE

- When treated according to the directions for post planting protection against listed pests Four way VAP will also provide protection during post treatment storage of the seed against damage from the following stored grain insects Indian Meal Moth (*Plodia interpunctella*) Rice Weevil (*Sitophilus oryza*) Red Flour Beetle (*Tribolium castaneum*) and Lesser Grain Borer (*Rhizopertha dominica*)
- If seed to be treated has existing infestations of stored grain insects it is recommended that the seed be fumigated prior to treating with Four way VAP and bagging

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage disposal or cleaning of equipment

Pesticide Storage

Store in a cool dry place Storage for extended periods above 90 F is not recommended

For minor spills leaks etc follow all precautions indicated on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility If these wastes cannot be used according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods

Container Handling *[5 gallons or less]*

Non refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Container Handling *[greater than 5 gal non refillable]*

Non refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container ¼ full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Container Handling *[greater than 5 gal refillable]*

Refillable container Refill this container with pesticide only Do not use this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill by incineration or by other procedures allowed by state and local authorities

Apron XL® the ALLIANCE FRAME the Syngenta Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company
Viton® is a registered trademark of E I DuPont de Nemours and Company
©2012 Syngenta

[optional]

For non emergency (e g current product information) call
*[Syngenta Crop Protection at 1 800 334 9481][Company name and optional alternate
phone number]*

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro North Carolina 27419 8300
SCP 1384A M(draft b legumes)
Four way VAP 1384A M(draft -b legumes)clean lg 8 1 12 000100 01384 20120801 b legumes pdf

(Container Label – 5 gal or less)

Four way VAP™

GROUP	4A	INSECTICIDE
GROUP	4 11 12	FUNGICIDES

Insecticide with Three Fungicides

A seed treatment product for protection against damage from listed insects seed borne diseases and seedling diseases on legume vegetables (including soybeans)

Active Ingredients

Thiamethoxam*	22.60%
Mefenoxam**	1.70%
Fludioxonil***	1.12%
Azoxystrobin****	0.90%
Other Ingredients	73.68%
Total	100.00%

*CAS No 153719 23 4

**CAS No 70630 17 0 and CAS No 69516 34 3

***CAS No 131341-86 1

****CAS No 131860 33 8

One gallon of Four way VAP contains 2.13 lb thiamethoxam 0.16 lb mefenoxam 0.11 lb fludioxonil and 0.08 lb azoxystrobin

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use in booklet

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 Refer to supplemental labeling under Agricultural Use Requirements in the Directions for Use section for information about this standard

EPA Reg No 100 1384

EPA Est XXXXX

SCP 1384A M(draft b legumes)

Net Contents

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation Harmful if swallowed Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet Avoid contact with eyes skin or clothing

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call [1 800 888 8372] [Alternate phone number]	

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment wash water.

Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops.

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Azoxystrobin in this product can be persistent for several months or longer.

Azoxystrobin degradation products and thiamethoxam have properties and characteristics similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use. Use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and

drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

The Syngenta Logo is Trademarks of a Syngenta Group Company
©2012 Syngenta

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro North Carolina 27419 8300

SCP 1384A M(draft b legumes)

Four way VAP 1384A M(draft b legumes)clean lg 8 1 12 000100 01384 20120801 b legumes pdf

(Container Label -greater than 5 gal)

Four-way VAP™

GROUP	4A	INSECTICIDE
GROUP	4	11 12
FUNGICIDES		

Insecticide with Three Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on legume vegetables (including soybeans)

Active Ingredients:

Thiamethoxam*	22.60%
Mefenoxam**	1.70%
Fludioxonil***	1.12%
Azoxystrobin****	0.90%
<hr/>	
Other Ingredients:	73.68%
Total:	100.00%

*CAS No. 153719-23-4

**CAS No. 70630-17-0 and CAS No. 69516-34-3

***CAS No. 131341-86-1

****CAS No. 131860-33-8

One gallon of Four-way VAP contains 2.13 lb. thiamethoxam, 0.16 lb. mefenoxam, 0.11 lb. fludioxonil and 0.08 lb. azoxystrobin.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1384

EPA Est. XXXXX

SCP 1384A-M(draft-b legumes)

_____gallons

Net Contents

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call <i>[1 800 888 8372] [Alternate phone number]</i>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation Harmful if swallowed Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet Avoid contact with eyes skin or clothing

Environmental Hazards

This product is toxic to wildlife freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates Runoff may be hazardous to aquatic organisms in neighboring areas Exposed treated seed may be hazardous to wildlife Do not contaminate water when disposing of equipment wash water

Thiamethoxam is highly toxic to bees exposed to direct treatment and effects may be possible as a result of exposure to translocated residues in blooming crops

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water This chemical may leach into the ground water if used in areas where soils are permeable particularly where the water table is shallow

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination

Azoxystrobin in this product can be persistent for several months or longer

Azoxystrobin degradation products and thiamethoxam have properties and characteristics similar to chemicals which are known to leach through soil to groundwater under certain conditions as a result of agricultural use Use of these chemicals in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination

Physical and Chemical Hazards

Do not use pour spill or store near heat or open flame

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Crop injury ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather or crop conditions presence of other materials or other influencing factors in the use of the product which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller To the extent permitted by

applicable law Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions To the extent permitted by applicable law (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA and (2) Buyer and User assume the risk of any such use TO THE EXTENT PERMITTED BY APPLICABLE LAW SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL

To the extent permitted by applicable law in no event shall SYNGENTA be liable for any incidental consequential or special damages resulting from the use or handling of this product TO THE EXTENT PERMITTED BY APPLICABLE LAW **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT**

SYNGENTA and Seller offer this product and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 Refer to supplemental labeling under Agricultural Use Requirements in the Directions for Use section for information about this standard

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage disposal or cleaning of equipment

Pesticide Storage

Store in a cool dry place Storage for extended periods above 90 F is not recommended

For minor spills leaks etc follow all precautions indicated on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility If these wastes cannot be used according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods

Container Handling*(non refillable)*

Non refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Container Handling*(refillable)*

Refillable container Refill this container with pesticide only Do not use this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill by incineration or by other procedures allowed by state and local authorities

The Syngenta Logo is Trademarks of a Syngenta Group Company
©2012 Syngenta

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro North Carolina 27419 8300

SCP 1384A M(draft b legumes)

Four way VAP 1384A M(draft b legumes)clean lg 8 1 12 **000100 01384 20120801 b legumes pdf**