



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

March 4, 2024

Patsy D. Laird  
Regulatory Product Manager  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419

**Subject:** Notification per PRN 98-10 – Adding a new pest to product label  
**Product Name:** Fortenza® Red  
**EPA Registration Number:** 100-1418  
**Application Date:** March 28, 2023  
**Case Number:** 481791

Dear Patricia Laird:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Andrés Garzón at (202) 566-2690 or at [GarzonMoreno.Andres@epa.gov](mailto:GarzonMoreno.Andres@epa.gov).

Sincerely,

 For

Gene Benbow, Product Manager 7  
Invertebrate & Vertebrate Branch 3  
Registration Division (7505P)  
Office of Pesticide Programs

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

03/04/2024

[Master label]

**Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

**CYANTRANILIPROLE    GROUP 28    INSECTICIDE**

**Fortenza® Red**

Insecticide

A seed treatment product for protection against early-season damage caused by [Colorado potato beetle and European corn borer on potato;] cutworms and flea beetles on rapeseed crop subgroup 20A; cutworms and wireworms on sunflower crop subgroup 20B and cottonseed crop subgroup 20C; cutworms, grubs, wireworms, fall armyworm, and seedcorn maggot on corn; bean leaf beetle, **soybean gall midge**, thrips, grubs, and wireworms on soybeans; and rice water weevil on rice (dry-seeded)

Active Ingredient:	
Cyantraniliprole <sup>1</sup> .....	48.8%
<hr/>	
Other Ingredients:	51.2%
<hr/>	
Total:	100.0%

<sup>1</sup>CAS No. 736994-63-1

Fortenza Red is formulated as a flowable suspension that contains 5.0 lb/gal [600 g/L] of cyantraniliprole (FS). One fluid ounce of Fortenza Red contains 17.72 grams of cyantraniliprole.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

See additional precautionary statements and directions for use [in booklet] [on label].

EPA Reg. No. 100-1418  
EPA Est. xxxxx

---

Net Contents

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOTLINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-888-8372</b>	

---

**PRECAUTIONARY STATEMENTS**

---

**Hazards to Humans and Domestic Animals****CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.

## Personal Protective Equipment (PPE)

### Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, polyethylene, polyvinyl chloride (PVC) or Viton™  $\geq 14$  mils
- Shoes plus socks

### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

#### Users should:

- Wash thoroughly with soap and water after handling.
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide is toxic to aquatic invertebrates. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not contaminate water when disposing of equipment washwater.

---

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

---

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception:** If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, natural rubber  $\geq$  14 mils, polyethylene, polyvinyl chloride (PVC) or Viton  $\geq$  14 mils
- Shoes plus socks

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.**

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta,

no claims are made to guarantee the germination of carry-over seed [or propagating material] for all crop seed.

---

## **PRODUCT INFORMATION**

---

Cyantraniliprole is a broad-spectrum insecticide belonging to the chemical class of diamides. Cyantraniliprole products are effective on the larval stages of lepidopteran insects and some other insect pests, including some coleopterans and dipterans. The length of control of the major insect pests will vary depending on the product use rate, insect pressure, crop growth and maturity, and soil and environmental conditions. When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.

### **Resistance Management**

Fortenza Red contains cyantraniliprole, a Group 28 insecticide. Cyantraniliprole is a systemic insecticide belonging to the diamide class of chemistry. Diamides cause muscle contraction and paralysis in insects by activating muscle ryanodine receptors.

Insect populations may contain individuals naturally resistant to Group 28 insecticides, and, if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refugia (untreated areas). Sound management programs also consider cultural and biological control practices.

### **In order to maintain susceptibility to this class of chemistry:**

- Use products at their full, specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless



otherwise excluded by product labels. Do not exceed the maximum amount of this insecticide's mode of action allowed per growing season.

- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

Syngenta encourages responsible product stewardship to ensure effective long term control of the insect pests on this label.

**For additional information on Insect Resistance Management:**

- Contact Syngenta representatives at 1-866-796-4368.
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org>.

---

**MIXING PROCEDURES**

---

**Apply Fortenza Red only with Syngenta-approved seed-treating equipment.**

Potatoes may be treated either on farm by the grower with Syngenta-approved equipment or at a commercial seed-treatment facility. All other crops must **only** be treated at Syngenta-approved commercial seed-treatment facilities. Not for use in hopper box, planter box, slurry box, or other farmer applied applications.

**Important:** Thoroughly recirculate or agitate the container of Fortenza Red prior to use.

Apply Fortenza Red as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect or disease control. Thoroughly mix the specified amount of Fortenza Red into the required amount of water for the slurry treater and dilution rate to be used. (See **Fortenza Red in Tank Mixtures.**) Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment.

- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155 (c).
- Allow seed to dry before bagging.
- Store away from feed and foodstuffs.

Fortenza Red has been found to be compatible with some liquid inoculant products. Fortenza Red may be mixed or applied sequentially with approved liquid inoculants. Consult the maker of the liquid inoculants and a Syngenta Crop Protection representative for directions before applying Fortenza Red with inoculants.

### **Fortenza Red in Tank Mixtures:**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

### **SEED BAG LABEL REQUIREMENTS**

---

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with cyantraniliprole insecticide.
- Do not use treated seed for food, feed or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with Fortenza Red (cyantraniliprole):

- Store treated seed away from feed and foodstuffs.
- Do not allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Use of Fortenza Red as a seed treatment must be communicated to all personnel involved in seasonal insect control recommendations.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crop or weeds. Ensure that the planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops and weeds.
- Corn seeds must be planted at a minimum soil depth of 1½ inches.
- Potato seed tubers must be planted at a minimum soil depth of 2 inches.
- Rapeseed crop subgroup 20A seeds must be planted at a minimum soil depth of ¼ inch.
- Rice (dry-seeded) must be planted at a minimum soil depth of ½ inch.
- Soybean must be planted at a minimum soil depth of 1 inch.
- Sunflower crop subgroup 20B seeds must be planted at a minimum soil depth of 1 inch.

- Cottonseed crop subgroup 20C seeds must be planted at a minimum soil depth of ½inch.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Immediately cover or collect seeds spilled during loading and planting. Do not allow treated seed to remain uncovered on the soil surface.
- Dispose of all excess treated seed by burying seed away from bodies of water. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate bodies of water when disposing of excess treated seed or washwaters of planting equipment.
- Excess treated seed may be used for ethanol production only if (1) ethanol production by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used for agronomic practice.
- Dispose of seed packaging in accordance with local requirements.
- For all crops: Do not apply a total of more than 0.4 lb ai per acre of cyantraniliprole-containing products per calendar year; this is the total cyantraniliprole applied by seed treatment, soil application and foliar application.
- For corn: This seed has been treated with a maximum of 0.5 mg cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- For potato: This seed has been treated with a maximum of 0.0135 lb cyantraniliprole per 100 lb seed tuber. Do not plant more than 3000 tubers per acre when treated at 0.0135 lb ai/100 lb of seed.
- For rapeseed crop subgroup 20A [based on flax seed]: This seed has been treated with a maximum of 0.05 mg active cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- For rice (dry-seeded): This seed has been treated with a maximum of 0.03 mg cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- For soybean: This seed has been treated with a maximum of 0.076 mg cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- For sunflower crop subgroup 20B: This seed has been treated with a maximum of 0.2 mg active cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- For cottonseed crop subgroup 20C: This seed has been treated with a maximum of 0.9 mg active cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre. Do not apply more than 0.17 lb active cyantraniliprole per acre as a seed treatment application.
- Including the Fortenza Red seed treatment, make no more than two applications of cyantraniliprole or other Group 28 products per generation to the same insect species on a crop or within a 30-day period (count planting date as day 1 if using

treated seed). Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide).

### **Crop Rotation**

- There is no plant back restriction for conversion of a treated field or for making a new or replacement planting into established orchards or fields of Bushberries (Crop Subgroup 13-07B); Citrus (Crop Group 10-10); Pome Fruits (Crop Group 11-10); Stone Fruits (Crop Group 12); Low Growing Berries (Crop Subgroup 13-07G); or Tree Nuts (Crop Group 14-12).
- Crops on this label and the following crops or crop groups may be planted immediately following the last application of Fortenza Red: Brassica Leafy Vegetables (Crop Group 5); Bulb Vegetables (Crop Group 3-07); Corn (Field, Pop, Seed and Sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Fruiting Vegetables (Crop Group 8-10); Leafy Vegetables (except Brassica) (Crop Group 4); Leaves of Root and Tuber Vegetables (Crop Group 2); Legume Vegetables (Crop Groups 6 and 7); Low Growing Berries (Crop Subgroup 13-07H); Oilseeds (Crop Group 20); Peanuts; Root and Tuber Vegetables (Crop Subgroups 1B and 1C); Tobacco.
- The following crops or crop groups may be planted 30 days following the last application of Fortenza Red: Cereal Grains (Crop Group 15); Forage, Fodder and Straw of Cereal Grains (Crop Group 16); Grass Forage, Fodder and Hay (Crop Group 17); Nongrass Animal Feeds (forage, fodder, straw and hay) (Crop Group 18); Sugar beets.
- All other crops cannot be planted until 12 months after the last application of cyantraniliprole.

**CROP USE DIRECTIONS**

**CORN (FIELD, POP, SEED AND SWEET) – NOT FOR USE IN CALIFORNIA**

Crop	Early-season Protection Against Damage Caused by:	Use Rate	
		mg ai/seed	fl oz product/100 lb seed (based on 1800 seeds/lb)
<b>Corn</b> Field Pop Seed Sweet <b>Not for Use in California</b>	Cutworms Grubs Wireworms Fall armyworm Seedcorn maggot	0.125 – 0.5	1.24 – 5.0
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>Fortenza Red at labeled rates provides protection against damage caused by cutworms, grubs, wireworms, fall armyworm, and seedcorn maggot during early-season growth and development of corn.</li> <li>Full-season protection or complete reduction of damage is <b>not</b> expected against the listed insects.</li> <li>Based on numerous popcorn seed-safety studies, popcorn use rates &gt;0.125 mg ai/seed (2.75 fl oz/100 lb seed, based on 4,000 seeds/lb) on individual hybrids should be tested for seed safety to evaluate specific genetic tolerance to higher rates of Fortenza Red.</li> <li>It is recommended to apply Fortenza Red with compatible and registered seed treatment fungicides such as Apron XL<sup>®</sup>, Dynasty<sup>®</sup>, Maxim<sup>®</sup> 4FS, Maxim Quattro, and Vibrance<sup>®</sup>, which are proven to provide protection from seed and seedling diseases; Cruiser<sup>®</sup> 5FS for additional early season insect pest protection; and Avicta<sup>®</sup> Complete Corn for early-season disease, insect and nematode protection.</li> <li>Follow planter manufacturer instructions for use of talc or other hopper box additives at planting.</li> </ul>			
<b>USE RESTRICTION</b>			
<ul style="list-style-type: none"> <li><b>DO NOT</b> apply more than 0.4 lb ai of cyantraniliprole-containing products per acre per calendar year, including all types of applications (seed treatment, soil, foliar).</li> </ul>			

## POTATO SEED PIECE TREATMENT

Fortenza Red is to be used as an integral part of potato pest management strategy. This strategy includes the use of high quality certified potato seed, crop rotation, monitoring proper insect population thresholds, appropriate comprehensive insect control measures, optimal harvest time of potato tubers and proper handling of the potatoes. Consult your local agricultural extension agent for more detailed information on insect management practices.

Crop	Protection Against Damage Caused by:	Use Rate (See Tables 1 and 2 for additional rate information.)		
		[grams ai/100 kg seed tubers]	lb ai/100 lb seed tubers	fl oz product/100 lb seed tubers
Potato	Colorado potato beetle European corn borer	[6 – 13.5]	0.006 – 0.0135	0.15 – 0.35
<b>Additional Information</b>				
<ul style="list-style-type: none"> <li>• Potatoes may be treated either on farm by the grower with Syngenta-approved equipment or at a commercial seed-treatment facility.</li> <li>• Fortenza Red used at labeled rates provides protection against damage caused by Colorado potato beetle and European corn borer during early- to mid-season growth and development of potatoes.</li> <li>• Full-season protection or complete reduction of damage caused by labeled pests is <b>not</b> expected with the use of Fortenza Red.</li> <li>• If a crop failure occurs, treated potato seed pieces may be replanted if the total cyantraniliprole rate applied does not exceed 0.4 lb ai per acre.</li> <li>• For protection against certain seed- and soil-borne diseases of potatoes, Fortenza Red may be applied with Maxim 4FS, CruiserMaxx® Vibrance Potato, CruiserMaxx Potato, or CruiserMaxx Potato Extreme family of products.</li> <li>• Application of Fortenza Red with CruiserMaxx Potato or CruiserMaxx Potato Extreme also provides protection against potato aphid, green peach aphid, beet leaf hopper and psyllids.</li> <li>• If an inert dust (fir bark or talc etc.) or a dust-based fungicide is applied, apply Fortenza Red prior to applying the dust treatments.</li> </ul>				
<b>USE RESTRICTIONS</b>				
<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 0.4 lb ai of cyantraniliprole-containing products per acre per calendar year, including all types of applications (seed treatment, soil, foliar).</li> <li>• <b>DO NOT BAG POTATO SEED THAT IS TREATED WITH ANY LIQUID SEED TREATMENTS.</b></li> </ul>				

### Application of Fortenza Red

Apply Fortenza Red utilizing Syngenta-approved seed treating systems designed to apply liquid seed treatments of potatoes. Uneven or incomplete seed coverage may not give the desired level of insect control. For slurry treatment, thoroughly mix the specified amount of Fortenza Red into the required amount of water for the slurry treater and dilution rate to be used. Maintain constant agitation of the slurry during the seed treatment process. Follow the manufacturer’s application instructions for the seed treatment equipment being used with appropriate set-up and calibration. Calibrate the equipment so that every potato seed tuber is uniformly coated with a fine layer of the slurry mix without any excess dripping off of the treated seed.

### Storing Treated Potato Seed Pieces

If the treated potato seed piece needs to be stored or held for a few days (2 to 3 weeks maximum), make sure the potato seed piece is stored in well ventilated areas that allow air to move through and out the treated potato seed piece. An ideal air temperature is

60 degrees Fahrenheit at a relative humidity of 85 to 90 percent. Do not allow free moisture to form within or around the treated potato seed piece during storage. If possible, allow treated potato seed pieces to dry during transit and planted the same day of treatment.

**Syngenta, LLC makes no claims as to the effect of this product or delivery systems on germination of the potato seed. The user, buyer or applicator of the seed treatment assumes all risks from such application.**

**Note:** Treatment of highly damaged or bruised potato seed, or seed known to be of low vigor and poor quality, or potato seed that is deemed “physiologically old”, may result in reduced germination and/or reduction of seed and seedling vigor and multiple stems from germination of the seed. When in doubt or if the status/condition of the potato seed tubers is unknown, treat a small sample batch of the same potato seed load with Fortenza Red using specified rates, equipment, and application procedures; before treating the total seed lot. Conduct this test on a small batch of the potato seed and observe the germination, emergence, stem count from the germinating seed. Consult with local experts in the region or conduct the test with university or area experts. Only if the data confirm that the seed treated with Fortenza Red are acceptable should the rest of the seed load from which the sample was taken be treated. **Due to seed quality, seed condition and seed storage conditions beyond the control of Syngenta LLC, no claims are made to guarantee the germination and/or performance of the potato seed tuber from treatment with Fortenza Red.**

**Table 1. Empirical Calculations based on popular potato seeding rates:**

Seeding rate	Expected lb ai per acre at popular seeding rates and expected use rate of Fortenza Red			
	Pounds/acre	0.006 lb ai/100 lb	0.01 lb ai/100 lb	0.0135 lb ai/100 lb
1500		0.090	0.150	0.203
2000		0.120	0.200	0.270
2500		0.150	0.250	0.338
3000		0.180	0.300	0.405
3500		0.210	0.350	Do not apply. <sup>1</sup>

<sup>1</sup> The lb ai applied per acre at this seeding density and application rate exceeds the maximum allowable 0.4 lb ai per acre.

**Table 2. Empirical Calculations based on popular potato seeding rates:**

Seeding rate	Expected fl oz per acre at popular seeding rates and expected use rate of Fortenza Red			
	Pounds/acre	0.15 fl oz/100 lb	0.26 fl oz/100 lb	0.35 fl oz/100 lb
1500		2.3	3.8	5.2
2000		3.1	5.1	6.9
2500		3.8	6.4	8.6
3000		4.6	7.7	10.4
3500		5.4	9.0	Do not apply. <sup>1</sup>

<sup>1</sup>The fl oz applied per acre at this seeding density and application rate exceeds the maximum allowable 11 fl oz per acre.



**OILSEED CROP GROUP 20**

Crops	Early-season Protection Against Damage Caused by:	Use Rate	
		lb ai/100 lb seed	fl oz product/100 lb seed
<b>Rapeseed Crop Subgroup 20A:</b> Borage Crambe Cuphea Echium Flax seed Gold of Pleasure Hare's Ear Mustard Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Oil radish Poppy seed Rapeseed (including Canola) Sesame Sweet Rocket Cultivars, varieties, and/or hybrids of these	Cutworms	0.30	7.7
	[Flea beetles]	[0.80]	[20.4]
<b>Additional Information</b> <ul style="list-style-type: none"> <li>• Fortenza Red at labeled rates provides protection against the damage caused by cutworms during early-season growth and development of rapeseed crop subgroup 20A.</li> <li>• Full-season protection or complete reduction of damage from the use of Fortenza Red is <b>not</b> expected against the listed insects.</li> <li>• For protection against damage caused by flea beetles when applied at 7.7 fl oz product per 100 lb seed, Fortenza Red must be tank mixed with Helix® Vibrance or other thiamethoxam-containing seed treatment product.</li> <li>• In all cases, it is recommended to apply Fortenza Red with Helix Vibrance for control of seedling diseases and other insect pests of seedling crops.</li> <li>• Follow planter manufacturer instructions for use of talc or other hopper-box additives at planting.</li> </ul>			
<b>USE RESTRICTION</b>			
<b>DO NOT</b> apply more than 0.40 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).			

Crops	Early-season Protection Against Damage Caused by:	Use Rate (based on an average of 4,500 seeds/lb)		
		[mg ai/seed]	lb ai/100,000 seed	fl oz product/100,000 seed
<b>Sunflower Crop Subgroup 20B:</b> Calendula Castor Oil plant Chinese Tallowtree Euphorbia Evening Primrose Jojoba Niger seed Rose hip Safflower Stokes Aster Sunflower Tallowwood Tea Oil plant Vernonia Cultivars, varieties, and/or hybrids of these	Cutworms	[0.1 – 0.2]	0.022 – 0.044	0.56 – 1.1
<b>Additional Information</b> <ul style="list-style-type: none"> <li>Fortenza Red at labeled rates provides protection against the damage caused by cutworms during early-season growth and development of sunflower crop subgroup 20B.</li> <li>Full-season protection or complete reduction of damage from the use of Fortenza Red is <b>not</b> expected against the listed insects.</li> <li>[When applied at these rates in combination [with Cruiser 5FS, Fortenza Red may provide additional protection against wireworms.]</li> <li>It is recommended to apply Fortenza Red with compatible and registered seed treatment fungicides such as Apron XL, Dynasty, Bion®, Maxim 4FS or Plenaris® 200FS, which are proven to control seed and seedling diseases, and with Cruiser 5FS for additional early-season insect pest protection.</li> <li>Follow planter manufacturer instructions for hopper box additives at planting.</li> </ul>				
<b>USE RESTRICTION</b>				
<b>DO NOT</b> apply more than 0.40 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).				

Crops	Early-season Protection Against Damage Caused by:	Use Rate	
		mg ai/seed	fl oz product/100 lb seed (based on an average of 4,500 seeds/lb)
<b>Cottonseed Crop Subgroup 20C</b> Cottonseed Cultivars, varieties, and/or hybrids of these	Cutworms	0.1 – 0.9	2.55 – 23.0
<b>Additional Information</b> <ul style="list-style-type: none"> <li>• Fortenza Red at labeled rates provides protection against the damage caused by cutworms during early-season growth and development of cotton.</li> <li>• Full-season protection or complete reduction of damage from the use of Fortenza Red is <b>not</b> expected against the listed insects.</li> <li>• [When applied at these rates in combination with Cruiser 5FS, Fortenza Red may provide additional protection against wireworms.]</li> <li>• It is recommended to apply Fortenza Red with compatible and registered seed treatment fungicides such as Apron XL, Dynasty, Dynasty CST, Bion, Maxim 4FS, Vibrance, or Vibrance CST, which are proven to provide early-season protection from seed and seedling diseases, and with Cruiser 5FS for additional early-season insect pest protection.</li> <li>• Follow planter manufacturer instructions for hopper box additives at planting.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 0.17 lb active cyantraniliprole per acre as a seed treatment application.</li> <li>• <b>DO NOT</b> apply more than 0.40 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).</li> </ul>			

**RICE (DRY SEEDED ONLY) – NOT FOR USE IN CALIFORNIA**

Crop	Early-season Protection Against Damage Caused by:	Use Rate	
		mg ai/seed	fl oz product/100 lb seed (based on an average of 21,000 seeds/lb)
Rice	Rice water weevil	0.03	3.47

**Additional Information**

- Fortenza Red at labeled rates provides protection against the damage caused by rice water weevil during early-season growth and development of rice.
- Fortenza Red-treated rice seed may be planted by drill or broadcast (ground or aerial) on soil.
- Full-season protection or complete reduction of damage from the use of Fortenza Red is **not** expected against the listed insects.
- It is recommended to apply Fortenza Red with compatible and registered seed treatment fungicides such as Apron XL, Dynasty, or Maxim 4FS, which are proven to provide early-season protection from seed and seedling diseases, and Cruiser 5FS for additional early-season insect pest protection.

**USE RESTRICTIONS**

- When broadcasting Fortenza Red-treated seeds on soil, the rice seeds must be incorporated into the soil.
- Fortenza Red-treated dry rice seed cannot be soaked or pre-germinated before seeding.
- **DO NOT** apply more than 0.17 lb active cyantraniliprole per acre as a seed treatment application.
- **DO NOT** apply more than 0.4 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).
- **DO NOT** use Fortenza Red-treated rice fields for the aquaculture of edible fish or crustacea (including crawfish) during the rice growing season from planting through harvest.

**SOYBEAN (INCLUDING SOYBEAN VEGETABLE) – NOT FOR USE IN CALIFORNIA**

Crop	Early-season Protection Against Damage Caused by:	Use Rate	
		mg ai/seed	fl oz product/100 lb seed (based on an average of 3,000 seeds/lb)
<b>Soybean (including soybean vegetable)</b>  <b>Not for Use in California</b>	Bean leaf beetle <u>Soybean gall midge</u> Thrips Grubs Wireworms	0.038 – 0.076	0.627 – 1.25
<b>Additional Information</b> <ul style="list-style-type: none"> <li>Fortenza Red at labeled rates provides protection against the damage caused by bean leaf beetle, <u>soybean gall midge</u>, thrips, grubs and wireworms during early-season growth and development of soybeans.</li> <li>Full-season protection or complete reduction of damage from the use of Fortenza Red is <b>not</b> expected against the listed insects.</li> <li>It is recommended to apply Fortenza Red with compatible and registered seed treatment fungicides such as Apron XL, Dynasty, Maxim 4FS, Mertect® 340-F, and Vibrance, which are proven to provide protection from seed and seedling diseases; with Cruiser 5FS for additional early-season insect pest protection; or with a combination fungicide/insecticide product such as CruiserMaxx Vibrance for early-season disease and insect protection.</li> <li>Follow planter manufacturer instruction for use of talc or other hopper box additives at planting.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li><b>DO NOT</b> apply more than 0.4 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).</li> </ul>			

---

## ROTATIONAL CROP RESTRICTIONS

---

- There is no plant back restriction for conversion of a treated field or for making a new or replacement planting into established orchards or fields of Bushberries (Crop Subgroup 13-07B); Citrus (Crop Group 10-10); Pome Fruits (Crop Group 11-10); Stone Fruits (Crop Group 12); Low Growing Berries (Crop Subgroup 13-07G); or Tree Nuts (Crop Group 14-12).
- Crops on this label and the following crops or crop groups may be planted immediately following the last application of Fortenza Red: Brassica Leafy Vegetables (Crop Group 5); Bulb Vegetables (Crop Group 3-07); Corn (Field, Pop, Seed and Sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Fruiting Vegetables (Crop Group 8-10); Leafy Vegetables (except Brassica) (Crop Group 4); Legume Vegetables (Crop Group 6); Legume Vegetables (Crop Groups 6 and 7); Low Growing Berries (Crop Subgroup 13-07H); Oilseeds (Crop Group 20); Peanuts; Root and Tuber Vegetables (Crop Subgroups 1B and 1C); Tobacco.
- The following crops or crop groups may be planted 30 days following the last application of Fortenza Red: Cereal Grains (Crop Group 15); Forage, Fodder and Straw of Cereal Grains (Crop Group 16); Grass Forage, Fodder and Hay (Crop Group 17); Nongrass Animal Feeds (forage, fodder, straw and hay) (Crop Group 18); Sugar beets.
- All other crops cannot be planted until 12 months after the last application of cyantraniliprole.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

### **Pesticide Storage**

Store in a cool, dry place. Do not store above 90°F for extended periods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

### **Pesticide Disposal**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

### **Container Handling (less than or equal to 5 gallons)**

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**

**Refillable container.** Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining

contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

---

Fortenza<sup>®</sup>, Apron XL<sup>®</sup>, Avicta<sup>®</sup>, Bion<sup>®</sup>, Cruiser<sup>®</sup>, CruiserMaxx<sup>®</sup>, Dynasty<sup>®</sup>, Helix<sup>®</sup>, Maxim<sup>®</sup>, Mertect<sup>®</sup>, Plenaris<sup>®</sup>, Vibrance<sup>®</sup>, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC  
© 20XX Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

Fortenza Red 1418 MAS 1118 NOTIF-B 0423-HI – jeb – 02/20/2024  
000100-01418.20230412B.FORTENZA\_RED.NOTIF.APR2023-HI.PDF