

100-1433

2/1/2012

JACKIE 10 of 23



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1433

Date of Issuance:

FEB 01 2012

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Breezeway™

Name and Address of Registrant (include ZIP Code):

Fred. J. Pearson
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Page 1 of 2

Signature of Approving Official:

Shaja B. Joyner
Shaja B. Joyner, Product Manager (20)
Fungicide Branch, Registration Division (7505P)

Date:

FEB 01 2012

20/23

2. Revise the "EPA Registration Number to read, "EPA Reg. No. 100-1433.
3. Update the fourth bullet point in the "If Swallowed" section of the FIRST AID statements on pages 2 and 18 to read, "Do not give anything to an unconscious person."
4. Update the Container Handling sections on pages 15 and 20 to remove the third sentence that reads, "Offer for recycling if available" as this statement already appears at the end of the paragraph.
5. On pages 2 and 18 under the Precautionary Statements, you may remove the last two sentences beginning with "Wash thoroughly with soap....." and ending with ".....contaminated clothing before reuse." Such language is found in the User Safety Recommendations section.
6. On page 3, place the User Safety Requirements text in a box.
7. On page 6 remove language from the first sentence of the last "DO NOT" statement so that it reads, "DO NOT spray when conditions favor drift."
8. On page 7, carry over the Resistance Management heading to the following page.
9. On page 8 remove the language "beyond the target area" from the first sentence under Spray Drift Management so that the sentence reads "To avoid spray draft, do not apply when conditions favor drift."
10. On page 10, under the Application Instructions, add the turf restrictions language:
 - Do not apply by air
 - Do not apply through irrigation systems (chemigation)
11. On page 14 add in the seasonal maximum application for azoxystrobin as you have done for difenoconazole.
12. On page 15 and page 19, add a box around the Storage and Disposal text.
13. Update box 5 of your CSFs with the correct PM and team number the next time you submit them (Shaja Joyner/20).
14. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,



Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7505P)

Enclosure: Label stamped "Accepted with Comments"

Breezeway™

Fungicide

A broad-spectrum fungicide for prevention and control of certain diseases in golf course turfgrasses only.

Active Ingredients:	
Azoxystrobin*	18.2%
Difenoconazole**	11.4%
<hr/>	
Other Ingredients:	70.4%
<hr/>	
Total:	100.0%

*CAS No. 131860-33-8
**CAS No. 119446-68-3

Contains 1.67 lb. of azoxystrobin active ingredient and 1.05 lb. of difenoconazole active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-XXXX

EPA Est.

Product of

_____ gallons
Net Contents

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

**ACCEPTED
with COMMENTS
In EPA Letter Dated
FEB 01 2012**
**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
100-1433**

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p>HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F or G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Difenoconazole is toxic to fish, mammals and aquatic invertebrates. Drift and runoff may be hazardous to **estuarine/marine** organisms in water adjacent to treated area.

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. This chemical may leach into ground water if used in areas where permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product has a potential for runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach surface water.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

90423

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

PRODUCT INFORMATION

Breezeway fungicide is a broad-spectrum product containing two fungicides. It has preventative, systemic and curative properties. Breezeway fungicide provides excellent disease control of many major turfgrass diseases. Breezeway fungicide may be applied as a foliar spray and integrated into a resistance management program. All applications should be made according to the use directions that follow.

USE PRECAUTIONS AND RESTRICTIONS

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN TURF INJURY AND/OR POOR DISEASE CONTROL.

ATTENTION

DO NOT apply by air.

DO NOT apply through irrigation systems (chemigation)

DO NOT spray Breezeway fungicide where spray drift may reach apple trees. Breezeway fungicide is extremely phytotoxic to certain apple varieties.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

USE INFORMATION

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application.

For control of foliar diseases, allow Breezeway fungicide applications to completely dry prior to irrigating. For control of soil-borne diseases, Breezeway fungicide can be watered in after application.

Adjuvants: When an adjuvant is to be used with this product, Syngenta recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of Breezeway fungicide has been used. If resistant isolates to Group 3 or Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure with highly susceptible varieties, or when environmental conditions are conducive to disease.

Breezeway fungicide is recommended for control of certain pathogens causing foliar, stem and root diseases including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass plants.

Integrated Pest Management (IPM): Breezeway fungicide should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required.

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce susceptibility to disease. Immunoassay detection kits and Extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Resistance Management

GROUP	11	3	FUNGICIDES
-------	----	---	------------

Breezeway fungicide contains two fungicides - azoxystrobin, a strobilurin fungicide in Group 11 and difenoconazole, a triazole fungicide in Group 3. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating with products having different modes of action or limiting the total number of applications per season. Syngenta encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label. Breezeway fungicide should not be alternated or tank mixed with any fungicide to which resistance has already developed.

Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. Breezeway fungicide should be alternated with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Since Breezeway fungicide is a QoI fungicide (Group 11 fungicide) and a DMI fungicide (Group 3 fungicide), avoid alternations with other QoIs such as pyraclostrobin, fluoxastrobin and trifloxystrobin or other DMI's such as propiconazole, tebuconazole and metaconazole. Addition of, or alternation to, a fungicide with Pythium activity such as Subdue® MAXX fungicide may be required if Pythium diseases are a major concern.

Turfgrass Tolerance: Breezeway fungicide plant tolerance has been found to be acceptable for all turfgrass species on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the turf to ensure that a phytotoxic response will not occur as a result of application.

Spray Drift Management: To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

MIXING AND APPLICATION METHODS

Spray Equipment

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.

- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be *16-mesh or coarser*.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - (1) Maintain 35-40 psi at nozzles.
 - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

Mixing Instructions

Breezeway fungicide is a suspension concentrate (SC) formulation.

Thoroughly clean spray equipment before using this product.

Make up only the amount of spray required for the immediate use.

To prepare a spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Breezeway fungicide to the tank, allowing time for good dispersion, then add an adjuvant, if recommended.

Do not use silicone based products with Breezeway fungicide due to possible phytotoxicity.

If tank mixes are required, products should be added to the spray tank in the following order: Breezeway fungicide, WG or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the spray tank to the desired volume to obtain the proper spray concentration.

Maintain agitation throughout the spraying operation. Do not allow spray mixture to stand overnight or for prolonged periods. If spray tank mixture is unsprayed for more than 18 hours (overnight), resuspend product with agitation for 20 minutes.

Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Breezeway Fungicide Alone (No Tank Mix)

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add Breezeway fungicide to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Breezeway fungicide has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Breezeway Fungicide + Tank Mixtures: Breezeway fungicide is usually compatible with many tank-mix partners registered for use on turf.

If tank mixes are desired, observe all directions, precautions and limitations on labeling of all products used. Consult compatibility charts or your local or State turf authority for compatibility information. Do not combine Breezeway fungicide in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use.

To determine the physical compatibility of Breezeway fungicide with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Breezeway fungicide to the spray tank.
- Allow Breezeway fungicide to completely disperse.
- Spray the mixture with the agitator running.

Application Instructions

Breezeway fungicide may be applied with all types of spray equipment commonly used for making ground applications. Proper adjustments and calibration of spraying equipment to give good coverage is essential for good disease control.

Ground Application

- Apply in sufficient water to provide good coverage. Typical application volumes range from 30 to 450 gallons of spray per acre.
- Thorough coverage is necessary to provide good disease control.

SPECIFIC DIRECTIONS FOR USE

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Use Rate (fl. oz. product per acre)	Application Interval (days)	Remarks*
Anthracnose (<i>Colletotrichum cereale</i>) (formerly known as <i>C. graminicola</i>)	0.3 to 0.725	13.1 to 31.6	14	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. For optimum disease control, alternate Breezeway fungicide with contact fungicides such as Daconil® brand products.
Brown Patch and Large Patch/Zoysia Patch (<i>Rhizoctonia solani</i>)	0.3 to 0.725	13.1 to 31.6	14-28	For cool season turfgrasses, apply when conditions are favorable for disease development. For large patch of all warm season turfgrasses, make 1 or 2 applications in fall prior to infection or when conditions are favorable for infection. A spring application prior to green-up may be required based on length of the infection period. Initiate first application for large patch/zoysia patch when soil temperature at a 2-4" depth averages 65° F.

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Use Rate (fl. oz. product per acre)	Application Interval (days)	Remarks*
Leaf and Sheath Spot (<i>Waitea circinata</i> var. <i>zeae</i> / <i>Chrysorhiza zeae</i> , formerly <i>R. zeae</i>)	0.5 to 0.725	21.8 to 31.6	14-28	For leaf and sheath spot, apply when conditions are favorable for infection such as sequential days, periods of temperatures at or above 90°F. Curative control may necessitate several applications. Use higher rates at shorter intervals for curative control. Target spray applications at crown of turfgrass.
Brown Ring Patch <i>Waitea circinata</i> var. <i>circinata</i>	0.5 to 0.725	21.8 to 31.6	14-28	Apply when conditions are favorable for disease development.
Cool Weather Brown Patch, Yellow Patch (<i>Ceratorhiza cerealis</i> , formerly <i>Rhizoctonia cerealis</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Make 1 or 2 applications in fall or when conditions are favorable for disease development.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.3 to 0.725	13.1 to 31.6	14-21	Apply preventatively when conditions are favorable for disease development. For optimum disease control, alternate Breezeway fungicide with contact fungicides such as a Daconil brand product. If dollar spot is active, use higher rates and combine with a Daconil brand product.

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Use Rate (fl. oz. product per acre)	Application Interval (days)	Remarks*
Fairy Ring (<i>Lycoperdon</i> spp., <i>Agrocybe</i> <i>pediades</i> , and <i>Bovistra plumbea</i>)	0.5 to 0.725	21.8 to 31.6	14-28	<p>For preventive control of fairy ring, apply early in the spring prior to the development of symptoms. Apply in 2-4 gallons of water per 1000 sq. ft. Irrigate into the thatch prior to the spray drying. Repeat the application within 14 to 28 days after the first application.</p> <p>For curative control, apply as soon as possible after fairy ring symptoms develop. Apply in 2-4 gallons of water per 1000 sq. ft. and irrigate lightly after application. Add the recommended rate of a wetting agent to the final spray. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. If area is hydrophobic, use wetting agents and irrigate prior to application(s) of Breezeway fungicide. Reapplication after 28 days may be required in some cases.</p>
Microdochium Patch (Formerly known as Fusarium Patch) (<i>Microdochium</i> <i>nivale</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
**Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.5 to 0.725	21.8 to 31.6	14-21	Use Breezeway fungicide in a preventive disease control program. Begin applications before disease is present and alternate with other fungicide chemistries that control gray leaf spot.
Leaf Rust Stem Rust Stripe Rust (<i>Puccinia</i> spp.)	0.5 to 0.725	21.8 to 31.6	14-28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Use Rate (fl. oz. product per acre)	Application Interval (days)	Remarks*
Leaf Spot (<i>Bipolaris sorokiniana</i> , <i>Bipolaris cynodontis</i> , <i>bipolaris</i> spp.)	0.5 to 0.725	21.8 to 31.6	14-21	Apply when conditions are favorable for disease development.
Melting Out (<i>Drechslera poae</i>)	0.5 to 0.725	21.8 to 31.6	14-21	Apply when conditions are favorable for disease development
Necrotic Ring Spot (<i>Ophiosphaerella korrae</i>)	0.725	31.6	14-28	Apply when conditions are favorable for disease development.
Pink Patch (<i>Limonomyces roseipellis</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Apply when conditions are favorable for disease development.
Powdery Mildew (<i>Erysiphe graminis</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Red Thread (<i>Laetisaria fuciformis</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Apply when conditions are favorable for disease development.
Southern Blight (<i>Sclerotium rolfsii</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Apply when conditions are favorable for disease development.
Summer Patch (<i>Magnaporthe poae</i>)	0.5 to 0.725	21.8 to 31.6	14-28	Initiate applications when soil temperatures reach 65°F at a 2 inch soil depth.
Take-All Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.5 to 0.725	21.8 to 31.6	28	Begin applications prior to disease symptom development. Make two applications 28 days apart in the spring and two applications 28 days apart in the fall.

*Do not exceed 0.50 gallons Breezeway fungicide per acre per year (1.45 fl. oz. Breezeway fungicide per 1000 square feet per year).

**Do not apply more than two sequential applications of Breezeway fungicide for control of gray leaf spot. For all other diseases when gray leaf spot is not present, do not apply more than three sequential applications of Breezeway fungicide.

One gallon (128 fl. oz.) Breezeway fungicide contains 1.67 lb. azoxystrobin and 1.05 lb. difenoconazole. Do not apply more than 0.52 lb. difenoconazole per acre per year.

Breezeway Fungicide: Rate Conversion Chart

Fl. Oz. Product Per 1000 Sq. Ft.	Fl. Oz. Product Per Acre	Maximum Yearly Applications
0.30	13.1	5
0.50	21.8	3
0.725	31.6	2

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original container only. Store in a cool, dry and well-ventilated place. Protect from excessive heat. Keep container closed when not in use. Do not store near food or feed.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [5 gallons or less]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

For Bulk and Minibulk Containers:

Container Handling

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Breezeway™, Daconil®, Subdue® MAXX, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

©201X Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

[NON-DETACHABLE CONTAINER LABEL]

GROUP 11 | 3 FUNGICIDES

Breezeway™

Fungicide

A broad-spectrum fungicide for prevention and control of certain diseases in golf course turfgrasses only.

Active Ingredients:

Azoxystrobin*	18.2%
Difenoconazole**	11.4%
<hr/>	
Other Ingredients:	70.4%
Total:	100.0%

*CAS No. 131860-33-8

**CAS No. 119446-68-3

Contains 1.67 lb. of azoxystrobin active ingredient and 1.05 lb. of difenoconazole active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-XXXX

EPA Est.

_____ gallons
Net Contents

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p>HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards

Difenoconazole is toxic to fish, mammals and aquatic invertebrates. Drift and runoff may be hazardous to **estuarine/marine** organisms in water adjacent to treated area.

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Azoxystrobin has

degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. This chemical may leach into ground water if used in areas where permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product has a potential for runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach surface water.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original container only. Store in a cool, dry and well-ventilated place. Protect from excessive heat. Keep container closed when not in use. Do not store near food or feed.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling [5 gallons or less]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Breezeway™ and the Syngenta logo are trademarks of a Syngenta Group Company.

©201X Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

Breezeway 100-XXXX 10-06-11 – df
000100-0XXXX.20111006.BREEZEWAY.PDF