

100-1434

9/26/2012

jacket 1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Thomas J Parshley
Syngenta Crop Protection, LLC
P O Box 18300
Greensboro, NC 27419

SEP 26 2012

Subject	Application for Pesticide Notification (PRN 2007-4)
Submission date	8/14/2012
Product Name	Medallion WDG
EPA Reg No	100-1434
EPA Decision Number	469587

Dear Mr Parshley

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10.

The Agency acknowledges the revisions to the Storage and Disposal section.

The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have questions concerning this letter, please contact Erin Malone at 703-347-0253.

Sincerely,

A handwritten signature in cursive script, appearing to read "C. Giles-Parker".

Cynthia Giles-Parker
Acting Product Manager 20
Fungicide Branch
Registration Division (7505P)

Medallion® WDG

[BOOKLET]

GROUP 12 FUNGICIDE

Fungicide

For control of certain foliar, stem, and root diseases in ornamentals grown in interiorscapes field nursery plantings, container nurseries forest nurseries, residential and commercial landscapes, greenhouses lath and shade houses or other enclosed structures

Active Ingredient	
Fludioxonil*	50 0%
Other Ingredients	50 0%
Total	100 0%

*CAS No 131341-86-1

NOTIFICATION

Medallion WDG is a 50% water dispersable granule

SEP 26 2012**KEEP OUT OF REACH OF CHILDREN****CAUTION**

See additional precautionary statements and directions for use inside booklet
EPA Reg No 100-1434

EPA Est 67545-AZ-1

Product of Switzerland

Formulated in the USA

SCP 1434A-L1 0412

[Product ID 51008 – Front Booklet/base]

8 ounces

Net Weight

EPA Est 67545-AZ-1

Product of Switzerland

Formulated in the USA

SCP 1434A-L2 0412

[Product ID 51008 – Back Booklet/base]

8 ounces

Net Weight

EPA Est 70992-CHE-003

Product of Switzerland

SCP 1434A-L3 0412

[Product ID 51334 – 99 lb (45 kg) Bulk Booklet]

99 pounds (45 kg)

Net Weight

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment	
NOTE TO PHYSICIAN	
If ingested induce emesis or lavage stomach Treat symptomatically	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals Causes moderate eye irritation Avoid contact with skin eyes or clothing Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet Remove and wash contaminated clothing before reuse

Personal Protective Equipment (PPE)

Applicators and other handlers must wear

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as Natural Rubber Selection Category A)

- Shoes plus socks
- Protective eyewear such as goggles or face shield

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users must

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsates.

Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

5/19

Surface Water Advisory

This chemical may contaminate water through drift of spray in wind. This chemical has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this chemical will be reduced by avoiding applications when conditions favor runoff (such as when soils are saturated and/or significant rainfall is forecast in the next 48 hours). Sound erosion control practices will reduce this chemical's contribution to surface water contamination.

Physical or Chemical Hazards

Do not use or store near heat or open flame. Do not use with or store near any oxidizing agents.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT**

SYNGENTA and Seller offer this product, and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons, either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides It contains requirements for training decontamination, notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours

Exception If the product is applied by drenching, the Worker Protection Standard, under certain circumstances allows workers to enter the treated area if there will be no

7/19

contact with anything that has been treated

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170) The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses

Do not enter treated areas without protective clothing until sprays have dried

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR PLANT INJURY

USE RESTRICTIONS AND PRECAUTIONS

Hawaii Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures

Nassau and Suffolk Counties, New York Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures

Do not apply Medallion WDG with any type of aircraft

RESISTANCE MANAGEMENT

Medallion WDG is a protective fungicide used to aid in the control of soil and foliar diseases Medallion WDG contains fludioxonil which is in the phenylpyrrole class of chemistry and has a unique mode of action which prevents fungal respiration (Fungicide Action Group 12) Fungal isolates with acquired resistance to Group 12 may eventually dominate the fungal population if Group 12 fungicides are used repeatedly or in successive years as the primary method of control for targeted species This may result in partial or total loss of control of those species by fludioxonil or other Group 12

8/19

fungicides A disease management program that includes alternation or tank mixes between Medallion WDG and other labeled fungicides that have a different mode of action may prevent pathogen populations from developing resistance Sanitation and other cultural practices to minimize disease are also recommended to aid in control as well as to assist in preventing/delaying resistance development

MIXING PROCEDURES

Prepare no more spray mixture than is needed for the immediate operation Thoroughly clean spray equipment before using this product Vigorous agitation is necessary to maintain uniformity of the spray mixture Maintain maximum agitation throughout the spraying operation Do not let the spray mixture stand overnight in the spray tank Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area

To determine the physical compatibility of Medallion WDG with other products, use a jar test as described below

Jar Compatibility Test Using a quart jar, add the proportionate amounts of the products to 1 qt of water Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last After thoroughly mixing, let stand for at least 5 minutes If the combination remains mixed or can be remixed readily it is physically compatible Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank

If using Medallion WDG in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label Do not exceed the dosage rates on any label Follow the most restrictive label precaution Do not mix this product with any other product whose label prohibits such mixing **Tank mixtures are permitted only in those states where the tank-mix partner is registered**

Note Use with oils or adjuvants may cause plant damage

Plant tolerance has been found acceptable for ornamentals listed in the ORNAMENTALS section of this label However, not all possible tank-mix combinations have been tested under all conditions When possible, it is recommended to test the combinations on a small portion of target plants to ensure a phytotoxic response will not occur as a result of application

Medallion WDG Alone Add $\frac{1}{3}$ of the required amount of water to the spray or mixing tank With the agitator running, add the labeled amount of Medallion WDG into the spray tank all at once Continue agitation while adding the remainder of the water Begin application of the spray solution after the material has completely dispersed into the mix water Maintain agitation until all of the mixture has been applied

Medallion WDG + Tank Mixtures Medallion WDG is usually compatible with Banner MAXX®, Subdue MAXX®, Daconil Ultrex®, Daconil WeatherStik® and other commonly used fungicides, insecticides and foliar nutrient products. However, the physical and biological compatibility of Medallion WDG with tank-mix partners should be tested before use. If in doubt, run a “jar compatibility test” or consult with Syngenta company representatives, other users, university or extension personnel before proceeding.

Add 1/3 of the required amount of water to the spray or mixing tank. With the agitator running, add the labeled amount of Medallion WDG into the tank all at once. Continue agitation while adding the remainder of the water. Allow Medallion WDG to dissolve and the product to completely disperse into the mix water. Then add the desired amount of other products recommended for tank mixture and allow them to become completely dispersed. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

Other wettable powders or water dispersible granules should be added to the water in the tank next, followed by flowable products, and emulsifiable concentrates added last. Provide sufficient mechanical or bypass agitation during mixing and application.

APPLICATION INSTRUCTIONS

Apply Medallion WDG at rates and timings as described in this label.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS

- Do not apply within 75 ft of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries.
- For all plantings within 150 ft of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
- Shut off the sprayer when at row ends.
- Spray the last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas.
- Do not cultivate within 10 ft of aquatic areas as to allow a vegetative filter strip.
- Do not apply when weather conditions favor drift to aquatic areas. Do not apply when gusts or sustained winds exceed 10 mph.
- Do not apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.

Directions for Use Through Microjet or Drip Chemigation Systems

For applications of Medallion WDG as soil drench applications only.

Do not use in overhead irrigation systems

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place

Medallion WDG alone or in tank mixture with other pesticides which are registered for microjet or drip application may be applied in irrigation water at rates specified in this label

Applications to the soil may be made through microjets or drip irrigation. With microjet systems, apply additional water after application is complete to remove residues from the foliage. Apply enough supplemental water to wet the root zones of the plants.

Do not apply this product through any other type of irrigation system.

Plant injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts.

Thorough coverage is necessary to provide good disease control. Where distribution patterns do not overlap sufficiently, unacceptable control may result. Where distribution patterns overlap excessively, injury to desirable plants may result.

Spray Preparation Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Fill the mix (supply) tank with the required amount of water. Start agitation in the tank. Agitate the solution until the Medallion WDG has completely dispersed into the solution. Then add desired amount of tank-mix partners. Maintain agitation in the tank and inject this mixture into the irrigation system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the fungicide in suspension.

Meter into irrigation water during the beginning of the irrigation cycle.

Operation Instructions

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed,

11/19

solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise

Specific Instructions for Public Water Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year
- 2 Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe
- 3 The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump
- 4 The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

12/19

- 5 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6 Systems must use a metering pump, such as a positive displacement injection pump (e g , diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7 Do not apply when wind speed favors drift beyond the area intended for treatment

ORNAMENTALS

Medallion WDG is a protectant fungicide for control of certain foliar, stem, and root diseases in ornamentals grown in interiorscapes, field nursery plantings, container nurseries, forest nurseries residential and commercial landscapes greenhouses, lath and shade houses or other enclosed structures

Hawaii Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures

Nassau and Suffolk Counties, New York Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures

Medallion WDG controls foliar diseases of ornamentals caused by *Rhizoctonia* spp , *Botrytis* spp , *Cercospora* spp , *Alternaria* spp , *Septoria* spp , and *Myrothecium* spp when applied on a regular schedule as a full coverage spray

Medallion WDG also will provide control of stem and root diseases caused by *Rhizoctonia* spp , *Fusarium* spp (e g , *F oxysporum*), *Cylindrocladium* spp , *Sclerotium* spp , and *Thielaviopsis* spp , when mixed with the potting media or as a drench to the root zone of plants

For control of stem and root diseases caused by *Pythium* spp and *Phytophthora* spp , tank mix Medallion WDG with labeled rates of Subdue MAXX See mixing instructions for these tank mixes

Maximum Use Rates

For indoor drench applications, use up to 30 oz of Medallion WDG per 1000 sq ft (80 lb product/A) per year or crop cycle Applications to pre-potting mix can be made up to 4 oz of Medallion WDG per cu yd

13/19

The high use rates specified for container nurseries, greenhouses, or other enclosed structures are due to the high organic matter soil mixes used in these systems and the high binding affinity of Medallion WDG for organic matter

For field grown and landscape ornamentals, apply up to a maximum of 1.5 oz /1000 sq ft /year (4 lb product/A/year) For outdoor container grown ornamentals, apply up to a maximum of 3 oz /1000 sq ft /year (8 lb product/A/year)

Plant Species

Medallion WDG has been tested and found to be safe on the ornamentals listed in this table at specified rates For plants not listed in the table, see the **NOTICE TO USER** box at the bottom of the table Numbers in parentheses refer to diseases controlled See Table 1

African Violets (1-12)	Coreopsis* (1-12)	Petunia (1-12)
Ageratum (1-10)	Cyclamen (1-12)	Pittosporium (8-12)
Alyssum (1-12)	Daffodil (4 9 10)	Poinsettia (1,4, 9-12)
Aster (1-12)	Dahlia* (1-12)	Portulaca* (1-12)
Azalea (4, 11, 12)	Daisy* (1-12)	Pothos (1-12)
Begonia (1-12)	Fern** (1-12)	Rose (1 4 9)
Bleeding Heart* (1-12)	Fuchsia* (1-4)	Salvia (1-12)
Bridal Veil* (1-12)	Gerbera Daisy (1-12)	Snapdragon (1-12)
Caladium (1-12)	Gomphrena (1-12)	Spathiphyllum (1-12)
Calendula (1-12)	Iris (4 9 10)	Sunflower* (1-12)
Carnation (1-12)	Lantana* (1-12)	Tobacco, Flowering* (1-12)
Celosia (1-12)	Lily (4 9 10)	Tulip (4 9 10)
Centrosa* (1-12)	Lysianthus (8-12)	Verbena (1-12)
Chenille* (1-12)	Marigold (1-12)	Vinca (1-12)
Christmas Cactus (1-12)	Mexican Heather* (1-7)	Wandering Jew* (1-12)
Chrysanthemums (1-12)	Nephtytis* (1-12)	Zinnia (1-12)
Coleus (1-12)	Pansy (1-12)	

*Indicates that only foliar applications have been tested for plant safety

**Do not apply Medallion WDG to leather leaf fern

Note

- Drench or at seedling applications to Impatiens or New Guinea Impatiens may cause stunting and/or chlorosis
- Foliar or drench applications to Geranium can cause stunting or chlorosis Responses may vary depending on environmental conditions Medallion WDG should be tested on a limited area to evaluate for any possible damage before proceeding with treatment of the entire crop

NOTICE TO USER Plant tolerance to Medallion WDG has been found to be acceptable for the specific genera and species listed on this label. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to Medallion WDG. Neither the Manufacturer nor the Seller has determined whether or not Medallion WDG can be used safely on ornamental plants not specified on this label. The professional user should determine if Medallion WDG can be used safely prior to commercial use. In a small area, test the required rates on a small number of plants for phytotoxicity prior to widespread use.

Table 1 Diseases Controlled by Medallion WDG

Foliar Diseases

- 1 Aerial Blight (*Rhizoctonia* spp)
- 2 Alternaria Leaf Blight (*Alternaria* spp)
- 3 Alternaria Leaf Spot (*Alternaria* spp)
- 4 Botrytis Blight (*Botrytis* spp)
- 5 Cercospora Leaf Spot (*Cercospora* spp)
- 6 Myrothecium Leaf Spot and Blight (*Myrothecium* spp)
- 7 Septoria Leaf Spot (*Septoria* spp)

Stem, Crown, and Root Rots

- 8 Cylindrocladium Stem and Root Rot (*Cylindrocladium* spp)
- 9 Fusarium Stem and Root Rot (*Fusarium* spp)
- 10 Rhizoctonia Stem and Root Rot (*Rhizoctonia* spp)
- 11 Southern Blight (*Sclerotium rolfsii*)
- 12 Black Root Rot (*Thielaviopsis* spp)

Foliar Spray

For control of *Rhizoctonia* spp , *Alternaria* spp , *Septoria* spp , *Myrothecium* spp , and *Cercospora* spp , use 1-2 oz /100 gal of water and spray to runoff at 7- to 14-day intervals while conditions are favorable for disease development.

For control of *Botrytis* spp use 2-4 oz /100 gal of water and spray to runoff at 7- to 14-day intervals while conditions are favorable for disease development. For management of the potential development of resistance in the *Botrytis* population, use no more than two consecutive applications of Medallion WDG before rotating to another effective product registered for *Botrytis* control on ornamentals with a different mode of action.

Notes (1) Under severe conditions, use the highest rate and/or the shortest application interval. (2) Use sufficient spray volume to wet the plants to the point of drip. (3) For a single foliar application, apply up to a maximum of 4 oz /100 gal (1 oz /25 gal).

Pre-Potting Growing Media Mix

Medallion WDG can be mixed with the potting media before seeding or transplanting by uniformly mixing 1-2 oz per cu yd of potting media. The desired amount of Medallion WDG should be mixed with 0.50-1 gal of water and applied to 1 (one) cu yd of potting media. It is recommended that Medallion WDG treatment should be made just before the plants are seeded or potted up. Uniform mixing can be accomplished by placing the potting mix in a rotating drum and spraying the Medallion WDG solution onto the mix while the drum is rotating.

Growing Medium Drench

At Seeding

For the control of damping-off, root, and stem diseases, mix 1 oz per 100 gallons of water. When using Medallion WDG for control of *Rhizoctonia* spp, apply sufficient mix to wet the upper one-half of the growing medium. For control of other root and stem diseases, completely drench the growing medium. Make only one application to the seeding crop prior to transplanting or transfer to larger containers.

Transplants and Cuttings

For the control of root and stem diseases, mix 1-2 oz per 100 gallons of water. When using Medallion WDG for control of *Rhizoctonia* spp, apply sufficient mix to wet the upper one-half of the growing medium. For control of other root and stem diseases, completely drench the growing medium. If needed, retreat transplants and cuttings with Medallion WDG as described above at 21- to 28-day intervals. Two applications per year during conditions favorable for disease development are usually adequate to control diseases of ornamentals.

Notes (1) Under severe conditions, use the highest rates and/or the shortest application interval. (2) For control of *Pythium* and *Phytophthora* diseases in addition to *Rhizoctonia* spp, *Cylindrocladium* spp, *Thielaviopsis* spp, *Fusarium* spp, and *Sclerotium* spp, tank mix Medallion WDG with labeled rates of Subdue MAXX. (3) Drench applications can be made at up to a maximum of 2 pt/sq ft to wet the root zone of plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry, secure place. Do not store this product under wet conditions.

Pesticide Disposal

Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling [For boxes and paper and plastic bags]

Non-refillable container Do not reuse or refill this container. Completely empty box or bag into application equipment. Then offer for recycling or dispose of empty box or bag in a sanitary landfill or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [fiber drums with liners]

Non-refillable container Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then offer for recycling if available or dispose of liner in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of it in a manner required by its liner.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER

Medallion® Banner MAXX®, Daconil Ultrex® Daconil WeatherStik®, Subdue MAXX®, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

©2012 Syngenta

<p>For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481</p>

Manufactured for
Syngenta Crop Protection, LLC
P O Box 18300
Greensboro, North Carolina 27419-8300

SCP 1434A-L1 0412
SCP 1434A-L2 0412
SCP 1434A-L3 0412

17/19

[NON-DETACHABLE CONTAINER LABEL]

GROUP 12 FUNGICIDE

Medallion® WDG

Fungicide

For control of certain foliar stem and root diseases in ornamentals grown in interiorscapes field nursery plantings, container nurseries forest nurseries, residential and commercial landscapes, greenhouses, lath and shade houses, or other enclosed structures

Active Ingredient	
Fludioxonil*	50 0%
Other Ingredients	50 0%
Total	100 0%

*CAS No 131341-86-1

Medallion WDG is a 50% water dispersable granule

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use inside booklet

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard

EPA Reg No 100-1434

EPA Est 67545-AZ-1

SCP 1434A-L1 0412 [Product ID 51008 – Front Booklet/base]
SCP 1434A-L2 0412 [Product ID 51008 – Back Booklet/base]

8 ounces
Net Weight

18/19

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses if present after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment	
NOTE TO PHYSICIAN	
If ingested induce emesis or lavage stomach Treat symptomatically	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals Causes moderate eye irritation Avoid contact with skin eyes, or clothing Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet Remove and wash contaminated clothing before reuse

[For small booklets, use the following reference statement]

Refer to Environmental Hazards, Ground Water Advisory Statement, Physical or Chemical Hazards, and Storage and Disposal sections in attached booklet

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER

19/19

Medallion® and the Syngenta logo are trademarks of a Syngenta Group Company

©2012 Syngenta

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro, North Carolina 27419-8300

MED WDG 1434 MAS 0412 AMEND-B 0812 CLEAN – df – 9/21/12
000100-01434 20120813B MED_WDG-AMEND-B-0812-CLEAN PDF