



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 3, 2025

Cathy Mazo
Regulatory Manager
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419

Subject: Label Amendment - Registration Review Mitigation for Abamectin
Product Name: Epi-Mek SCL
EPA Registration Number: 100-1439
Case Number: 481265
Application Dates: October 3, 2019

Dear Cathy Mazo:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Abamectin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'M. Perch', with a long, sweeping horizontal line extending to the right.

Maryam K. Muhammad-Perch, Team Lead
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

[Master]

<p align="center">RESTRICTED USE PESTICIDE</p> <p align="center">TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.</p>
--

Epi-Mek™ SCL

Miticide/Insecticide

ABAMECTIN	GROUP 6	INSECTICIDE
------------------	----------------	--------------------

Active Ingredient:	
Abamectin ¹	4.2%*
Other Ingredients:	95.8%
Total:	100.0%

¹CAS No. 71751-41-2

*1 gallon contains 0.35 lb abamectin formulated as a suspension concentrate.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1439

EPA Est. XXXX

Product of XXXXX
Formulated in XXXXX

Net Contents

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. <p>Call a poison control center or doctor for treatment advice.</p>
<p style="text-align: center;">NOTE TO PHYSICIAN</p> <p>Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).</p> <p>If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p style="text-align: center;">SYNGENTA HOTLINE NUMBER</p> <p style="text-align: center;">For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed or inhaled. Do not breathe vapor or spray mist. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or Viton™ ≥ 14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish and wildlife.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks to months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of abamectin from runoff water and sediment.

RUNOFF PREVENTION

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Attention: This product contains a chemical known to the state of California to cause birth defects or other reproductive harm.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers are allowed in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short pants and short-sleeved shirt
- Chemical-resistant gloves made of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride \geq 14 mils, or Viton \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, AND/OR ILLEGAL RESIDUES.

USE INFORMATION

Epi-Mek SCL is an aqueous-based soluble concentrate that will control specified pests on the crops listed on this label when the product is applied according to the **Directions for Use**. Thorough coverage of foliage is essential for good mite and insect control.

NOTE:

- **Adjuvant Requirement:** To avoid illegal crop residues, Epi-Mek SCL **must always** be mixed with a non-phytotoxic, non-ionic activator type wetting, spreading and/or penetrating spray adjuvant or horticultural oil (not a dormant oil) as specified in the **Directions for Use** for each crop on this label. Non-ionic activator type wetting, spreading and/or penetrating spray adjuvants include non-ionic surfactants (NIS) with at least 75% surface active agent and crop oil concentrates (COC), vegetable oil concentrates (VOC), methylated seed/vegetable oils (MSO) and organosilicones (OS) with at least 15% emulsifiers/surfactants and include blends of these non-ionic activator type spray adjuvants. Spray adjuvants must be compatible with Epi-Mek SCL and must be used at concentrations specified on the **spray adjuvant product label** directions for use for the targeted crop unless more specific directions are provided in the **Directions for Use** for individual crops on this label. **Do not use binder or sticker type adjuvants because these type adjuvants may reduce translaminar movement of the active ingredient into the plant.** SYNGENTA recommends the use of a Chemical Producers and Distributors Association-certified spray adjuvant.
- **Phytotoxicity and Tank-Mix Compatibility:** Epi-Mek SCL has been tested for phytotoxicity and has a wide margin of safety on a variety of crops. Epi-Mek SCL has also been tested and shown to have broad physical, biological, and non-phytotoxic compatibility with many commonly used pesticides, spray adjuvants and nutritional products. However, since it is not possible to test all potential mixtures, it is recommended that the user contact the supplier of all desired tank-mix products for advice and/or pre-test, prior to commercial application, any previously unproven proposed mixtures with Epi-Mek SCL. Conduct a jar test for physical compatibility of all components of the proposed mixture using proper concentrations of each mixture component. Pre-test the proposed mixture on a small area of the target crop under environmental and application conditions similar to the intended commercial application to ensure physical, biological, and non-phytotoxic compatibility. **Agricultural products formulated with binder or sticker type components may reduce performance of Epi-Mek SCL.**
- **Adjuvant Phytotoxicity Precaution:** Since Epi-Mek SCL must always be mixed with a spray adjuvant as instructed in the directions above, and spray adjuvants alone are known to cause Phytotoxicity to certain crops under certain environmental conditions, do not use Epi-Mek SCL on a spray-adjuvant-sensitive crop unless the spray adjuvant supplier can confirm a known non-phytotoxic labeled use rate for the intended spray adjuvant on the target crop.

- **New York State Restriction:** Do not apply Epi-Mek SCL with aircraft in New York state.
- **Chemigation:** Do not apply this product through any type of irrigation system.

SPRAY DRIFT MANAGEMENT

Spray Drift

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S641).
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must not exceed 65% of the wingspan for fixed wing aircraft or 75% of the rotor diameter for helicopters. Otherwise, the boom length must not exceed 75% of the wingspan for fixed wing aircraft or 90% of the rotor diameter for helicopters.
- Applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field.
- Do not apply during temperature inversions.

Air Blast Applications:

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 mph at the application site
- User must turn off outward pointing nozzles at the row end and when spraying outer row.
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceeds 15 mph at the application site.
- Do not apply during temperature inversions.

NOTE: When states have more stringent regulations, they must be observed.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Spray Drift Precautions for Application with Aircraft or Ground Application Equipment

- Do not apply with ground application equipment within 25 ft of or with aircraft within 150 ft of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries, or commercial fish farm ponds.
- Do not cultivate within 25 ft of the aquatic area to allow growth of a vegetative filter strip.
- Do not allow this product to drift onto non-target areas. Drift may result in illegal residues or injury to non-target species. Risk of exposure to sensitive areas can be reduced by applying this product when the wind direction is away from the sensitive area.
- Do not apply when wind speed is below 2 mph because wind direction will vary and there is a high potential for inversion.
- Observe the following precautions when using ground application to spray tree crops or hops in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish ponds:
 - Do not apply Epi-Mek SCL when weather conditions favor drift to aquatic areas
 - Do not apply within 110 ft upwind of aquatic areas or when wind speed is above 8 mph.
 - Spray last 3 rows windward of aquatic areas using nozzles on one side only, with spray directed away from the aquatic areas.
 - Avoid spray going over tops of trees by adjusting or turning off top nozzles.

Resistance Management

ABAMECTIN	GROUP 6	INSECTICIDE
------------------	----------------	--------------------

Some insect or mite pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects or mites on this label.

Epi-Mek SCL contains a Group 6 miticide/insecticide (abamectin). Insect or mite biotypes with acquired or inherent resistance to Group 6 miticides/insecticides may eventually dominate the pest population if Group 6 miticides/insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Epi-Mek SCL or other Group 6 miticides/insecticides.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect or mite may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

To delay insecticide resistance, take the following steps:

- Rotate the use of Epi-Mek SCL or other Group 6 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.

- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact your local Syngenta representative.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 6 miticides/insecticides exclusively for season-long control of insect or mite species with more than one generation per crop season.
- For insect or mite species with successive or overlapping generations, apply Epi-Mek SCL or other Group 6 miticides/insecticides using a “treatment window” approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated) of the Group 6 miticides/insecticides. Do not exceed the maximum Epi-Mek SCL allowed per growing season.
- Following a treatment window of Group 6 miticides/insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 6 miticides/insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest’s ability to develop resistance to these classes of chemistry.
- If resistance is suspected, do not reapply Epi-Mek SCL or other Group 6 miticides/insecticides.

Other Insect or Mite Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect or mite control program
- Monitoring treated insect or mite populations for loss of field efficacy
- Using tank mixtures or premixes with miticides/insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied

For additional information on Insect or Mite Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

CROP USE DIRECTIONS

APPLES

For control of European red mite, McDaniel spider mite, Tentiform leafminer, Twospotted spider mite, and White apple leafhopper

Apply Epi-Mek SCL using conventional dilute or concentrate ground application sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Apples

Pest	For Dilute Sprays ¹ fl oz/ 100 gal	For Concentrate Sprays ² fl oz/A	Instructions
European red mite McDaniel spider mite Tentiform leafminer Twospotted spider mite White apple leafhopper	1.0 - 2.0	4.5 - 8.5	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a spray adjuvant. Horticultural spray oil (not a dormant oil) is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Apply Epi-Mek SCL when spider mite or insect thresholds are reached. Make a second application, if needed, to maintain control. (See Use Restrictions below.) Spider Mites: For best results, apply before a threshold of 5 spider mites per leaf is reached. Residual spider mite control is greater from spray deposits on newer leaves compared to older. For best results apply Epi-Mek SCL for spider mite control in the tree development period extending from petal fall through 6 weeks following petal fall. Tentiform Leafminer: For best results, apply against egg (to control new hatch) and early sap feeder stages of first- and second-generation tentiform leafminers when locally established thresholds have been reached. Do not apply during bloom.
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	0.25% or 1 gal/A	Minimum of 1 gal/A	

			<ul style="list-style-type: none"> • White Apple Leafhopper (Not for use west of the Rocky Mountains): Application of Epi-Mek SCL is limited only to first-generation white apple leafhoppers. Apply soon after petal fall.
--	--	--	---

¹The rate of Epi-Mek SCL per 100 gallons is based on a volume of 400 gal/A dilute spray.

²To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product/A in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This can result in use of less than 4.5 fl oz/A on small trees.

Use Restrictions – Apples

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the targeted apple variety. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.
 - When used alone or when other products are applied sequentially, the combination of Epi-Mek SCL with horticultural spray oil can injure the fruit of certain apple varieties (e.g., russetting on light-skinned varieties such as Golden Delicious).
 - Applying the combination of Epi-Mek SCL and horticultural spray oil fewer than 14 days before or after applying Captan® or other sulfur-containing products can result in phytotoxicity and crop loss.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 8.5 fl oz/A (or 0.023 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated orchards.

AVOCADOS

For control of Avocado thrips and Persea mite

NOTE: Aerial vs. Ground Application: Apply Epi-Mek SCL using ground application equipment or aircraft. Aerial application is permitted because of its importance to the avocado growing industry. However, aerial application is not the preferred method of application for the best control of thrips. With aerial application, the resulting level and duration of control of thrips could be less than with ground application.

Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good mite and insect control.

Avocados

Pest	Fl oz/A	Instructions
Avocado thrips (<i>Scirtothrips perseae</i>)	4.5 - 8.5	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a spray adjuvant. 1.0% to 4.0% v/v of a horticultural spray oil (not a dormant oil) approved for use on avocados is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) For best results, apply when immature thrips are first observed but before numbers exceed 5 immature thrips per leaf/fruit. Use 4.5-6.5 fl oz of product /A for low* to moderate* infestations and 6.5-8.5 fl oz of product/A for severe* infestations. Make a second application, if needed, to maintain control, but wait at least 30 days after the first application to retreat. (See Use Restrictions below.) For Ground Application: If spray volume is greater than 400 gal/A, apply Epi-Mek SCL at a rate of 1.0 fl oz per 100 gallons of water.
Persea mite (<i>Oligonychus perseae</i>)	Do not use less than 4.5 fl oz of product/A	
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	Minimum 1 to 4% v/v	

*Thrips Infestation Level	Number of immature thrips per leaf/fruit
Low	1-2
Moderate	3-4
Severe	More than 5

Use Restrictions - Avocados

- PHI:** Do not apply within **14 days** of harvest.
- Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use

on avocados. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.

Note: To prevent crop injury, observe all precautions and restrictions on the horticultural spray oil label. When using horticultural spray oil concentrations above 2.0%, treat a small test area before making a large-scale application.

- **Application Method:** Ground and aerial application are permitted.
- **Amount of Water:** Do not apply in less than 100 gallons of water with ground application equipment or 50 gallons of water with aircraft.
- **Maximum Amount per Application:** Do not apply more than 8.5 fl oz/A (or 0.023 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 30 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per season.
- **Maximum Amount per Season:** Do not apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per season.
- **Grazing:** Do not allow livestock to graze in treated orchards.

CELERIAC (*Apium graveollens*)

For control of Twospotted spider mite

Apply Epi-Mek SCL using conventional dilute or concentrate sprayers. Thorough coverage of the crop canopy is essential for optimum results. Inadequate crop coverage can result in reduced control.

Celeriac

Pest	Fl oz/A	Instructions
Twospotted spider mite	7.0	<ul style="list-style-type: none">To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)Apply when mites first appear and repeat as necessary to maintain control. (See Use Restrictions below.)

Use Restrictions – Celeriac

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on celeriac. Carefully follow the **Directions for Use** and **Precautions** on the adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Apply with ground application equipment only.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** After transplanting the celeriac, do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per season.

CITRUS FRUIT CROP GROUP (CROP GROUP 10)

Crops in this group are: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Sour orange, Sweet orange, Pummelo, and Satsuma mandarin.

For control of Asian citrus psyllid, Broad mite, Citrus bud mite, Citrus leafminer, Citrus rust mite, Citrus thrips, and Twospotted spider mite

Thorough spray coverage is essential for good mite and insect control. Adjust gallons of spray per acre based on size and number of trees per acre as well as density of foliage.

- **Ground Application**

Apply using conventional dilute or concentrate ground application sprayers calibrated to deliver sufficient water for thorough coverage (or outside coverage for Asian citrus psyllid, citrus leafminer, or citrus thrips).

- **Aerial Application**

Aerial application is permitted **only** for control of citrus leafminer and Asian citrus psyllid. When using aerial application, the resulting level and duration of control of pests on this label could be reduced compared to ground application. It is recommended to use the higher end of the rate range (7.5 to 8.5 fl oz/A for concentrated sprays) when applying this product by air.

Citrus Fruit Crop Group

Pest	For Dilute Sprays ¹ fl oz/100 gal	For Concentrate Sprays ² fl oz/A	Instructions
Asian citrus psyllid	--	4.5 - 8.5 ³	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a spray adjuvant. It is recommended for best results that a minimum of 0.20% horticultural spray oil (not a dormant oil) be used in the spray mixture or not less than 1.0 gallon of horticultural spray oil/A (except when specified differently in footnotes below). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Asian Citrus Psyllid: Apply Epi-Mek SCL to protect newly expanding foliage flush during the spring, summer or fall. Epi-Mek SCL will only control adults and nymphs present at the time of application. Use 7.5 to 8.5 fl oz/A with aerial application. Mites: Apply when mites first appear during spring, summer, and/or fall. Citrus Bud Mite: For best results, time the spray at "bud swell." Citrus Leafminer: Apply to protect new growth during spring, summer, or fall. Use 7.5 to 8.5 fl oz/A with aerial application. Citrus Thrips: Application targeted for citrus thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun – preferably early to mid-hatch).
Broad mite	0.5 - 0.8	4.5 - 8.5 ³	
Citrus bud mite	0.5 - 0.8	4.5 - 8.5 ⁴	
Citrus leafminer	--	2.0 - 8.5 ³	
Citrus rust mite	0.25 - 0.8	2.0 - 8.5 ³	
Citrus thrips	--	4.5 - 8.5 ⁵	
Twospotted spider mite	0.5 - 0.8	4.5 - 8.5	

¹The rate of Epi-Mek SCL per 100 gallons is based on a volume of 1,000 gal/A dilute spray.

²For concentrate sprays, adjust the dosage to apply an amount/A equal to that used in full cover dilute spray.

³Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural spray oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 MPH.

⁴For best results, use a minimum of 500 gal/A of spray mix with a minimum of 0.5% horticultural spray oil (not a dormant oil).

⁵For best results, use 100-250 gal/A of spray mix applied to achieve outside coverage.

Use Restrictions – Citrus Fruit Crop Group

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the Use Information section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the citrus crop group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Refer to the **Directions for Use** table above for additional spray adjuvant instructions.
- **Application Method:**
 - **To control citrus leafminer and Asian citrus psyllid** – Ground or aerial application is permitted. For aerial application, use a minimum of 10 gallons of finished spray volume per acre. For best results, use at least 25 gallons of finished spray volume per acre. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
 - **All other pests** - Ground application only. Do not apply with aircraft.
- **Prohibited Use:** To manage resistance, do not use in citrus nurseries.
- **Application Interval:** Wait at least 30 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:**
 - **Do not** apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in any growing season.
 - **Do not** make more than 3 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product in any growing season.
- **Grazing:** Do not allow livestock to graze in treated orchards.

COTTON

For control of spider mites

Mix with water as indicated below. Thorough coverage of the crop is essential for optimum results. Inadequate coverage may result in reduced control.

Note:

- Epi-Mek SCL may be applied with ground application equipment or aircraft.
- For best control of spider mites, apply Epi-Mek SCL with ground application equipment. With aerial application, spray coverage and the resulting level and duration of control of mites may be less than with ground application.

Pest	Fl oz/A	Instructions
Carmine spider mite Pacific spider mite Strawberry spider mite Twospotted spider mite	2.0 - 2.5 (Early-Season Cotton - See Instructions) 3.5 - 7.0	<ul style="list-style-type: none">• To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)• Apply when mites first appear. The use rate depends on the size of the plant and density of the foliage.• The lower use rates of 2.0 - 2.5 fl oz/A are only recommended if cotton is early season and under 10 inches in height. Do not use less than 2.0 fl oz/A.• West of the Rocky Mountains - The lower use rates may only be used on cotton less than 10 inches in height and applied only with ground equipment.• Repeat application, if needed, to maintain control. See Use Restrictions below.

Use Restrictions

- **PHI:** Do not apply within **20 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a surfactant approved for use on cotton. Do not use binder or sticker type adjuvants. Carefully follow the **Directions for Use** and **Precautions** on the surfactant label as well as the precautions in official spray guides or on product labels.
- **Application Method:** Ground or aerial application is permitted.
- **Minimum Amount of Water:** Do not apply in fewer than 5 gallons of water per acre. Use sufficient volume to ensure thorough coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application interval:** Wait at least 21 days before repeating application.
- **Maximum Amount per Season:** Do not apply more than 14.0 fl oz/A (or 0.038 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per season.
- **Grazing Restriction:** Do not feed or allow livestock to graze treated cotton.

CUCURBIT VEGETABLES CROP GROUP (CROP GROUP 9)

Crops in this group are: Chayote, Chinese waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourd (includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo*, includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), and Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*).

For Control of Leafminers and Spider Mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control of leafminers and spider mites, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting level and duration of control of leafminers and spider mites could be less than with ground application.

Cucurbit Vegetables Crop Group

Pest	Fl oz/A	Instructions
Leafminers Spider mites	3.5 - 7.0	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz/A for severe infestations. Apply when adult leafminer flies or spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

Use Restrictions – Cucurbit Vegetables Crop Group

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the cucurbit crop group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.

DRY BEANS

Crops include: *Cicer arietinum* (chickpea, garbanzo bean), *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine), *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean), *Vicia faba* (broad bean, fava bean), and *Vigna* spp. (including adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea*, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean)

*For use on cowpeas that are grown only for dry seed. Do not allow livestock to graze cowpea forage and do not harvest cowpea forage or hay for use as livestock feed.

For control of *Liriomyza* leafminers and Spider Mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting control of leafminers and spider mites could be less than with ground application.

Dry Beans

Pest	Fl oz/A	Instructions
<i>Liriomyza</i> leafminers Spider mites	3.5 - 7.0	<ul style="list-style-type: none">• To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)• Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz/A for high infestations.• Leafminers and Spider Mites: Apply when adult leafminer flies or spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

Use Restrictions – Dry Beans

- **PHI:** Do not apply within **7 days** of harvest (7-day PHI).
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the dry bean crop list above. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Amount of Water:** Do not apply in less than 10 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 6 days before repeating application.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.

FRUITING VEGETABLES CROP GROUP (EXCEPT CUCURBITS) (CROP GROUP 8)

Crops in this group are: Eggplant, Groundcherry, Pepino, Peppers (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, and Tomato

For control of Broad mite, Colorado potato beetle, *Liriomyza* leafminers, Spider mites, *Thrips palmi*, Tomato pinworm, Tomato psyllid, and Tomato russet mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control of mites and insects, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting level and duration of control of mites and insects could be less than with ground application.

Fruiting Vegetables Crop Group

Pest	fl oz/A	Instructions
Broad mite Colorado potato beetle <i>Liriomyza</i> leafminers Spider mites Thrips palmi Tomato psyllid Tomato russet mite	3.5 - 7.0	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Broad, russet, and spider mites: Apply when mites first appear. Thrips palmi: Apply when thrips are first observed.
Tomato pinworm	7.0	<ul style="list-style-type: none"> Tomato pinworm: Application can be made from the time moth activity is detected up to, but no later than, the time when newly emerged larvae are present. Mites, leafminers, Thrips palmi, and Colorado potato beetle: Use 3.5 fl oz/A for low to moderate infestations and 7.0 fl oz/A for severe infestations. Repeat application as necessary to maintain control. (See Use Restrictions below.)

Use Restrictions – Fruiting Vegetables Crop Group

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the fruiting vegetables crop group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Prohibited Use:** For resistance management, do not use on fruiting vegetables grown for transplanting.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.

- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.

GRAPES

For control of Pacific spider mite, Twospotted spider mite, Variegated leafhopper, Western grape leafhopper, Western grapeleaf skeletonizer, and Willamette spider mite

Apply Epi-Mek SCL using conventional ground sprayers calibrated to deliver sufficient water for thorough coverage. Thorough coverage is essential for good spider mite and insect control. Do not spray alternate rows. Epi-Mek SCL must be applied to both sides of each row for maximum coverage.

Grapes

Pest	Fl oz/A	Instructions
Pacific spider mite Twospotted spider mite Variegated leafhopper Western grape leafhopper Western grapeleaf skeletonizer Willamette spider mite	3.5 - 7.0	<ul style="list-style-type: none"> • To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) • Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz for severe infestations. • Repeat application if needed. (See Use Restrictions below.) • Spider mites: Apply when mites first appear but before motiles exceed 5 per leaf. • Western grapeleaf skeletonizer: Apply when larvae are first observed. For optimum control, apply shortly after egg hatch. • Leafhoppers: Epi-Mek SCL provides contact knock-down only.

Use Restrictions – Grapes

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the targeted grape variety. Although Epi-Mek SCL has been tested in combination with a range of non-ionic activator type wetting, spreading and/or penetrating spray adjuvants for safety to grapes, it is impossible to test Epi-Mek SCL with all potential spray adjuvants on all grape varieties under the variety of conditions that could cause crop injury. Therefore, when using Epi-Mek SCL, carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment.
- When using an electro-static sprayer, less than 50 gallons of water/A can be used; however, do not use less than 5 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 14.0 fl oz/A (0.038 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated vineyards.

HERB CROP SUBGROUP (CROP SUBGROUP 19A)

Crops in this subgroup are: Angelica, Balm, Basil, Borage, Burnet, Camomile, Catnip, Chervil (dried), Chives, Chives (Chinese), Clary, Coriander (leaf), Cilantro (leaf), Costmary, Culantro (leaf), Curry (leaf), Dill weed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram (*Origanum* species), Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Summer and Winter savory, Sweet bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, and Wormwood.

For control of *Liriomyza* leafminers and Spider Mites

Mix with water as indicated below. Apply thoroughly with ground application equipment as a foliar spray and ensure good upper and lower leaf coverage.

Herb Crop Subgroup (Crop Subgroup 19A)

Pest	Fl oz/A	Instructions
<i>Liriomyza</i> leafminers Spider Mites	3.5 - 7.0	<ul style="list-style-type: none">To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)Use 3.5 fl oz/A for low to moderate infestations and 7.0 fl oz/A for severe infestations. Apply when adult leafminer flies or spider mites are first observed and repeat application, as necessary to maintain control. (See Use Restrictions below.)

Use Restrictions – Herb Crop Subgroup (Crop Subgroup 19A)

- PHI:** Do not apply within **14 days** of harvest. For **chives**, do not apply within **7 days** of harvest.
- Adjuvant Requirement:** To avoid illegal residues Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the herb crop subgroup 19A. Although Epi-Mek SCL has been tested in combination with a range of non-ionic activator type wetting, spreading and/or penetrating spray adjuvants for safety to herbs, it is impossible to test Epi-Mek SCL with all potential spray adjuvants on all herb types and varieties under the variety of conditions that could cause crop injury. Therefore, it is important to carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. Do not use binder or sticker type adjuvants.
- Application Method:** Apply with ground application equipment.
- Amount of Water:** Use a minimum of 20 gallons of water/A.

- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per single cutting (harvest).
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per year.

HOPS (Not approved for use in California)

For control of Twospotted spider mite

Mix with water as indicated below. Apply with ground application equipment as a foliar spray and ensure good upper and lower leaf coverage. Adjust gallons of spray per acre based on the size of plants and density of foliage. In any case, thorough coverage is essential for good control of twospotted spider mite. When using concentrate sprays, the amount of water is reduced, but the amount of Epi-Mek SCL per acre must remain the same as for dilute sprays.

Hops

Pest	Fl oz/A	Instructions
Twospotted spider mite	3.5 - 7.0	<ul style="list-style-type: none"> • To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) • Apply Epi-Mek SCL when twospotted spider mites reach treatment thresholds. • For applications at ½ trellis growth (6 - 8 ft height) apply 3.5 - 7.0 fl oz/A in a minimum of 40 gallons of water/A. • For applications beyond ½ trellis growth, do not use less than 7.0 fl oz/A in a minimum of 100 gallons of water/A.

Use Restrictions – Hops

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on hops. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Application Interval:** If a second application is necessary to maintain control, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product and repeat application of Epi-Mek SCL or any other foliar-applied abamectin-containing product only after an alternative miticide with a different mode of action has been used.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Maximum Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 14.0 fl oz/A (or 0.038 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated hops yards.

LEAFY VEGETABLES (except *Brassica* vegetables) CROP GROUP (CROP GROUP 4)

Crops in this group are: Amaranth (leafy Amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celtuce; Chervil; Chinese celery; Chrysanthemum, edible leaved and garland; Corn salad; Cress, garden and upland (yellow rocket and winter cress); Dandelion; Dock (Sorrel); Endive (escarole); Florence fennel; Lettuce, head and leaf; New Zealand Spinach; Orach; Parsley; Purslane, garden and winter; Radicchio (red chicory); Rhubarb; Spinach; Vine spinach; and Swiss chard.

For Control of Carmine spider mite, *Liriomyza* leafminers, and Twospotted spider mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control of leafminers and spider mites, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting level and duration of control of leafminers and spider mites could be less than with ground application.

Leafy Vegetables Crop Group

Pest	Fl oz/A	Instructions
Carmine spider mite <i>Liriomyza</i> leafminers Twospotted spider mite	3.5 - 7.0	<ul style="list-style-type: none">• To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)• Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz/A for severe infestations.• Spider mites: Apply when mites first appear, and repeat as needed to maintain control. (See Use Restrictions below.)• Leafminers: Apply when adult flies are first observed and repeat applications as needed to maintain control. (See Use Restrictions below.)

Use Restrictions – Leafy Vegetables Crop Group

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the leafy vegetables crop group. Although Epi-Mek SCL has been tested in combination with a range of non-ionic activator type wetting, spreading, and/or penetrating spray adjuvants for safety to leafy vegetables, it is impossible to test Epi-Mek SCL with all potential spray adjuvants on all leafy vegetable types and varieties under the variety of conditions that could cause crop injury. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Prohibited Use:** For resistance management, do not use on leafy vegetables grown for transplanting.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water per acre with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.

MINT (PEPPERMINT AND SPEARMINT)

For control of Twospotted spider mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control of spider mites, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting level and duration of control of spider mites could be less than with ground application.

Mint

Pest	Fl oz/A	Instructions
Twospotted spider mite	3.5 - 5.0	<ul style="list-style-type: none">• To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)• Treat when spider mites first appear.

Use Restrictions – Mint

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on mint. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions such as high temperatures, use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 5.0 fl oz/A (or 0.014 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.

- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product and do not apply more than 3 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per season.
- **Maximum Amount per Season:** Do not apply more than 15.5 fl oz/A (or 0.042 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per crop per season.
- **Grazing:** Do not allow livestock to graze or feed treated foliage to livestock.

ONION, BULB (CROP SUBGROUP 3-07A)

Crops in this group are: onion, bulb including daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; and cultivars, varieties, and/or hybrids of these.

For control of *Liriomyza* Leafminers and Thrips

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL with ground application equipment or aircraft.
- For best control, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting control of leafminers and thrips could be less than with ground application.

Onion, Bulb (Crop Subgroup 3-07A)

Pest	Fl oz/A	Instructions
<i>Liriomyza</i> leafminers Thrips	3.5 - 7.0	<ul style="list-style-type: none"> • To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) • Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz/A for high infestations. • Insect control can be reduced if Epi-Mek SCL is used in combination with a sticker or binder type product such as Bravo® WeatherStik®. • Leafminers: Apply when adult leafminer flies are first observed and repeat application as needed to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

		<ul style="list-style-type: none"> • Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.) Do not use Epi-Mek SCL as a rescue treatment for thrips control.
--	--	--

Use Restrictions – Onion, Bulb (Crop Subgroup 3-07A)

- **PHI:** Do not apply within **30 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on bulb onions or on the other intended target crop in the bulb onion crop subgroup 3-07A. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker or type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** Make 2 consecutive applications of Epi-Mek SCL, then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional Epi-Mek SCL applications. Do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.

PEARS (INCLUDING ORIENTAL PEAR TREES)

For control of European red mite, McDaniel spider mite, Pear psylla, Pear rust mite, Twospotted spider mite, and Yellow mite

Apply Epi-Mek SCL using conventional dilute or concentrate ground application sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Pears

Pest	For Dilute Sprays ¹ fl oz/ 100 gal	For Concentrate Sprays ² fl oz/A	Instructions
European red mite McDaniel spider mite Pear psylla Pear rust mite Twospotted spider mite Yellow mite	1.0 2.0	4.5 - 8.5	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a spray adjuvant. Horticultural spray oil (not a dormant oil) is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	0.25% or 1 gal/A	Minimum of 1 gal/A	<ul style="list-style-type: none"> Apply Epi-Mek SCL when spider mite or insect thresholds are reached. Make a second application, if needed, to maintain control. (See Use Restrictions below.)

¹The rate of Epi-Mek SCL per 100 gallons is based on a volume of 400 gal/A dilute spray.

²To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product/A in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This can result in use of less than 4.5 fl oz/A on small trees.

Use Restrictions – Pears

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the Use Information section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on pears. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.

- Applying the combination of Epi-Mek SCL and horticultural spray oil fewer than 14 days before or after applying Captan® or other sulfur-containing products can result in phytotoxicity and crop loss.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 8.5 fl oz/A (or 0.023 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated orchards.

STONE FRUIT CROP GROUP (CROP GROUP 12)

Crops in this group are: apricot, cherry (sweet and tart), nectarine, peach, plum, plum (chickasaw, damson, and Japanese), plumcot, and prune (fresh)

For control of European red mite, Pacific spider mite, and Twospotted spider mite

Apply Epi-Mek SCL using conventional dilute or concentrate sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite control.

Stone Fruit Crop Group

Pest	For Dilute Sprays ¹ fl oz/ 100 gal	For Concentrate Sprays ² fl oz/A	Instructions
European red mite Pacific spider mite Twospotted spider mite	1.0 2.0	4.5 - 8.5	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant or with a horticultural spray oil (not a dormant oil). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Apply when mites first appear. Make a second application, if needed, to maintain control. (See Use Restrictions below.)
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	0.25% or 1 gal/A	Minimum of 1 gal/A	<ul style="list-style-type: none"> A horticultural spray oil (not a dormant oil) can improve efficacy but it could also increase the possibility of phytotoxicity to foliage and fruit. Use a minimum of 0.25% horticultural spray oil (not a dormant oil) in the dilute spray mixture and not less than 1.0 gallon of horticultural spray oil per acre in the final finished spray.

¹The rate of Epi-Mek SCL per 100 gallons is based on a volume of 400 gal/A dilute spray.

²To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This can result in use of less than 4.5 fl oz/A on small trees.

Use Restrictions – Stone Fruit Crop Group

- **PHI:** Do not apply within **21 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the stone fruit crop group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. See the **Directions for Use** table above for application instructions. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 8.5 fl oz/A (or 0.023 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is needed, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated orchards.

STRAWBERRIES

For control of Strawberry spider mite, Twospotted spider mite, and Carmine spider mite

Pest	Fl oz/A	Instructions
Strawberry spider mite Twospotted spider mite Carmine spider mite	7.0	<ul style="list-style-type: none">To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.)Make 2 applications 7-10 days apart when mites first appear. Repeat this application sequence, if needed, to maintain control. (See Use Restrictions below.)

Use Restrictions – Strawberries

- **PHI:** Do not apply within **3 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a surfactant approved for use on strawberries. Do not use binder or sticker type adjuvants. Carefully follow the **Directions for Use** and **Precautions** on the surfactant label, as well as the precautions in official spray guides on product labels.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Prohibited Use:** For resistance management purposes, do not use in strawberry nurseries.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using electro-static sprayers, do not apply in less than 10 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 21 days after the second application before repeating application.
- **Maximum Amount per Season:** Do not apply more than 28 fl oz/A (or 0.075 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.

TREE NUTS CROP GROUP (CROP GROUP 14) AND PISTACHIO

Crops in this group are: almond, beech nut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia nut, pecan, and walnuts (black and English)

For control of European red mite, Pacific spider mite, Strawberry spider mite, and Twospotted spider mite

Apply Epi-Mek SCL using conventional dilute or concentrate sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Tree Nuts Crop Group and Pistachio

Pest	For Dilute Sprays ¹ fl oz/ 100 gal	For Concentrate Sprays ² fl oz/A	Instructions
European red mite Pacific spider mite Strawberry spider mite Twospotted spider mite	1.0 2.0	4.5 - 8.5	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a spray adjuvant. Horticultural spray oil (not a dormant oil) is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Apply Epi-Mek SCL when spider mites first appear. Residual spider mite control is greater from spray deposits on newer leaves compared to older leaves. Make a second application, if needed, to maintain control. (See Use Restrictions below.)
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	0.25% or 1 gal/A	Minimum of 1 gal/A	

¹The rate of Epi-Mek SCL per 100 gallons is based on a volume of 400 gal/A dilute spray.

²To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. This can result in use of less than 4.5 fl oz/A on small trees.

Use Restrictions – Tree Nuts Crop Group and Pistachio

- **PHI:** Do not apply within **21 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a spray adjuvant as instructed in the Use Information section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the tree nuts crop group or on pistachio. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Application:** Do not apply more than 8.5 fl oz/A (or 0.023 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per application.
- **Applications per Season:** Do not make more than 2 applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product per growing season.
- **Maximum Amount per Season:** Do not apply more than 17.0 fl oz/A (or 0.047 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product in a growing season.
- **Grazing:** Do not allow livestock to graze in treated groves/orchards.

TUBEROUS AND CORM VEGETABLES CROP SUBGROUP (CROP SUBGROUP 1C)

Crops in this subgroup are: arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava (bitter and sweet); chayote (root); chufa; dasheen; ginger; leren; potato; sweet potato; tanier; turmeric; yam bean; and yam (true).

For control of Colorado potato beetle, *Liriomyza* leafminers, Potato psyllid, and Spider mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Epi-Mek SCL using ground application equipment or aircraft.
- For best control of mites, apply Epi-Mek SCL with ground application equipment. With aerial application, the resulting level and duration of control of insects and spider mites could be less than with ground application.

Tuberous and Corm Vegetables Crop Subgroup

Pests	Fl oz/A	Instructions
Colorado potato beetle <i>Liriomyza</i> leafminers Potato psyllid Spider mites	3.5 - 7.0	<ul style="list-style-type: none"> To avoid illegal residues, Epi-Mek SCL must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Epi-Mek SCL label and Use Restrictions below.) Use 3.5 - 5.0 fl oz/A for low to moderate infestations and 7.0 fl oz/A for severe infestations. Insect and spider mite control can be reduced if Epi-Mek SCL is used in combination with a binder or sticker type product such as Bravo® Weather Stik®. Colorado potato beetle: Make the first application after approximately 50% of the egg masses have hatched and larvae are present. If 2 applications are needed, limit them to a single Colorado potato beetle generation per crop. (See Use Restrictions below.) <i>Liriomyza</i> leafminers: Make the first application when adult flies are first observed. Repeat applications as needed to maintain control. (See Use Restrictions below.) Spider mites: Make the first application when mites first appear. Repeat application as needed to maintain control. (See Use Restrictions below.)

Use Restrictions – Tuberous and Corm Vegetables Crop Subgroup

- **PHI:** Do not apply within **14 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Epi-Mek SCL **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Epi-Mek SCL label. The spray adjuvant must be approved for use on the intended target crop in the tuberous and corm vegetables crop subgroup. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. See the **Directions for Use** table above for application instructions. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York state.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 7.0 fl oz/A (or 0.019 lb ai/A) per application.

- **Application Interval:** Wait at least 7 days before repeating application of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Epi-Mek SCL or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Crop:** Do not apply more than 20.5 fl oz/A (or 0.056 lb ai/A) of Epi-Mek SCL or any other foliar-applied abamectin-containing product per crop.
- **Grazing:** Do not allow livestock to graze or feed treated foliage to livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a tightly closed container in a cool, dry place.

Pesticide Disposal

Pesticide waste may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Epi-Mek™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Bravo® and Weather Stik® are trademarks of Makhteshim Agan of North America, Inc.
d/b/a ADAMA.

Captan® is a trademark of Tomen Agro, Inc.

Viton™ is a trademark of The Chemours Company FC, LLC

©20XX Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

Epi-Mek SCL 1439 MAS 0714 AMEND-B OCT2019-CL – jvb – 5/9/2024
000100-01439.20191003B.Epi-Mek SCL.AMEND-OCT2019-CL.pdf