



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1560

Date of Issuance:

8/28/15

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

SCHOLAR RTU

Name and Address of Registrant (include ZIP Code):

Teresa S. Cox
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 100-1560."
 - Add an appropriate EPA Establishment Number to the label.

Signature of Approving Official:

Hope Johnson, Product Manager 21
Fungicide Branch, Registration Division (7505P)

Date:

8/28/15

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/13/2015
- Alternate CSF 1 dated 08/13/2015

If you have any questions, you may contact Maryam K. Muhammad at 703-347-0301 or via email at Muhammad.maryam@epa.gov.

Enclosure: Stamped Accepted "Scholar RTU" Label
Acute Toxicity Review
Product Chemistry Review

Scholar® RTU

Fungicide

Scholar RTU is ready-to-use liquid fludioxonil for post-harvest thermal fogging on pome fruit.

Active Ingredient:

Fludioxonil* 10.0%

Other Ingredients: 90.0%

Total: 100.0%

*CAS No. 131341-86-1

Scholar RTU is a 10% hot fogging concentrate.

Scholar RTU contains 0.795 pounds of fludioxonil per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION


See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-XXXX

EPA Est.

2.5 gallons

Net Contents



| FIRST AID | |
|---|---|
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Harmful if inhaled or absorbed through skin. Avoid breathing vapor. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Handlers (mixers, loaders, persons cleaning or setting up the fogging equipment, and persons handling treated fruit prior to drying) must wear:

- Long-sleeved shirt and long pants
- Protective eyewear (goggles, safety glasses, or a face shield)
- Chemical-resistant gloves made from any waterproof material: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton® ≥ 14 mils
- Shoes plus socks

In addition, fumigating applicators must wear:

- Full-face respirator with organic vapor filter

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Physical or Chemical Hazards

This product contains a highly flammable ingredient. It may cause a fire or explosion if not used properly. Follow the "Directions for Use" on this label very carefully.



**Highly Flammable Ingredient
Ingrediente Altamente Inflamable**

Combustible. Do not use or store near heat or open flame. Do not use with or store near any oxidizing. Do not use with or store near water or 10% mono-ammonium phosphate solution (fire extinguishant).

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be

modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND RESTRICTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL.

PRODUCT INFORMATION

A20335 is a protective fungicide used to aid in the control of post-harvest diseases in pome fruit in storage facilities.

RESISTANCE MANAGEMENT

Scholar RTU contains fludioxonil which is in the phenylpyrrole class of chemistry and has a unique mode of action, inhibition of a MAP/histidine kinase (FRAC Group 12). Fungal isolates with acquired resistance to Group 12 may eventually dominate the fungal population if Group 12 fungicides are used repeatedly in the same field or packing facility or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by fludioxonil or other Group 12 fungicides. A disease management program that includes alternation or tank mixes between Scholar RTU and other labeled fungicides that have a different mode of action may prevent pathogen populations from developing resistance. Use sanitation and other cultural practices to minimize pathogen populations in order to control disease and prevent or delay resistance development.

MIXING PROCEDURES

If using Scholar RTU in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label. Do not exceed any label dosage and apply the most restrictive label precautions. Do not mix Scholar RTU with any other product whose label prohibits such mixing. Tank mixtures are permitted only in those states where the tank-mix partner is registered.

THE CROP SAFETY OF ALL POTENTIAL TANK MIXES INCLUDING ADDITIVES AND OTHER PESTICIDES ON ALL CROPS HAS NOT BEEN TESTED. BEFORE APPLYING ANY TANK MIXTURE, THE SAFETY TO THE TARGET CROP SHOULD BE CONFIRMED.

APPLICATION INSTRUCTIONS

Apply Scholar RTU at rates and timings as described in this label.

Follow all label directions and restrictions for each ingredient in the tank mixture.

Preparation for Scholar RTU Treatment

Posting

Notify workers of application by warning them orally and posting warning signs at entrances to treated areas.

Re-Entry

Entry into the treatment area by another person other than properly-trained and -equipped handlers is prohibited from the start of application until the treated area is ventilated as follows:

- 8 hours with no ventilation followed by 1 hour of mechanical ventilation
- 24 hours with no ventilation

In case it is necessary to enter the treatment area during treatment or before ventilation requirements have been met, handlers must wear chemical resistant headgear and a self-contained breathing apparatus (SCBA, MSHA/NIOSH approval number prefix TC-13F) in addition to PPE listed under the PRECAUTIONARY STATEMENTS. Follow all applicable safety precautions for entering a controlled atmosphere storage facility, such as monitoring oxygen and carbon monoxide meters to assure safe levels before re-entry.

Application Procedures

- Do not breathe the fog.
- Turn off room cooling systems and humidifiers 12 hours prior to and during treatment.
- Turn off room circulation fans immediately prior to and during treatment.
- If necessary, open a small vent in the room during fogging to avoid back-pressure.
- Turn on room circulation fans one hour after the treatment.

- Visually check that the fog has totally disappeared (about 5 hours post-treatment) before restarting the cooling systems.

Note: When rooms are treated with less than 75% fruit capacity, calculate the dosage by adding a tonnage value of 20% of the empty volume to the real tonnage of the fruit.

Thermal Fogger Operation

- Place the thermal fogger machine horizontally outside the treatment room at a height of about 3 feet from the floor. Insert the fogging pipe through the porthole in the door to the storage room.
- Seal the opening around the nozzle with thermally resistant tape.
- Adjust the pump and mount the nozzle to deliver up to 120 mL per minute.
- Turn on the thermal fogger machine.
- Insert the insulated input pipe into the mixture. Start the pump.
- If used in tank mix, ensure the mixture is agitated during application.
- Maintain fogger inlet temperature between 500-600°F. Do not exceed 650° F fogger temperature (i.e., inner temperature of the fogger barrel at the point of injection).
- Do not use any elbow or allow any restriction at the end of the fog barrel which might cause back-pressure or increase the temperature in the fogger barrel.

USE RESTRICTIONS

- Only apply 1 post-harvest application of fludioxonil-containing product.
- Do not apply to fruit previously treated with fludioxonil via drench or dip/wash.
- Apply only with a thermal fogging machine.
- Only other fungicides compatible with Scholar RTU, diphenylamine (DPA), and approved for fogging applications can be tank mixed with Scholar RTU.
- Do not exceed 650° F fogger temperature (i.e., inner temperature of the fogger barrel at the point of injection).
- Do not use any elbow or allow any restriction at the end of the fog barrel which might cause back-pressure or increase the temperature in the fogger barrel.

CROPS USE DIRECTIONS - POST-HARVEST

Pome Fruit

Complete List of Pome Fruit: Apple (*Malus domestica*); Azarole (*Crataegus azarolus*); Crabapple (*Malus* spp.); Loquat (*Eriobotrya japonica*); Mayhaw (*Crataegus aestivalis*, *C. opaca*, and *C. rufula*); Medlar (*Mespilus germanica*); Pear (*Pyrus communis*); Pear, Asian (*Pyrus* spp.); Quince (*Cydonia oblonga*); Quince, Chinese (*Chaenomeles speciosa*); Quince, Japanese (*Chaenomeles japonica*); Tejocote (*Crataegus mexicana*) and cultivars, varieties and/or hybrids of these.

Use Scholar RTU as a post-harvest thermal-fog aerosol application to reduce damage from post-harvest diseases caused by:

| Application Method | Disease | Rate ml (grams) | Remarks |
|---|--|---|--|
| Fogging into Storage Facility | Blue mold Gray mold Bitter rot Speck rot White rot Phacidiopycnis rot Sphaeropsis rot Alternaria rot and surface mold Rhizopus rot Bull's-eye rot | 10.5 ml/bin or 1 gram fludioxonil /bin (900 lb of fruit /bin) | <ul style="list-style-type: none"> • Ensure proper distribution of the product on the fruit. • For best efficacy, treat only dry fruit. • Apply to fruit as soon as possible after harvest, and ideally within 15 days (after harvest). • Load bins in the storage room with normal spacing between bins and head space. • For best efficacy, do not cover top bins. • Cover the front face of the bins directly in front of the fogging nozzle. |
| <p>For maximum decay control during storage, treat fruit before storage with Scholar RTU.</p> <p>Specific Use Restrictions</p> <ol style="list-style-type: none"> 1) Do not apply to fruit previously treated with fludioxonil via drench or dip/wash. 2) Apply only with a thermal fogging machine. 3) Do not apply more than 1 post-harvest application of fludioxonil-containing product. | | | |

Conversion Chart

| Number of 900 lb Fruit Bins | US Tons of Fruit | Amount of Scholar RTU in liters (grams of fludioxonil) |
|------------------------------------|-------------------------|---|
| 100 | 45 | 1.05 (100) |
| 500 | 225 | 5.25 (500) |
| 1000 | 450 | 10.50 (1000) |
| 1500 | 675 | 15.75 (1500) |
| 2000 | 900 | 21.00 (2000) |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers in a cool, dry place. Do not store this product under wet conditions. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons - bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Scholar®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company.

Viton® is a trademark of E.I. du Pont de Nemours and Company.

©20XX Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCHOLAR RTU (A20335D) – NEW – C - CLEAN – df – 8/21/15
000100-XXXXX.20141201C.SCHOLAR_RTU-A20335D-NEW-CLEAN.PDF