



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1597

Date of Issuance:

12/19/16

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Clariva[®] Complete Beans 500

Name and Address of Registrant (include ZIP Code):

Patrick McCain
Senior Regulatory Product Manager
Syngenta Crop Protection
P.O. Box 18300
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate-Vertebrate Branch 3, Registration Division (7505P)

Date:

12/19/16

2. You are required to comply with the data requirements described in the DCI's identified below:
 - a. Thiamethoxam GDCI-060109-1309
 - b. Mefenoxam GDCI-113502-940
 - c. Fludioxonil GDCI-071503-1243

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: http://www.epa.gov/oppsrrd1/contacts_prd.htm

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 100-1597."
4. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/29/2016
- Alternate CSF dated 08/29/2016

If you have any questions, please contact Dee Colby by phone at 703-347-8657, or via email at colby.deanna@epa.gov.

Enclosure: stamped label



(Master label)

GROUP	4A	INSECTICIDE
GROUP	4 7 12	FUNGICIDES

Clariva® Complete Beans 500

Insecticide and Nematicide with Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, seedling diseases and Soybean Cyst Nematode on soybeans

Active Ingredients:

Thiamethoxam ¹	12.50%
Mefenoxam ²	1.88%
Fludioxonil ³	0.63%
Sedaxane ⁴	0.63%
<i>Pasteuria nishizawae</i> – Pn1 ^{*5}	5.07%

Other Ingredients: 79.29%

Total: 100.00%

* Contains at least 3.34 x 10⁹ spores per mL or 1.3 x 10¹³ spores per gal.

¹CAS No. 153719-23-4

²CAS No. 70630-17-0 and CAS No. 69516-34-3

³CAS No. 131341-86-1

⁴CAS No. 874967-67-6

⁵CAS No. None

Clariva Complete Beans 500 contains 1.14 lb thiamethoxam, 0.17 lb mefenoxam, 0.06 lb fludioxonil and 0.06 lb sedaxane per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

[See additional precautionary statements and directions for use [in booklet] [on label].

EPA Reg. No. 100-

EPA Est. XXXXX

SCP

Net Contents

FIRST AID	
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton® ≥ 14 mils
- Shoes plus socks
- A NIOSH-approved particulate respirator with any R or P filter with NIOSH approval number prefix TC-84A or a NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C.

Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife, freshwater and estuarine/marine fish, oysters, and shrimp and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Groundwater Advisory

Fludioxonil and Thiamethoxam have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame. Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities only. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.
This product is to be used in liquid or slurry treaters only.

Do not use this product until you have read the entire label. Keep unprotected persons out of treatment area until seeds have dried or been packaged.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with Clariva Complete Beans 500.

USE INFORMATION

Clariva Complete Beans 500 is a seed treatment product containing the active ingredients thiamethoxam (insecticide) and fludioxonil, mefenoxam, sedaxane (fungicides) and *Pasteuria nishizawae* – Pn1 (nematicide). Clariva Complete Beans 500 protects against damage from listed early-season insects, soil-borne and seed-borne diseases on soybeans as well as offers season-long protection from soybean cyst nematode (*Heterodera glycines*)

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam protects against listed chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against *Pythium*, *Phytophthora* and systemic downy mildew.

Fludioxonil fungicide is active against *Fusarium* and *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Sedaxane fungicide is active against seed decay, seedling blight and damping-off caused by *Rhizoctonia* species.

Pasteuria nishizawae (Pn1) is a biological nematicide for protection against soybean cyst nematode (*Heterodera glycines*)

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum seasonal total for the active ingredient as stated on the label of the product containing the lowest seasonal total on that crop.

RESISTANCE MANAGEMENT

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

Clariva Complete Beans 500 contains thiamethoxam, a Group 4A insecticide. Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists.

Insect populations may contain individuals naturally resistant to Group 4A insecticides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refugia (untreated areas). Sound management programs also consider cultural and biological control practices.

In order to maintain susceptibility to this class of chemistry:

- Use products at their full, specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless

otherwise excluded by product labels. Do not exceed the maximum amount of this insecticide's mode of action allowed per growing season.

- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

GROUP	4	7	12	FUNGICIDES
--------------	----------	----------	-----------	-------------------

Clariva Complete Beans 500 contains mefenoxam, a Group 4 fungicide; fludioxonil, a Group 12 fungicide; and sedaxane, a Group 7 fungicide. Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with a MAP/histidine protein kinase in osmotic signal transduction. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry which disrupts cellular respiration and energy generation.

Fungal populations may contain individuals naturally resistant to Group 4, 7, and 12 fungicides, and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as alternation with fungicides with a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Insecticide or Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org> or the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>

MIXING PROCEDURES

Important: Always re-circulate Clariva Complete Beans 500 thoroughly before using.

Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Apply Clariva Complete Beans 500 as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Clariva Complete Beans 500 into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. Clariva Complete Beans 500 is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Syngenta representative for directions before applying Clariva Complete Beans 500 with inoculants.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution. Clean out any unused product from the treater after treating or maintain constant agitation if the leftover slurry will be maintained overnight.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Use an EPA approved dye/colorant that imparts an unnatural color to the seed as required in 40CFR 153.155(c).

Allow seed to dry before bagging.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting.

USE RESTRICTIONS

- Do not allow children, pets or livestock to have access to treated seed.
- Do not graze or feed livestock on treated areas for 45 days after planting seed treated with Clariva Complete Beans 500.
- Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 g) per acre, 0.004 lb fludioxonil (2.0 g) per acre and 0.013 lb mefenoxam (5.7 g) per acre per calendar year as a seed treatment application.

- This seed has been treated with 0.0756 mg thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed.
- Do not use more than 0.125 lb of thiamethoxam (56.7 g) per calendar year, including all thiamethoxam products and application methods (for example, seed treatment, in-furrow, and foliar).
- Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with Clariva Complete Beans 500.

CROP ROTATIONAL RESTRICTIONS

In the event of crop failure or harvest of a crop grown from Clariva Complete Beans 500 treated seed, the field may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Canola Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat Soybean	Alfalfa <i>Brassica</i> (Cole) Leafy Vegetables Crop Group 5 Cereal Grains: Buckwheat, Corn, Pearl Millet, Proso Millet, Popcorn, Rice (dry-seeded), Sorghum, Teosinte, Wild Rice Cotton Cucurbit Vegetables Crop Group 9 Fruiting Vegetables Crop Group 8 Leafy Vegetables Crop Group 4 Legume Vegetables (Succulent or Dried) Crop Group 6 Mint: Peppermint and Spearmint Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower Onion, Bulb Peanut Root Vegetables (including Sugarbeet) Crop Subgroup 1A Strawberry Sunflower Tobacco Tuberous and Corm Vegetables (including Potato) Crop Subgroup 1C

- For any other crops the minimum plant back interval is 120 days from the date the Clariva Complete Beans 500 treated seeds were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide, fludioxonil, mefenoxam, sedaxane fungicides, and *Pasteuria nishizawae* - Pn1 nematicide.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with Clariva Complete Beans 500:

- Groundwater Advisory:
Fludioxonil and Thiamethoxam have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Pollinator Precautions:
Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.

- Do not graze or feed livestock on treated areas for 45 days after planting seed treated with Clariva Complete Beans 500.
- In the event of a crop failure or harvest of a crop grown from Clariva Complete Beans 500 treated seed, the field may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Canola Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat Soybean	Alfalfa <i>Brassica</i> (Cole) Leafy Vegetables Crop Group 5 Cereal Grains: Buckwheat, Corn, Pearl Millet, Proso Millet, Popcorn, Rice (dry-seeded), Sorghum, Teosinte, Wild Rice Cotton Cucurbit Vegetables Crop Group 9 Fruiting Vegetables Crop Group 8 Leafy Vegetables Crop Group 4 Legume Vegetables (Succulent or Dried) Crop Group 6 Mint: Peppermint and Spearmint Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower Onion, Bulb Peanut Root Vegetables (including Sugarbeet) Crop Subgroup 1A Strawberry Sunflower Tobacco Tuberous and Corm Vegetables (including Potato) Crop Subgroup 1C

- For any other crops the minimum plant back interval is 120 days from the date the Clariva Complete Beans 500 treated seeds were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

- Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 g) per acre, 0.004 lb fludioxonil (2.0 g) per acre and 0.013 lb mefenoxam (5.7 g) per acre per calendar year as a seed treatment application.
- This seed has been treated with 0.0756 mg ai thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed.
- Do not use more than 0.125 lb of thiamethoxam (56.7 g) per calendar year, including all thiamethoxam products and application methods (for example, seed treatment, in-furrow, and foliar).
- Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with Clariva Complete Beans 500.

CROP USE DIRECTIONS

When applied according to the **CLARIVA COMPLETE BEANS 500 RATE TABLE**, Clariva Complete Beans 500 provides early season protection against injury by aphids, bean leaf beetle, grape colaspis, leaf miners, leaf hoppers, Mexican bean beetle, seed corn maggot, three-cornered alfalfa hopper, thrips, white grubs, and wireworm.

Clariva Complete Beans 500 provides protection against damping-off and seed borne rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* species and early season *Phytophthora* root rot . Clariva Complete Beans 500 also suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Clariva Complete Beans 500 provides protection against soybean cyst nematode.

Clariva Complete Beans 500 Rate Table

Crop	Rate of Clariva Complete Beans 500					
	fl oz per 100 lb seed ¹ or fl oz per 140,000 seeds ¹		g ai per 100 kg		mg ai per seed	
Soybean including soybean, vegetable	Clariva Complete Beans 500	Apron XL	Clariva Complete Beans 500	Apron XL	Clariva Complete Beans 500	Apron XL
		5.6 fl oz/100 lb seed or 2.6 fl oz per 140,000 seeds	0.32 fl oz Apron XL/100 lb seed or 0.15 fl oz Apron XL/140,000 seeds	Thiamethoxam 50 g Mefenoxam 7.5 g Fludioxonil 2.5 g Sedaxane 2.5 g Pasteuria nishizawae Pn1 20 g Total = 82.5 g	Mefenoxam 7.5 g	Total of all active ingredients = 0.125 mg

¹The mg ai per seed, fl oz 500 per 100 lb seed, and fl oz per 140,000 seeds rates are based on 3,000 seeds per pound.

Additional Apron® XL or similar mefenoxam product, may be necessary for high levels of *Phytophthora* or *Pythium*. Refer to the table above to determine the appropriate use rate of Apron XL, if needed. Read and follow all label direction for Apron XL use.

For additional protection from early season infection of *Fusarium virguliforme* (causal organism of Sudden Death Syndrome or SDS), add 20 g ai/100 kg seed or 0.64 fl oz /100 lb seed of Mertect® 340-F (or 0.29 fl oz /140,000 seeds of Mertect 340-F).

STORED GRAIN PROTECTION

When treated according to the directions for post-planting protection against listed pests, Clariva Complete Beans 500 will also provide protection during post treatment storage of the seed listed on this label against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If the seed to be treated has existing infestations of stored grain insects, fumigate the seed with a registered product approved for such use prior to treating with Clariva Complete Beans 500 and bagging.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store between a range of 32°F to 100°F.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Apron®, Clariva®, Mertect®, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Viton® trademark of E. I. du Pont de Nemours & Company, Inc.

©20XX Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

CLARIVA COMPLETE BEANS 500 XXXX MAS NEW-D-Clean – rk – 12-12-16.docx
000100-XXXXX.20151112D.CLARIVA COMPLETE BEANS _NEW-D_NOV2015.CL.pdf