

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Biopesticides and Pollution Prevention Division (7511P) 1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

EPA Reg. Number:	Date of Issuance:
100-1599	7/27/2016

Term of Issuance:

Unconditional

Name of Pesticide Product:

Inosco

Name and Address of Registrant (include ZIP Code):

Syngenta Crop Protection, LLC PO Box 18300 Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
- 2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Date:

7/27/2016

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)
Office of Pesticide Programs

- 3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 100-1599."
- 4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

Basic CSF dated 04/26/2016

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Menyon Adams of my team by phone at (703) 347-8496 or via email at adams.menyon@epa.gov.

Sincerely,

Andrew Bryceland, Team Leader Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511P)

andrew . Buycelow

Office of Pesticide Programs

Enclosure

Inosco™

A systemic fungicide for the control of downy mildew, damping off, late blight, root rot, and suppression of bacterial blight in the production of ornamental plants, ornamental bulbs, corms and tubers, vegetables and vegetable plants grown in greenhouses and nurseries (including field and container grown plants grown outdoors and in shade houses, lath houses and other ornamental production structures) and conifer nurseries

ACTIVE INGREDIENT:

Potassium Phosphite*	54.50%
OTHER INGREDIENTS:	45.50%
TOTAL:	100.0%

^{*}Phosphorous acid equivalent: 34.30% (4.2 lbs/gal).

KEEP OUT OF REACH OF CHILDREN CAUTION

	CACTION	
	FIRST AID	
If swallowed	 Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person. 	
If in eyes	 Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. 	
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 	
	 Call a poison control center or doctor for treatment advice. 	
If on skin or	Take off contaminated clothing.	
clothing	 Rinse skin immediately with plenty of water for 15 – 20 minutes. 	
	Call a poison control center or doctor for treatment advice.	
LIOT LINE NUMBER		

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-888-8372 for emergency medical information.

See inside booklet for additional Precautionary Statements and for Directions for Use.

EPA Reg. No.:	
EPA Est. No.:	

Manufactured for:

Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina 27419-8300 Customer Service 1-800-334-9481 SCP XXXX-XX

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE: 1-800888-8372

ACCEPTED

07/27/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 100-1599

Net Contents:

AGRICULTURAL CHEMICAL DO NOT SHIP OR STORE WITH FOODS. FEEDS, DRUGS OR CLOTHING

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals:

CAUTION. Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 4 hours unless wearing appropriate PPE.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- coveralls
- waterproof gloves
- shoes plus socks

Chemigation:

General Chemigation Information

- 1. Apply this product only through drip (trickle), microjet, center pivot, solid set, hand move, or moving wheel irrigation system that does not contain aluminum components.
- Do not apply this product through any other type of irrigation system.
- 2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.
- 4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- 6. Equip the pesticide supply tank with a means for continuous agitation either recirculation or mechanical agitator. Fill the supply tank with the appropriate amount of water and add the Inosco slowly followed by any other products that are compatible and labeled for chemigation. Observe all cautions and limitations on the label of all products used in the mixtures. For fixed position irrigation systems such as center pivot etc., apply the pesticide toward the end of the irrigation cycle. Exact timing will depend upon the desired Inosco application rate and the calibration of the system. For moving systems, apply the Inosco continuously. In all cases, maintain thorough coverage of the crop.

For Chemigation Systems Connected to Public Water Systems:

- 1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the reduced-pressure zone, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- 4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

For Drip (Trickle) Chemigation and Sprinkler Chemigation Systems:

- 1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- 2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Product Information:

Inosco is a systemic product, which contains phosphorous acid. Phosphorous acid is effective in prevention and control of diseases by activation of the plants natural resistance mechanism, as well as by direct activity on the fungus. Inosco is intended for use as part of an Integrated Pest Management (IPM) system. When used in conjunction with good cultural management practices and as part of a complete disease control program, Inosco will control downy mildew, damping off, root rot and late blight, and suppression of bacterial blight in the production of ornamental plants, ornamental bulb, corm and tuber crops, vegetables and vegetable plants grown in greenhouses and nurseries (including field and container grown plants grown outdoors and in shade houses, lath houses and other ornamental production structures and conifer nurseries. In order to achieve maximum results in controlling the listed diseases, treatments should be initiated when environmental conditions are favorable for disease development.

Under periods of high disease pressure, Inosco may be tank-mixed and/or used ion rotation with other fungicides, such as Heritage[®] Fungicide, Micora[™] Fungicide or Subdue MAXX[®] Fungicide, per label directions, for improved disease control.

Mixing Instructions:

Fill spray tank halfway with clean water. Add the required amount of Inosco to the tank as indicated in Table 1. While agitating, add the remaining amount of water.

Avoid mixing Inosco with products that are either strongly acidic (pH < 5.5) or strongly alkaline (pH > 7.8).

Compatibility:

Inosco is believed to be compatible with most commonly used pesticides. Consult specific product labels for additional information. Conduct a tank spray compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, followed by flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank.

Inosco and Inosco tank-mix combinations have been evaluated for phytotoxicity on a large variety of crops under normal greenhouse and nursery conditions. However, testing all plant species in all mixtures and combinations is not feasible. Test a small portion of the area to be treated for phytotoxicity, before treating the entire area.

When tank mixing Inosco with another registered plant protection product, follow the more restrictive labeling requirements of any tank mix partner. Do not tank mix with products whose label prohibits tank mixing.

Important:

Do not apply on plants when they are under water stress, or during severe weather conditions such as high or very low temperatures. Begin spraying when environmental conditions are favorable for the development of disease or as indicated below.

Application Instructions:

Apply spray solution directly to foliage and drench solutions to the growing media surface, unless directed otherwise in Table 1. Make foliar applications in sufficient water to ensure thorough plant coverage.

Unless specified, there is no limit to the number of applications. When applying Inosco, be sure the spray solution has a **pH between 5.5 and 7.8** to limit the possibility of phytotoxicity to the crop, and that it is applied in sufficient water (> **30 gal/acre** for foliar applications to small fruits and vegetables).

Use the higher listed application rate when conditions favor disease development and/or with increased disease pressure.

Use Restrictions

Do not apply higher than the listed application rates.

Do not apply Inosco with fertilizers in the spray solution.

Table 1. Use Rates.

CROP	DISEASE	RATE/100	REMARKS
	D (1)	gal.	D
Brassica vegetables, such as, broccoli, cabbage, cauliflower, kale,	Pythium spp.	0.75 to 1.5 pints	Drench*
mustard greens	Downy mildew	4 pints	Make preventative foliar applications on a 7 day interval.
Bulb vegetables, such as, garlic, onion, shallot	Pythium spp.	0.75 to 1.5 pints	Drench*
	Downy mildew	4 pints	Make preventative foliar applications on a 7 day interval.
Citrus seedlings	Phytophthora root rot	2 gallons	Drench using 0.5 pint per square foot of bench every 2 months.
Note: Do not apply before or after a severe pruning.			Do not tank-mix with other products.
Conifers	Pythium spp. Phytophthora spp.	4 pints	Pre-plant dip. Thoroughly wet the root mass and plant.
	-11		Apply foliar applications up to run-off every 4 to 6 weeks.
Cucurbits, such as cucumber, melon,	Pythium spp.		Drench*
squash, pumpkin	Downy mildew		Make preventative foliar applications on a 7day interval, in combination with a protectant fungicide, such as Micora Fungicide or Mancozeb.
Fruiting vegetables, such as, eggplant, tomato, pepper	Pythium spp.	0.75 to 1.5 pints	Drench*
	Late blight Bacterial blight - Xanthomonas	4 pints	Make preventative foliar applications on a 7 day interval.
	spp.		Suppression of bacterial diseases may be improved when Inosco is applied in a tank-mix with Kocide® 3000 and Mancozeb.

Herbs and Spices	Pythium spp. Phytophthora spp.	0.75 to 1.5 pints	Drench* Apply as needed on a 7 to 21 day interval.
	Downy mildew	4 pints	Make preventative foliar applications on a 7 day interval. Due to the numerous kinds of herbs and spices, it is recommended that a phytotoxicity trial be initiated on a limited area of the crop prior to treating the entire crop.
Leafy vegetables, such as, lettuce, spinach	Pythium spp.	0.75 to 1.5 pints	Drench*
	Downy mildew	4 pints	Make a preventative foliar application at the 2-leaf stage. Repeat applications on a 7 day interval. During periods of high disease pressure, you may apply Inosco in a tank-mix with Micora Fungicide for improved disease control.
Strawberry	Pythium spp. Phytophthora leather rot, or Red steele	4 pints	Pre-plant dip – thoroughly wet the root mass. Make foliar applications on a 14 to 28 day schedule.

Ornamentals Breeding Crops Bulb, corm and Tuber Crops Conifers Christmas Trees Cut Flowers Evergreens, including Conifers Flowering Plants Foliage plants Ground Covers Perennial Plants Pot and Bedding Plants Shrubs Trees**, including non-bearing fruit and nut trees. Vines**, non- bearing.	Pythium spp. Phytophthora spp.	0.75 to 1.5 pints	Drench*
	Pythium spp. Phytophthora spp.	2 to 4 pints	Make preventative foliar applications on a 7 day interval. Apply no greater than 400 gallons of spray solution per acre. During periods of high disease pressure, Inosco may be applied in a tank-mix with Heritage Fungicide or Micora Fungicide for improved disease control.
	Downy mildew	4 pints	Make preventative applications on a 7 day interval. Apply no greater than 400 gallons of spray solution per acre. During periods of high disease pressure, you may apply Inosco in a tank-mix with Heritage Fungicide, Micora Fungicide or Subdue MAXX Fungicide for improved disease control.
	Bacterial blight - Xanthomonas spp.	4 pints	Preventative foliar applications on a 7 day schedule will provide disease suppression.

Preharvest interval (PHI) for Inosco for all listed fruit and vegetable crops is 0 days.

*Drench applications:

- Apply 1 to 2 pints of drench solution per square foot of surface or growing area unless directed otherwise.
- For small seedlings growing in flats and trays, apply 0.75 pints of drench solution per square foot of surface or growing area.

^{**}Non-bearing fruit and nut trees and vines are plants that will not bear edible fruit or nuts for one year after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a dry, cool place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Container sizes of 5 gallons or less: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Container sizes of greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Read and follow the Directions for Use of this product. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Inosco[™], Heritage®, Micora®, Subdue Maxx®, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Kocide® trademark of E.I. du Pont de Nemours and Company

©20XX Syngenta

 $Inosco\ NEW\ PRODUCT\ JUL2016\ CLEAN\ -rk-7/6/16\\ 000100-0XXXX.20160708.INOSCO.NEW.JUL2016.CL.pdf$