



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
**WASHINGTON, DC 20460**

**OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION**

February 16, 2023

Rachel Hardie  
Regulatory Product Manager  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419

**Subject:** Registration Amendment – Amended Terms and Conditions, Amended Labeling  
Including Collateral Labeling  
Product Name: A21472 PLUS VAPORGRIP TECHNOLOGY (ABN TAVIUM  
PLUS VAPORGRIP TECHNOLOGY)  
EPA Registration Number: 100-1623  
Application Date: 12/15/2022  
Last Resubmission: 02/13/2023  
Decision Number: 589649

Dear Rachel Hardie:

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. The labeling associated with this approval supersedes the previously approved labeling. This amendment does not affect any terms and conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them. This includes the stipulation that this registration remains time-limited and will automatically expire on December 20, 2025, unless amended by EPA.

Syngenta Crop Protection, LLC (Syngenta) submitted a label amendment pursuant to 40 C.F.R. § 152.44 to change the directions for use on this product to further restrict use of this product when used in Iowa, Illinois, Indiana, and South Dakota. Subject to the additional terms and conditions of registration described in this letter, EPA approves the labeling proposed by Syngenta and has determined that the amendment meets the standard of registration. The amendment approved through this letter includes additional, state-specific application date restrictions intended to further reduce off-target movement of the active ingredient dicamba. The states incorporated in this amendment were proposed by the registrant because they accounted for a significant percentage of total off-target movement complaints reported to state lead agencies in the last 3 years and these states did not object in writing to these labeling changes being made.

The ecological risk assessment (available on [www.regulations.gov](http://www.regulations.gov) in docket EPA-HQ-OPP-2020-0492) relied upon in the registration of the product on 10/27/2020 supports the use of measures to address off-target movement, such as application cut-off dates, to avoid applications

of dicamba on days with temperatures favoring off-target movement. As discussed in that risk assessment, ambient temperature has been demonstrated to be directly related to the volatility of dicamba, with higher temperatures leading to increased volatility. Restricting the application to a time when temperature is reduced both on the day of application and in the days following application is likely to reduce the potential for volatilization of dicamba. The registrants of three dicamba-based products registered for over-the-top use, including Syngenta, proposed a June 12<sup>th</sup> cut-off date for Iowa, Illinois, and Indiana. The registrants also proposed a June 20<sup>th</sup> cut-off date for South Dakota, as recommended by that state. A registrant-provided rationale for the June 12<sup>th</sup> date was based on a reduction in off-target movement as a result of decreased temperatures, reduced plant height earlier in the season, the opportunity for applications to be spread more evenly through the season, and indications of improvement in Minnesota during the 2022 season using the June 12<sup>th</sup> cut-off date. EPA views these dates as directionally correct to reduce temperatures at which applications are performed.

Because this amendment approves application cut-off dates earlier than the cut-off dates considered in the Agency's ecological risk assessment developed for the registration of the product on 10/27/2020, this amendment does not allow for exposures beyond what was considered in the Agency's previous ecological risk assessment.

If the terms outlined below are not complied with, the registration may be subject to cancellation in accordance with FIFRA section 6. Syngenta must comply with all the following terms and conditions:

1. To ensure that products are not misbranded or misused, and appropriate labeling is in the possession of users for the 2023 growing season, the following terms and conditions are required.
  - a. For all existing product in the possession of Syngenta or released for shipment and are in the channels of trade prior to date of this letter, Syngenta must make the amended labeling available as described below.
    - i. By 2/22/2023, Syngenta's website where existing collateral labeling is held must be updated to reflect the changes within this amendment where the user will be able to access all collateral labeling for this registration in a downloadable format.
    - ii. By 2/22/2023, copies of the amended labeling must be announced and posted in a prominent location on all Syngenta websites that are part of this product's labeling and be retained until the expiration date of this registration unless superseded by subsequent approved labeling.

- iii. By 2/22/2023, a non-closeable banner containing the following must be prominently displayed at the top of all Syngenta websites that are part of this product's labeling. This banner must be posted until the expiration date of this registration unless superseded by subsequent approved labeling.
  1. "ATTENTION: YOU MUST HAVE ALL LABELING APPLICABLE TO YOUR LOCATION IN YOUR POSSESSION FOR A21472 PLUS VAPORGRIP TECHNOLOGY (ABN TAVIUM PLUS VAPORGRIP TECHNOLOGY) TO BE LAWFULLY APPLIED AFTER March 15, 2022."
  2. A copy or link to a copy of the amended labeling with approval date unless prominent labeling links are visible upon reading banner.
  3. "EPA has approved amended labeling for this product which must be followed."
  4. "It is a violation of FIFRA Section 12 to use a registered pesticide in a manner inconsistent with its labeling."
- b. Syngenta must communicate the availability of and requirement for users to follow the amended labeling in the annual A21472 PLUS VAPORGRIP TECHNOLOGY (ABN TAVIUM PLUS VAPORGRIP TECHNOLOGY) education and training program, including in written educational communications. This information must be incorporated into the training by 2/22/2023. The training course must provide users with the amended labeling, if applicable in the state covered by the training course and must discuss the requirement to visit the website listed on the product label. Materials must emphasize that:
  - i. Users must obtain a copy of the amended labeling, if applicable, which must be in possession of the user at the time of application.
  - ii. After March 15, 2022, users of this product must follow the amended labeling they receive during training, if applicable.
- c. Syngenta must take the following actions to contact all users who have already completed the 2023 training in states where there are EPA-approved state-specific restrictions implemented through the amended labeling to communicate the amended labeling. This will be done by sending all users who have already completed the 2023 training either (1) an email requesting a response via certification statement acknowledging that they have read and understood the new requirements or (2) a physical letter containing a reply postcard acknowledging receipt stating they have read and understood the new labeling requirements. If Syngenta does not receive the email response or reply postcard within two weeks, Syngenta must contact that user two additional times. The second contact attempt must be made by repeating the initial method of contact. Next, if no reply is received one week after the second contact attempt, Syngenta must follow-up another time with an alternative

reasonable means of contact to receive documentation from the user acknowledging that they have read and understood the new labeling requirements. If a valid phone number is used as the method for making the third contact attempt, Syngenta must solicit the user during that phone call to either mail or email them an acknowledgement that they have read and understand the new labeling requirements. Records of these contact attempts and replies must be kept by Syngenta and shared upon request.

- d. Syngenta must send a report to EPA by six weeks after date of this letter detailing its efforts to accomplish all of the above. A copy or a link to the copy of the training materials must be included with this report. Syngenta must maintain copies of any correspondence it sends to all sellers, distributors, and purchasers and make them available to EPA upon request.
2. Syngenta must communicate the availability and requirement to adhere to the amended labeling.
    - a. Syngenta must add the amended labeling to its training and educational materials for the relevant states and have website links to copies of the current approved labeling.
    - b. The educational and training materials supporting this labeling change must be disseminated to state pesticide authorities and state agricultural extension services in the relevant states to support those authorities in assisting users in their local area.

References to the company's website on the label becomes labeling under FIFRA and therefore the website is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA Section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to EPA's attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of the container and collateral labeling is enclosed. This labeling supersedes all previously accepted labeling. Syngenta must submit one printed copy of the final amended labeling for the record prior to adding the amended labeling to its website as required under 1.a.ii. If Syngenta fails to comply with these terms, Syngenta has agreed in prior written acceptance of these terms that EPA may cancel the registration under an expedited process under FIFRA 6(e).

Page 5 of 5  
EPA Reg. No. 100-1623  
Decision No. 589649

You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved container labeling for 18 months from the date of this letter, noting that based on the above conditions the collateral labeling will be posted on the website and go into effect immediately. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions, please contact Lindsay Roe by phone at (202)566-2878, or via email at [roe.lindsay@epa.gov](mailto:roe.lindsay@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Lindsay M.", written in a cursive style.

Lindsay Roe, Chief  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

[Master]

**RESTRICTED USE PESTICIDE**

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

**This EPA registration expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

DICAMBA	GROUP	<b>4</b>	HERBICIDE
S-METOLACHLOR	GROUP	<b>15</b>	HERBICIDE

**A21472 Plus VaporGrip® Technology must only be used for the uses specified on this label and only in the following states, subject to county restriction as noted: Alabama, Arizona, Arkansas, Colorado, Delaware, Florida (excluding Palm Beach County), Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, New York (excluding Nassau and Suffolk Counties), North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee (excluding Wilson County), Texas (excluding use on cotton in Gaines County), Virginia, West Virginia, Wisconsin.**


**Users must comply with additional requirements including state restrictions found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). The user must check [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com) no more than 7 days before application of this product for additional labeling.**

**Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

Primary Brand Name:  
**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:  
**Tavium® Plus VaporGrip® Technology**

**Herbicide**



Foliar systemic broadleaf herbicide with residual grass and certain broadleaf weed control for dicamba-tolerant cotton and dicamba-tolerant soybeans

Active Ingredients:

Diglycolamine salt of dicamba*:	17.7%
S-metolachlor**:	24.0%
<hr/>	
Other Ingredients:	58.3%
Total:	100.0%

\*CAS No. 104040-79-1

\*\*CAS No. 87392-12-9

A21472 Plus VaporGrip Technology is a capsule suspension (CS) formulation containing 1.12 pounds of dicamba acid equivalent (ae) and 2.26 pounds of S-metolachlor per U.S. gallon.

**KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

EPA Est.

---

Net Contents

## TABLE OF CONTENTS

### 1.0 FIRST AID

### 2.0 PRECAUTIONARY STATEMENTS

#### 2.1 Hazards to Humans and Domestic Animals

#### 2.2 Personal Protective Equipment (PPE)

2.2.1 User Safety Requirements

2.2.2 Engineering Controls

2.2.3 User Safety Recommendations

#### 2.3 Environmental Hazards

2.3.1 Groundwater Advisory

2.3.2 Surface Water Advisory

2.3.3 NON-TARGET ORGANISM ADVISORY

2.3.4 Reporting Ecological Incidents

2.3.5 Mixing/Loading Instructions

#### 2.4 Physical or Chemical Hazards

## DIRECTIONS FOR USE

### 3.0 PRODUCT INFORMATION

#### 3.1 Weed Resistance Management Practices

3.1.1 Principles of Herbicide Resistant Weed Management

### 4.0 APPLICATION DIRECTIONS

#### 4.1 Training

#### 4.2 Record Keeping

#### 4.3 Methods of Application

#### 4.4 Application Equipment

#### 4.5 Application Volume and Spray Coverage

#### 4.6 Equipment Ground Speed

#### 4.7 Optional Use of Drift Reduction Technology

4.7.1 Hooded/Shielded Broadcast Sprayer

4.7.2 Ground Application (Hooded In-Row and Directed Layby)

#### 4.8 Tank Mix Partners and Compatibility Testing

4.8.1 Tank-Mix Restrictions

4.8.2 Tank-Mix Precautions

#### 4.9 Sprayer Cleanout

### 5.0 REPLANT AND ROTATIONAL CROP

#### 5.1 Rotational Crop Restrictions

### 6.0 RESTRICTIONS AND PRECAUTIONS

#### 6.1 Use Restrictions

#### 6.2 Use Precautions

#### 6.3 Mandatory Spray Drift Management

6.3.1 Importance of Droplet Size

6.3.2 Controlling Droplet Size – Ground Boom

6.3.3 BOOM HEIGHT – Ground Boom



6.3.4 BOOMLESS GROUND APPLICATION

6.3.5 SHIELDED SPRAYERS

6.3.6 Wind

6.3.7 Temperature and Humidity

6.3.8 Temperature Inversions

6.3.9 Spray Buffers

6.3.10 Sensitive Crops

## **7.0 WEEDS CONTROLLED BY A21472 PLUS VAPORGRIP TECHNOLOGY**

7.1 **Weeds Controlled by A21472 Plus VaporGrip Technology Applied Prior to Weed Emergence**

7.2 **Weeds Controlled by A21472 Plus VaporGrip Technology Applied Postemergence to Weeds**

## **8.0 CROP USE DIRECTIONS**

8.1 **Dicamba-Tolerant Cotton**

8.1.1 Preplant, At-Planting, Preemergence or Postemergence (In-Crop) Application

8.2 **Dicamba-Tolerant Soybean**

8.2.1 Preplant, At-Planting, Preemergence or Postemergence (In-Crop) Application

## **9.0 STORAGE AND DISPOSAL**

## **10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

## **11.0 [APPENDIX]**

11.1 **[A21472 Plus VaporGrip Technology Use Summary Table]**

11.2 **[State Additional Labeling]**

11.3 **[Additional Iowa-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]**

11.4 **[Additional Minnesota-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]**

11.5 **[Additional Illinois-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]**

11.6 **[Additional Indiana-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]**

11.7 **[Additional South Dakota-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]**

## 1.0 FIRST AID

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p><b>HOTLINE NUMBER</b></p> <p>For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-888-8372</b></p>	

## 2.0 PRECAUTIONARY STATEMENTS

### 2.1 Hazards to Humans and Domestic Animals

#### **CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### 2.2 Personal Protective Equipment (PPE)

**All mixers, loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

#### 2.2.1 User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## 2.2.2 Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

## 2.2.3 User Safety Recommendations

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## 2.3 Environmental Hazards

**DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash water or rinsate. Apply this product only as directed on the label.

### 2.3.1 Groundwater Advisory

Dicamba and S-metolachlor are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Groundwater contamination may occur in areas where soils are permeable or coarse and groundwater is near the surface. **DO NOT** apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for groundwater contamination. **DO NOT** apply to soils classified as sand with less than 3% organic matter and where groundwater depth is shallow.

### 2.3.2 Surface Water Advisory

**DO NOT** apply if soil is saturated with water or when rainfall that may exceed soil field capacity is forecasted to occur within 48 hours.

Under some conditions, dicamba has the potential for runoff several days and s-metolachlor for several months after application. Poorly draining, wet, or erodible soils with readily visible slopes toward adjacent sensitive areas are more prone to produce runoff. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Soil Conservation Service for recommendations in your use area.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip between areas to which this

product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of dicamba and S-metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

### 2.3.3 NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

### 2.3.4 Reporting Ecological Incidents

To report ecological incidents, including mortality, injury, or harm to plants and animals, call 1-800-888-8372.

### 2.3.5 Mixing/Loading Instructions

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or anti-siphoning devices must be used on all mixing equipment.

- This product must not be mixed or loaded within 50 ft of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs, wells, including abandoned wells, drainage wells, and sink holes.
- Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling, or application equipment or containers within 50 ft of any well are prohibited, unless conducted on an impervious pad that meets the following specifications. Containment capacities described below must be maintained at all times.
  - The pad must be constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad.
  - The pad must be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rain water that may fall on the pad.
  - Surface water must not be allowed to either flow over or from the pad, which means the pad must be self-contained and sloped.
  - An unroofed pad must contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad.
  - A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad, must have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad.

## 2.4 Physical or Chemical Hazards

**DO NOT** use or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### Endangered Species Protection Requirements

Prior to making an application of this product on dicamba-tolerant cotton or dicamba-tolerant soybeans, an applicator must visit <http://www.epa.gov/espp/> to determine if there are any additional restrictions on A21472 Plus VaporGrip Technology use within the area to be sprayed.

It is a Federal offense to use any pesticide in a manner that results in an unauthorized “take” (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act, Section 9. When using this product, you must follow the measures contained in the Endangered Species Bulletin for the area in which you are applying this product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product

**Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

Use A21472 Plus VaporGrip Technology only in accordance with specifications on this label, or in separately EPA-approved labeling instructions for this product.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective

equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

## APPLICATION REQUIREMENTS OVERVIEW

Read and follow all applicable restrictions, precautions, and directions on the container label and booklet and at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). For product questions or inquiries and/or to report any nonperformance of this product against any particular weed species, call 1-866-Syngent(a) (866-796-4368).

Product Use Requirements	Label Section
<p><b><u>Training Requirements Prior to Use:</u></b></p> <ul style="list-style-type: none"> <li>• Only certified applicators can apply A21472 Plus VaporGrip Technology.</li> <li>• Prior to applying this product, applicator(s) must complete dicamba-specific training.</li> <li>• This product is NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.</li> </ul>	4.1
<p><b><u>Record Keeping Requirements:</u></b></p> <ul style="list-style-type: none"> <li>• The certified applicator must keep application records specified in the Record Keeping requirements section for a period of two years.</li> <li>• All tank mixing partners must be recorded.</li> <li>• Application records must be generated as soon as practical but no later than 72 hours after application, unless state restrictions require records to be generated sooner.</li> </ul>	4.2
<p><b><u>General Application Requirements:</u></b></p> <ul style="list-style-type: none"> <li>• Only apply A21472 Plus VaporGrip Technology using approved nozzles designed to produce extremely coarse to ultra-coarse droplets listed at <a href="http://www.TaviumApplicationRequirements.com">www.TaviumApplicationRequirements.com</a>.</li> <li>• <b>DO NOT</b> exceed a boom height of 24 inches above target pest or crop canopy when applying this product.</li> <li>• A21472 Plus VaporGrip Technology must be applied in a minimum of 15 gallons of spray solution per acre.</li> <li>• <b>DO NOT</b> exceed a ground speed of 15 miles per hour.</li> <li>• All applications of A21472 Plus VaporGrip Technology must include an approved volatility reduction agent (VRA) / buffering agent (pH modifier) listed at <a href="http://www.TaviumApplicationRequirements.com">www.TaviumApplicationRequirements.com</a>. An approved drift reduction agent (DRA) must also be included in the spray solution, unless otherwise indicated on <a href="http://www.TaviumApplicationRequirements.com">www.TaviumApplicationRequirements.com</a>. Refer to the website for a list of approved DRAs and volatility reduction agents (VRAs) / buffering agents (pH modifiers).</li> </ul>	4.4-4.9

<ul style="list-style-type: none"> <li>• A21472 Plus VaporGrip Technology must only be tank-mixed with approved products listed at <a href="http://www.TaviumApplicationRequirements.com">www.TaviumApplicationRequirements.com</a>. <b>DO NOT</b> tank mix with products not included on this approved products list.</li> <li>• Spray system equipment cleanout: User must ensure entire sprayer system is properly cleaned before and after each application.</li> </ul>	
<p><b><u>Environmental Condition Requirements:</u></b></p> <ul style="list-style-type: none"> <li>• ONLY apply A21472 Plus VaporGrip Technology when wind speed is 3 to 10 MPH.</li> <li>• <b>DO NOT</b> apply A21472 Plus VaporGrip Technology during a temperature inversion.</li> <li>• <b>Only apply A21472 Plus VaporGrip Technology using a broadcast open-boom from one hour after sunrise to two hours before sunset.</b></li> <li>• When making applications in low relative humidity or high temperatures, set up equipment to deliver larger droplets to compensate for evaporation.</li> <li>• <b>DO NOT</b> apply this product if rainfall could exceed soil field capacity and result in soil runoff is expected in the next 48 hours.</li> </ul>	6.0
<p><b><u>Application Requirements for Dicamba Tolerant Cotton:</u></b></p> <ul style="list-style-type: none"> <li>• A21472 Plus VaporGrip Technology may be applied preplant or preemergence at a maximum rate of 3.53 pt/A (0.5 lb ae/a dicamba) to dicamba-tolerant cotton</li> <li>• A21472 Plus VaporGrip Technology may be applied postemergence at a maximum rate of 3.53 pt/A (0.5 lb ae/a dicamba) in dicamba-tolerant cotton through 6-leaf or up to and including July 30, whichever comes first.</li> </ul>	8.1
<p><b><u>Application Requirements for Dicamba Tolerant Soybean:</u></b></p> <ul style="list-style-type: none"> <li>• A21472 Plus VaporGrip Technology may be applied preplant or preemergence at a maximum rate of 3.53 pt/A (0.5 lb ae/a dicamba) to dicamba-tolerant soybean.</li> <li>• A21472 Plus VaporGrip Technology may be applied postemergence at a maximum rate of 3.53 pt/A (0.5 lb ae/a dicamba) in dicamba-tolerant soybean through V4 or up to and including June 30, whichever comes first.</li> </ul>	8.2
<p><b><u>Buffer Requirements:</u></b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply if sensitive plants and/or crops are planted in adjacent downwind fields or areas when using broadcast open-boom equipment.</li> <li>• Sensitive crop definitions can be found in section 6.3.8.</li> <li>• The user must ensure there is a <b>240 foot downwind</b> buffer when applying with broadcast open-boom equipment.</li> <li>• Prior to making an application of this product on dicamba-tolerant cotton or dicamba-tolerant soybeans, an applicator must visit <a href="http://www.epa.gov/espp/">http://www.epa.gov/espp/</a> to determine if there are any additional restrictions on A21472 Plus VaporGrip Technology use within the area to be sprayed.</li> </ul>	2.3 6.0

### 3.0 PRODUCT INFORMATION

A21472 Plus VaporGrip Technology is a foliar systemic broadleaf herbicide with residual control of grass and certain broadleaf weeds in:

- dicamba-tolerant cotton (preplant, at-planting, preemergence, postemergence (In-crop) application)
- dicamba-tolerant soybeans (preplant, at-planting, preemergence, postemergence (In-crop) application)

This product needs a minimum of ½ inch of either rainfall or irrigation following application to activate residual weed control. If rainfall or irrigation is not received within 10 days after application, residual weed control may be reduced. Under these conditions, cultivate or use other weed control measures if weeds develop.

**DO NOT** apply this product if rainfall could exceed soil field capacity and result in soil runoff is expected in the next 48 hours.

Rainfall or irrigation occurring within 4 hours after postemergence application may reduce effectiveness.

### 3.1 Weed Resistance Management Practices

DICAMBA	GROUP	4	HERBICIDE
S-METOLACHLOR	GROUP	15	HERBICIDE

For resistance management, please note that A21472 Plus VaporGrip Technology contains both a Group 4/(dicamba) and a Group 15/(S-metolachlor) herbicide. Any weed population may contain plants naturally resistant to Group 4 and/or Group 15 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Contact your local Syngenta representative, retailer, crop advisor or extension agent for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes. **DO NOT** assume that each listed weed is being controlled by multiple modes of action. Premixes are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product. If resistant biotypes have been reported, use the full labeled rate of this product, apply at the labeled timing, and tank-mix with an additional different mode of action product so there are multiple effective modes of application for each suspected resistant weed. To delay herbicide resistance, take one or more of the following steps:

#### 3.1.1 Principles of Herbicide Resistant Weed Management

Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural ( e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

#### Scout and know your field

- Know weed species present in the field to be treated through scouting and field history. An understanding of weed biology is useful in designing a resistance management strategy. Ensure the weed management program will control all weeds present.
- Fields should be scouted prior to application to determine species present and growth stage. Always apply this herbicide at the full labeled rate and correct timing for the weeds present in the field.



### **Utilize non-herbicidal practices to add diversity**

- Use diversified management tactics such as cover crops, mechanical weed control, harvest weed seed control, and crop rotation as appropriate. Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

### **Use good agronomic practices, start clean and stay clean**

- Use good agronomic practices that enhance crop competitiveness.
- Plant into weed-free fields utilizing tillage or an effective burndown herbicide for control of emerged weeds.
- Sanitize farm equipment to avoid spreading seed or vegetative propagules prior to leaving fields.

### **Difficult to control weeds**

- Fields with difficult to control weeds should be planted in rotation with crops that allow the use of herbicides with an alternative mode of action or different management practices.
- Difficult to control weeds may require sequential applications, such as a broad spectrum preemergence herbicide followed by one or more postemergence herbicide applications. Utilize herbicides containing different modes of action effective on the target weeds in sequential applications.

### **DO NOT overuse the technology**

- **DO NOT** use more than two applications of this or any other herbicide with the same mode of action in a single growing season unless mixed with an herbicide with a different mode of action which provides overlapping spectrum for the difficult to control weeds.

### **Scout and inspect fields following application**

- Prevent an influx of weeds into the field by controlling weeds in field borders.
- Scout fields after application to verify that the treatment was effective.
- Indicators of possible herbicide resistance include:
  - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - A spreading patch of non-controlled plants of a particular weed species; and
  - Surviving plants mixed with controlled individuals of the same species.

For further information or to report suspected resistance your Syngenta retailer, Syngenta representative, or call 1-866-Syngent(a) (866-796-4368).

- If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- **Prevent weed escapes before, during, and after harvest**
- **DO NOT** allow weed escapes to produce seed or vegetative structures such as tubers or stolons which contribute to spread and survival.

## 4.0 APPLICATION DIRECTIONS

### 4.1 Training

Prior to applying this product, all applicators must complete dicamba-specific training on an annual basis. This product is NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers. If training is available and required by the state where the applicator intends to apply this product, the applicator must complete that training before applying this product. If your state does not require dicamba-specific training, then the applicator must complete dicamba-specific training provided by one of the following sources: a) a registrant of a dicamba product approved for in-crop use with dicamba-tolerant crops, or b) a state or state-authorized provider.

### 4.2 Record Keeping

Record keeping is required for each application of this product. **The certified applicator must keep the following records for a period of two years;** records must be generated as soon as practical but no later than 72 hours after application, unless state restrictions require records to be generated sooner. Records must be made available to State Pesticide Control Official(s), USDA, and EPA upon request. An example form summarizing record keeping requirements can be found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com).

1. All Items required by 7 CFR Part 110 (RECORDKEEPING ON RESTRICTED USE PESTICIDES BY CERTIFIED APPLICATORS) including:
  - a. The brand or product name
  - b. The EPA registration number
  - c. The total amount applied
  - d. The month, day, and year of application
  - e. The location of the application
  - f. The crop, commodity, stored product, or site of application
  - g. The size of treated area
  - h. The name of the certified applicator
  - i. The certification number of the certified applicator
2. Training: Date and provider of required training completed and proof of completion.

3. Receipts of Purchase: Receipts or copies for the purchase of this product, and for the purchase of the required volatility reduction agent (VRA) / buffering agent (pH modifier) and drift reduction agent (DRA).
4. Product Label: A copy of the product label and any labeling that supplements the product label.
5. Buffer Requirement: Record of the buffer distance and any non-sensitive areas included in buffer distance. Non-sensitive areas refers to the exclusion areas that can be counted as part of the buffer, such as paved areas.
6. Sensitive Plants and/or Crops Awareness: Record that a sensitive crop registry was consulted and survey adjacent fields documenting the crops/areas surrounding the field prior to application. Records at a minimum must include the name of the sensitive crop registry and the date it was consulted and documentation of adjacent crops/areas and the date the survey was conducted.
7. Start and Finish Times of Each Application: Record of the time at which the application started and the time when the application finished.
8. Application Timing: Record of the type of application (for example: preemergence, postemergence).
9. Air Temperature: Record of the air temperature in degrees Fahrenheit at the start and completion of each application.
10. Wind Speed and Direction: Record of the wind speed and direction (the direction from which the wind is blowing) at boom height at the start and completion of each application of this product.
11. Nozzle and Pressure: Record of the spray nozzle manufacturer/brand, type, orifice size, and operating pressure used during each application of this product.
- 12. Tank Mix Products: Record of the brand names, EPA registration numbers (if available), and use rates for all products including volatility reduction agent (VRA) / buffering agent (pH modifier) and DRA that were tank mixed with this product for each application.**
13. Spray System Cleanout: At a minimum, records must include the confirmation that the spray system was clean before using this product and that the post-application cleanout was completed in accordance with the Sprayer Cleanout Section 4.9.

### 4.3 Methods of Application

Applications with A21472 Plus VaporGrip Technology alone or in tank mixtures are permitted with ground equipment only. This product may be applied using broadcast or hooded broadcast applications for postemergence weed control as well as residual control of susceptible weeds. Preplant, at-planting, preemergence and postemergence (In-crop) applications are allowed. All labeled application restrictions must be adhered to when using these application methods.

### 4.4 Application Equipment

- Configure spray equipment to provide accurate and uniform coverage of the target area and minimize potential for spray drift.
- Only use sprayers that provide accurate and uniform application with nozzles designed to

produce **extremely coarse to ultra-coarse** droplets in order to minimize drift (**Section 6.3.1**) and provide uniform coverage.

- To ensure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult spray equipment manufacturers and/or state recommendations.
- Avoid using screens and strainers finer than 50-mesh.
- All ground application equipment must be properly maintained.
- Equipment must be cleaned out after use following Spray System Cleanout procedures in Section 4.9.

## 4.5 Application Volume and Spray Coverage

- Apply alone or in tank mixtures **in a minimum of 15 gal/A of spray solution**.
- Good spray coverage of emerged weeds is essential for optimum control.
- When weed vegetation is dense, increase spray volume and pressures to ensure coverage of the target weeds.
- Spray boom and nozzle heights must be adjusted to provide coverage of target weeds but not more than 24 inches above the target.

## 4.6 Equipment Ground Speed

**DO NOT exceed a ground speed of 15 miles per hour.** Select a ground speed that will deliver the desired spray volume while maintaining the desired spray pressure, but slower speeds generally result in better spray coverage and deposition on the target area. Provided the applicator can maintain the required nozzle pressure, it is recommended that tractor speed is reduced to 5 miles per hour at field edges.

## 4.7 Optional Use of Drift Reduction Technology

This product may be optionally applied using a hooded/shielded broadcast sprayer or other types of drift reduction technology (DRT) for postemergence weed control as well as residual control of susceptible weeds. The applicator must use an approved nozzle within a specified pressure range as found at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). Use of drift reduction technology (DRT) in combination with approved nozzles is recommended to further reduce drift potential.

### 4.7.1 Hooded/Shielded Broadcast Sprayer

For hooded/shielded sprayers, all application nozzles must be contained within the enclosed area. Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product. Applicators must ensure the sprayer system is off or controlled during turns to further prevent spray drift. Refer to the hooded/shielded sprayer manufacturer use specifications prior to use.

*Requirements for Reduced Use Restrictions with Optional Hooded/Shielded Sprayer:*

**ONLY qualified** hooded/shielded sprayers listed on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com) are eligible for reduced use restrictions when applying this product. When using a qualified hooded/shielded sprayer, **the applicator must**

**always maintain a minimum 110-foot downwind buffer** between the last treated row and the nearest downwind field edge. Consult Endangered Species Protection Bulletins for ESA counties and restrictions. While this product may be applied with other (non-qualified) hooded/shielded sprayers, no reduction in use restrictions is associated with their use.

#### 4.7.2 Ground Application (Hooded In-Row and Directed Layby)

Using a hooded sprayer or other drift reduction technology in combination with approved nozzles may further reduce drift potential. When applying A21472 Plus VaporGrip Technology by hooded in-row or layby sprayers, determine the amount of herbicide and water volume needed using the following formula:

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} * \text{broadcast rate per acre} = \text{rate per treated acre}$$

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} * \text{broadcast volume per acre} = \text{spray volume per treated acre}$$

### 4.8 Tank Mix Partners and Compatibility Testing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

- All applications of this product require the inclusion of an approved volatility reduction agent (VRA) / buffering agent (pH modifier) listed at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com).
- All applications of this product require the inclusion of a drift reduction agent (DRA) in the tank mix, unless otherwise indicated on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com).
- A21472 Plus VaporGrip Technology may only be tank-mixed with products that have been tested and found not to adversely affect the offsite movement potential of A21472 Plus VaporGrip Technology. The applicator must check the website found at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com) no more than 7 days before applying A21472 Plus VaporGrip Technology.
- For preplant, at-planting and preemergence applications, the following additives may be used. A complete list of approved products can be found at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com).
  - Nonionic Surfactant (NIS) - Use NIS containing at least 80% active ingredient at 0.25% v/v (1 qt/100 gal) of the finished spray volume.
  - Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO) – Use a nonphytotoxic COC or MSO containing 15–20% approved emulsifier at 0.5–1.0% v/v (2-4 qt/100 gal) of the finished spray volume.
- For postemergence applications, use of a Nonionic Surfactant (NIS) additive described above is allowed.
  - Use of a Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO) for postemergence applications is not advised due to the potential for crop injury.

### 4.8.1 Tank-Mix Restrictions

- **DO NOT** tank mix any product with A21472 Plus VaporGrip Technology unless:
  - The intended tank-mix product is identified on the list of tested products found at [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com);
  - The intended products are not prohibited on either this label or the label of the tank mix product; and
  - All requirements and restrictions on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com); are followed.
- **DO NOT** apply A21472 Plus VaporGrip Technology with ammonium sulfate (AMS) containing additives, conditioners, or fertilizers.

### 4.8.2 Tank-Mix Precautions

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## 4.9 Sprayer Cleanout

As part of the Restricted Use Product requirements, applicators must document that they have complied with the Sprayer Clean-out section of this label

Severe crop injury may occur if any of this product remains in the spray system equipment following an application and the equipment is subsequently used for application to sensitive crops. After using this product, clean all mixing and spray equipment (including tanks, pumps, lines, filters, screens, and nozzles) with a strong detergent based sprayer cleaner. The rinsate must be disposed in compliance with local, state, and federal guidelines.

Inadvertent contamination can also occur in equipment used for bulk product handling and mixing prior to use in the spray system. Care should be taken to reduce contamination not only in the spray system but in any equipment used to transfer or deliver product. For example, bulk handling and mixing equipment containing this product should be segregated when possible to reduce potential for cross-contamination. Consider using block and check valves to avoid backflow during transfer. Piping should be reviewed to ensure there not potential for product build-up. Dedicated nurse trucks and tender equipment should be used when possible.

To avoid subsequent injury to other crops, thoroughly clean mixing and application equipment immediately after spraying using the triple rinse procedures below:

1. **DO NOT** clean sprayer near desirable vegetation, wells or other water sources.
2. Drain and flush tank walls, boom and all hoses with clean water.
3. Prepare a cleaning solution with a detergent or a commercial sprayer cleaner or ammonia according to the product's use directions.
4. Be sure to wash all internal parts of the tank, including the inside top surface with the cleaning solution. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.

5. Flush hoses, spray lines and nozzles for at least one minute with the cleaning solution.
6. Repeat steps 3-5 for two additional times.
7. Remove nozzles, screens and strainers, and clean separately in the cleaning solution after completing the above procedures.
8. Drain lines, filters and sump.
9. Rinse the complete spraying system with clean water.
10. Clean and wash off the outside of the entire sprayer and boom.
11. Dispose of all rinsate according to local, state and federal regulation.

## 5.0 REPLANT AND ROTATIONAL CROP

### 5.1 Rotational Crop Restrictions

The following crops may be planted at the specified interval following application of A21472 Plus VaporGrip Technology. Exclude counting days from application when the ground is frozen.

Crop	Plant-Back Interval
Dicamba-tolerant cotton Dicamba-tolerant soybeans Corn (field, pop, seed, sweet)*	0 days
Non-dicamba-tolerant soybeans	28 days following a minimum accumulation of 1 inch of rainfall or overhead irrigation
Non-dicamba-tolerant cotton	42 days following a minimum accumulation of 1 inch of rainfall or overhead irrigation
Barley Oats Rye Wheat	4 ½ months
Alfalfa Bean Beet Broccoli Brussels sprouts Cabbage Carrot Cauliflower Celery Garlic Lentil Onion Pea Peanut Pepper Potato Pumpkin Radish Sorghum Sunflower Sugar beet Sweet potato Tomato	6 months
Clover (seeded)	9 months
Buckwheat Rice Tobacco	In the next spring following treatment

All other crops not listed above	12 months
----------------------------------	-----------

\* User precaution for corn plantback: Application of this product to coarse-textured soils (sand, loamy sand, or sandy loam) or any soil with less than 2.5% organic matter under cool, wet conditions may result in transient crop injury.

## 6.0 RESTRICTIONS AND PRECAUTIONS

### 6.1 Use Restrictions

- **Users must comply with additional requirements including state restrictions found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). The user must check [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com) no more than 7 days before application of this product for additional labeling.**
- **DO NOT** sell, use or distribute this product in Nassau and Suffolk Counties in the State of New York.
- **DO NOT** use in nurseries, turf, or landscape plantings.
- **DO NOT** apply this product by air.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply A21472 Plus VaporGrip Technology with ammonium sulfate (AMS) containing additives, conditioners, or fertilizers.
- **DO NOT** apply this product at ground speed greater than 15 miles per hour.
- **DO NOT** apply this product in less than 15 gallons of spray solution per acre.
- **DO NOT** exceed a boom height of 24 inches above target pest or crop canopy when applying this product.
- **DO NOT** apply this product when the wind speeds are less than 3 mph or greater than 10 mph.
- **DO NOT** apply this product until at least one hour after sunrise and no later than two hours before sunset.
- **DO NOT** apply to soils classified as sand with less than 3% organic matter and where groundwater depth is shallow.
- **DO NOT** apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.
- **DO NOT** graze or feed to livestock, or harvest for food, any cover crop planted following an A21472 Plus VaporGrip Technology treated crop.
- **DO NOT** apply to frozen ground.
- **DO NOT** apply to any body of water.
- **DO NOT** contaminate irrigation ditches.
- **DO NOT** apply this product if rainfall that could exceed soil field capacity and result in soil runoff is expected in the next 48 hours.
- **DO NOT** apply to powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, you must ensure that the soil surface is first settled by rainfall or irrigation prior to application.
- **DO NOT** apply to impervious substrates, such as paved or highly compacted surfaces.
- **DO NOT** use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least ½ inch of rainfall has occurred between application and the first irrigation.
- Prior to making an application of this product on dicamba-tolerant cotton or dicamba-



tolerant soybeans, an applicator must visit <http://www.epa.gov/espp/> to determine if there are any additional restrictions on A21472 Plus VaporGrip Technology use within the area to be sprayed.

<b>A21472 Plus VaporGrip Technology must only be used for the uses specified on this label and only in the following states</b>	
Alabama	Nebraska
Arizona	New Jersey
Arkansas	New Mexico
Colorado	New York (excluding Nassau & Suffolk Counties)
Delaware	North Carolina
Florida (excluding Palm Beach County)	North Dakota
Georgia	Ohio
Illinois	Oklahoma
Indiana	Pennsylvania
Iowa	South Carolina
Kansas	South Dakota
Kentucky	Tennessee (excluding Wilson County)
Louisiana	Texas (excluding use on cotton in Gaines County)
Maryland	Virginia
Michigan	West Virginia
Minnesota	Wisconsin
Mississippi	
Missouri	

This product must only be used in the states listed above and is subject to area specific restrictions as required by <http://www.epa.gov/espp/> that must be consulted prior to making an application in dicamba-tolerant cotton or dicamba-tolerant soybeans.

## 6.2 Use Precautions

- A21472 Plus VaporGrip Technology requires actively growing green plant tissue to function fully for postemergence weed control. Application of this product to drought-stressed weeds or weeds with little green foliage (i.e., mowed, cut, or hailed on weeds); weeds covered with dust; weeds damaged by insects or diseases may result in reduced weed control.
- Drift may cause damage to nontarget vegetation.
- Avoid spray overlap, as crop injury may result.

## 6.3 Mandatory Spray Drift Management

### **Ground Boom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 24 inches above the ground or crop canopy.
- Applicators are required to use only nozzles producing extremely coarse to ultra-coarse droplets (ASABE S572).
- **DO NOT** apply during temperature inversions.

- **DO NOT** apply when the wind speed is less than 3 mph or greater than 10 mph.
- **DO NOT** apply when weather conditions may cause drift to nontarget areas.

**Boomless Ground Applications:**

- Applicators are required to use only nozzles producing extremely coarse to ultra-coarse droplets (ASABE S572.3) for all applications.
- **DO NOT** apply when the wind speed is less than 3 mph or greater than 10 mph.
- **DO NOT** apply during temperature inversions.

**SPRAY DRIFT ADVISORIES**

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering these factors when making a decision.

**6.3.1 Importance of Droplet Size**

- An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. Use only nozzles producing **extremely coarse to ultra-coarse** droplets listed on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com).
- While applying larger droplets will reduce drift potential, drift will be greater if applications are made improperly, or under unfavorable environmental conditions.

**6.3.2 Controlling Droplet Size – Ground Boom**

- **Volume** – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume not less than 15 gallons per acre for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size. **DO NOT** exceed the nozzle manufacturer's specified pressures or maximum pressures as listed for specific nozzles on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure. If sprayer is equipped with rate controller hardware, ensure it does not allow pressure increases that exceed the desired range.
- **Spray nozzles** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

**6.3.3 BOOM HEIGHT – Ground Boom**

- For ground equipment, the boom should remain level with the crop and have a minimal bounce. Use manufacturer's recommendation for boom height or 24 inches above the crop height, whichever is smaller. Excessive boom height will increase the drift potential.

**6.3.4 BOOMLESS GROUND APPLICATION**

- Setting nozzles at the lowest effective height will help reduce the potential for spray drift.

### 6.3.5 SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### 6.3.6 Wind

- Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**
- Drift potential is lowest when wind speeds are 3 to 10 mph. **DO NOT** apply this product when the wind speed is less than 3 mph or greater than 10 mph.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### 6.3.7 Temperature and Humidity

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation (for example: increase orifice size and/or increase spray volume as directed on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)).

### 6.3.8 Temperature Inversions

- Drift potential is high during a temperature inversion.
- **DO NOT apply this product during a temperature inversion.**
- **ONLY** apply this product between one hour after sunrise and two hours before sunset.
- Temperature inversions are characterized by increasing temperatures with altitude, and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning.
- The presence of an inversion can be indicated by ground fog; or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- The inversion will typically dissipate with increased winds (above 3 miles per hour) or at sunrise when the surface air begins to warm (generally 3°F from morning low).

### 6.3.9 Spray Buffers

- **DO NOT apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or the crops thereof may be rendered unfit for sale, use or consumption.**
- Apply A21472 Plus VaporGrip Technology only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or

endangered species, nontarget crops) is minimal (i.e., when wind is blowing away from the sensitive areas).

- When applying this product a **240 foot downwind buffer must be maintained** between the last treated row, and the closest downwind edge (in the direction in which the wind is blowing).
- To maintain this required buffer zone, no application swath can be initiated in, or into an area that is within the applicable buffer distance.
- The following areas may be included in the buffer distance calculation when adjacent to field edges:
  - Roads, paved or gravel surfaces, mowed grass areas, and areas of bare ground from recent plowing or grading that are contiguous to the treated field.
  - Planted agricultural fields containing: corn, dicamba-tolerant cotton, dicamba-tolerant soybeans, sorghum, proso millet, small grains and sugarcane and other crops approved for postemergence dicamba use. If the applicator intends to include such crops as dicamba-tolerant cotton and/or dicamba-tolerant soybeans in the buffer distance calculation, the applicator must confirm the crops are in fact dicamba-tolerant and not conventional cotton and/or soybeans.
  - Agricultural fields that have been prepared for planting.
  - Areas covered by the footprint of a building, silo, or other man-made structure with walls and/or roof.
- Applicators are required to ensure that they are aware of the proximity to sensitive areas, to avoid potential adverse effects from off-target movement of A21472 Plus VaporGrip Technology.

### 6.3.10 Sensitive Crops

To protect sensitive crops including cover crops, the following restrictions must be followed.

- The applicator must survey the application site for adjacent nontarget sensitive crops. The applicator must also consult applicable sensitive crop registries to identify any commercial specialty or certified organic crops that may be located near the application site. At a minimum, records must include the name of the sensitive crop registry and the date it was consulted and documentation of adjacent crops/areas and the date the survey was conducted.

**DO NOT APPLY** this product when the wind is blowing toward adjacent non-dicamba-tolerant sensitive crops and/or plants. **Sensitive Crops may be severely injured or killed if they are contacted by this product.**

The applicator must be aware that wind direction may vary during the application. If wind direction shifts such that the wind is blowing toward adjacent sensitive crops, the applicator must STOP the application.

Crops known to be sensitive include but are not limited to:

- non-dicamba-tolerant soybeans
- non-dicamba-tolerant cotton
- EPA Crop Group 6 (peas and beans)
- EPA Crop Group 8 (fruiting vegetables including peppers and tomatoes)
- EPA Crop Group 9 (cucurbit group including cucumbers and melons)
- flowers
- fruit trees
- grapes
- ornamental plantings including broadleaf ornamentals grown in greenhouses and shadehouses
- other broadleaf plants
- peanuts
- potatoes
- sweet potatoes
- sunflower
- tobacco

## 7.0 WEEDS CONTROLLED BY A21472 PLUS VAPORGRIP TECHNOLOGY

### 7.1 Weeds Controlled by A21472 Plus VaporGrip Technology Applied Prior to Weed Emergence

Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus powellii</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Crabgrass, large	<i>Digitaria ischaemum</i>
Crabgrass, smooth	<i>Digitaria sanguinalis</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria pumila</i>
Goosegrass	<i>Eleusine indica</i>
Nightshade, Eastern black	<i>Solanum ptychanthum</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>

Common Name	Scientific Name
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Pusley, Florida	<i>Richardia scabra</i>
Signalgrass, broadleaf	<i>Urochloa platyphylla</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
Witchgrass	<i>Panicum capillare</i>

## 7.2 Weeds Controlled by A21472 Plus VaporGrip Technology Applied Postemergence to Weeds

Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus powellii</i>
Amaranth, spiny	<i>Amaranthus spinosus</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>
Buffalobur	<i>Solanum rostratum</i>
Burcucumber	<i>Sicyos angulatus</i>
Buttercup	<i>Ranunculus</i> spp.
Carpetweed	<i>Mullugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Copperleaf, hophornbeam	<i>Acalypha ostryifolia</i>
Croton, tropic	<i>Croton glandulosus</i>
Cutleaf eveningprimrose	<i>Oenothera laciniata</i>
Falseflax, smallseed	<i>Camelina microcarpa</i>
Fleabane, annual	<i>Erigeron annus</i>
Goosefoot, nettleleaf	<i>Chenopodium murale</i>
Henbit	<i>Lamium amplexicaule</i>

Common Name	Scientific Name
Horseweed/Marestail	<i>Conyza canadensis</i>
Jimsonweed	<i>Datura stramonium</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mayweed	<i>Anthemis cotula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea.</i>
Morningglory, tall	<i>Ipomoea purpurea</i>
Mustard, black	<i>Brassica nigra</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, tansy	<i>Descurainia pinnata</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Mustard, wild	<i>Brassica kaber</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, cutleaf	<i>Solanum triflorum</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pepperweed, Virginia	<i>Lepidium virginicum</i>
Pigweed, prostrate	<i>Amaranthus, blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus, albus</i>
Prickly sida (Teaweed)	<i>Sida spinosa</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Rocket, London	<i>Sisymbrium irio</i>

Common Name	Scientific Name
Sesbania, hemp	<i>Sesbania exaltata</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Sicklepod	<i>Senna obtusifolia</i>
Smartweed (lady's thumb)	<i>Polygonum persicaria</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Spanish needles	<i>Bidens bipinnata</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, leafy	<i>Euphorbia esula</i>
Spurry, corn	<i>Spergula arvensis.</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Canada	<i>Cirsium arvense</i>
Thistle, Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>



## 8.0 CROP USE DIRECTIONS

### 8.1 Dicamba-Tolerant Cotton

#### 8.1.1 Preplant, At-Planting, Preemergence or Postemergence (In-Crop) Application

Crop			
Dicamba-tolerant cotton			
Target Weeds	Rate (pt/A)	Application Timing	Use Directions
Weeds listed in Section 7.1 & 7.2	3.53	<p><b>Burndown/Preplant Application:</b></p> <p>Apply prior to planting crop.</p> <p><b>At-Planting and Preemergence Application</b></p> <p>Apply during planting or after planting but <b>before</b> crop emergence.</p>	<p>Use only in: AR, KS, LA, MS, NM, OK, TN (excluding Wilson County), TX (excluding Gaines County) and the Boot Heel of MO.</p> <p>For emerged broadleaf weeds apply as a broadcast spray to small weeds that are less than 4 inches in height.</p>
		<p><b>Postemergence (In-crop) Application</b></p> <p>Postemergence in-crop applications can be made over-the-top of dicamba-tolerant cotton through 6-leaf cotton or up to and including July 30, whichever comes first, <b>unless additional state specific use restrictions are specified.</b></p>	<p>For use only in States specified in Section 6.1. This product must not be used in a county that has been explicitly prohibited on this label.</p> <p>Apply as a postemergence broadcast spray to small broadleaf weeds that are less than 4 inches in height.</p> <p>If at least ½ inch of rainfall does not occur within 10 days after application, cultivate shallowly.</p> <p>Crop canopy interference can reduce spray coverage on target weeds and soil, and hinder weed control. Use higher spray volumes (greater than 15 gallons per acre) under these conditions.</p> <p>For grass weed control, apply before grass weeds emerge or after clean cultivation.</p>
<p><b>Precautions:</b></p> <ul style="list-style-type: none"> <li>For preplant application, to the extent possible, avoid moving treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.</li> <li>If heavy rainfall occurs soon after application, crop injury may occur. Injury will be more severe in poorly drained areas where water stands for several hours or days, or where the seeding slit has not been properly closed.</li> </ul>			

**USE RESTRICTIONS**

- 1) Refer to **Section 6.1** for additional product use restrictions.
- 2) **Maximum Single Application Rate:** 3.53 pt/A (equivalent to 0.5 lb dicamba ae/A and 1.0 lb S-metolachlor/A)
- 3) **Maximum Annual Rate:** 7.06 pt/A/year (equivalent to 1 lb dicamba ae/A and 2.0 lb S-metolachlor/A)
  - a. **DO NOT** exceed 1.9 lb ai/A/year of S-metolachlor-containing products on coarse-textured soils.
  - b. **DO NOT** exceed 2.48 lb ai/A/year of S-metolachlor-containing products on medium- or fine-textured soils.
  - c. **DO NOT** exceed 2.0 lb ae/A/year of dicamba-containing products.
- 4) **DO NOT** apply less than 3.53 pt of this product/A (equivalent to 0.5 lb dicamba ae/A and 1.0 lb S-metolachlor/A).
- 5) **DO NOT** make more than one preplant or at-planting or preemergence application, and/or one postemergence (In-crop) application on medium-or fine-textured soils.
- 6) **DO NOT** make more than one application on coarse-textured soils.
- 7) **DO NOT** use on sand or loamy sand soils.
- 8) **DO NOT** use on Taloka silt loam.
- 9) **DO NOT** use where water is likely to “pond” over the bed.
- 10) **DO NOT** apply to non-dicamba-tolerant cotton.
- 11) **DO NOT** incorporate A21472 Plus VaporGrip Technology if applied prior to planting, or crop injury may result.
- 12) **DO NOT** use in Gaines County, TX; Wilson County, TN; or Palm Beach County, FL, or Nassau and Suffolk Counties, NY.
- 13) **DO NOT** graze or feed treated forage or fodder to livestock.
- 14) **Pre-harvest Interval (PHI):** 100 days

## 8.2 Dicamba-Tolerant Soybean

### 8.2.1 Preplant, At-Planting, Preemergence or Postemergence (In-Crop) Application

Crop			
Dicamba-tolerant soybeans			
Target Weed	Rate (pt/A)	Application Timing	Use Directions
Weeds listed in Section 7.1 & 7.2	3.53	<p><b>Preplant Application:</b></p> <p>Apply prior to planting crop.</p> <p><b>At-Planting and Preemergence Application:</b></p> <p>Apply during planting or after planting but <b>before</b> crop emergence.</p>	<p>For use only in States specified in <b>Section 6.1</b>. This product must not be used in a county that has been explicitly prohibited on this label.</p> <p>For emerged broadleaf weeds, apply as a broadcast spray to small weeds that are less than 4 inches in height.</p>
		<p><b>Postemergence (In-crop) Application</b></p> <p>Postemergence in-crop applications can be made over-the-top of dicamba-tolerant soybeans through V4 soybeans or up to and including June 30, whichever comes first, <b>unless additional state specific soybean use restrictions are specified.</b></p>	<p><b>For Postemergence Applications:</b></p> <p>For emerged broadleaf weeds, apply as a broadcast spray to small weeds that are less than 4 inches in height.</p> <p>For grass weed control, apply before grass weeds emerge.</p> <p>Dicamba-tolerant soybeans may exhibit leaf drooping following postemergence application. This response is transient and the soybeans will fully recover.</p>
<p><b>Precautions:</b></p> <ul style="list-style-type: none"> <li>For preplant application, to the extent possible, avoid moving treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate:</b> 3.53 pt/A (equivalent to 0.5 lb dicamba ae/A and 1.0 lb S-metolachlor/A)</li> <li><b>Maximum Annual Rate:</b> 7.06 pt/A/year (equivalent to 1 lb dicamba ae/A and 2.0 lb S-metolachlor/A) <ol style="list-style-type: none"> <li><b>DO NOT</b> exceed 3.71 lb ai/A/year of S-metolachlor-containing products.</li> <li><b>DO NOT</b> exceed 2.0 lb ae/A/year of dicamba-containing products.</li> </ol> </li> <li><b>DO NOT</b> apply less than 3.53 pt of this product/A (equivalent to 0.5 lb dicamba ae/A and 1.0 lb S-metolachlor/A).</li> <li><b>DO NOT</b> make more than one preplant or at-planting or preemergence application, and/or one</li> </ol>			

- postemergence (In-crop) application.
- 6) **DO NOT** apply to non-dicamba-tolerant soybeans.
  - 7) **DO NOT** use in Wilson County, TN; Palm Beach County, FL, or Nassau and Suffolk Counties, NY.
  - 8) **DO NOT** feed treated forage or hay to livestock for 30 days following a preplant, at-planting, or preemergence application.
  - 9) **DO NOT** graze or feed treated forage or hay to livestock following a postemergence application.
  - 10) **Pre-harvest Interval (PHI):** 75 days

## 9.0 STORAGE AND DISPOSAL

### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Keep container closed to prevent spills and contamination.

#### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

#### Container Handling (less than or equal to 5 gallons)

**Non-refillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling (greater than 5 gallons)

**Refillable container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Container Handling (greater than 5 gallons)**

**Non-refillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

## **10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold Syngenta and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES**

**(INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## 11.0 [APPENDIX]

### 11.1 [A21472 Plus VaporGrip Technology Use Summary Table]

[Start of Optional Text]

**IMPORTANT:** The table below is a summary of the Crop Use Directions for A21472 Plus VaporGrip Technology. However, it is important for the user to read and follow the complete instructions contained within this label.

Crop or Crop Group or Subgroup with examples	Maximum A21472 Plus VaporGrip Technology Rate per Application (pt/A)	Minimum Application Interval (days)	Pre-Harvest Interval (PHI days)	Maximum A21472 Plus VaporGrip Technology Rate per Year (fl oz/A)
Dicamba-Tolerant Cotton	3.53 <sup>1</sup>	NA	100	7.06 <sup>2</sup>
Dicamba-Tolerant Soybeans	3.53 <sup>1</sup>	NA	75	7.06 <sup>2</sup>

<sup>1</sup> 3.53 pt/A is equivalent to 0.5 lb dicamba ae/A and 1.0 lb S-metolachlor/A

<sup>2</sup> 7.06 pt/A is equivalent to 1 lb dicamba ae/A and 2.0 lb S-metolachlor/A

## 11.2 [State Additional Labeling]

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

DICAMBA	GROUP	4	HERBICIDE
S-METOLACHLOR	GROUP	15	HERBICIDE

Primary Brand Name:

**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:

**Tavium® Plus VaporGrip® Technology**

Active Ingredients:

Diglycolamine salt of dicamba\*.....17.7%

S-metolachlor\*\*.....24.0%

---

Other Ingredients: 58.3%

---

Total: 100.0%

\*CAS No. 104040-79-1

\*\*CAS No. 87392-12-9

A21472 Plus VaporGrip Technology is a capsule suspension (CS) formulation containing 1.12 pounds of dicamba acid equivalent (ae) and 2.26 pounds of S-metolachlor per U.S. gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the**

user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.



## **USE RESTRICTIONS**

Users must comply with additional requirements including state restrictions found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com). The user must check [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com) no more than 7 days before application of this product for additional labeling.

**Record keeping Requirement:** Maintain a copy of the product label and any labeling that supplements the product label.



### 11.3 [Additional Iowa-specific restrictions as found on www.TaviumApplicationRequirements.com]

#### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

**These instructions for A21472 Plus VaporGrip® Technology are only applicable to the state of Iowa.**

Primary Brand Name:

**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:

**Tavium® Plus VaporGrip® Technology**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Additional specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)**



### Application Directions

In order to apply A21472 Plus VaporGrip® Technology in Iowa, the following requirements apply:

#### Cotton:

- DO NOT apply after June 12 or 1<sup>st</sup> square, whichever comes first

Soybean:

- DO NOT apply after June 12 or V4, whichever comes first.

## 11.4 [Additional Minnesota-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

**These instructions for A21472 Plus VaporGrip® Technology are only applicable to the state of Minnesota.**

Primary Brand Name:  
**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:  
**Tavium® Plus VaporGrip® Technology**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

### Application Directions

In order to apply A21472 Plus VaporGrip® Technology in Minnesota, the following requirements apply:

DO NOT apply south of Interstate 94 after June 12. DO NOT apply north of Interstate 94 after June 30.

Statewide restriction: DO NOT apply if the air temperature of the field at the time of application is over 85 degrees Fahrenheit or if the National Weather Service's forecasted high temperature for the nearest available location for the day exceeds 85 degrees Fahrenheit. Forecasted temperature must be recorded at the start of application.

## 11.5 [Additional Illinois-specific restrictions as found on www.TaviumApplicationRequirements.com]

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

#### **Syngenta Crop Protection, LLC**

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

**These instructions for A21472 Plus VaporGrip® Technology are only applicable to the state of Illinois**

Primary Brand Name:

**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:

**Tavium® Plus VaporGrip® Technology**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Additional specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)**

**syngenta**

### **Application Directions**

In order to apply A21472 Plus VaporGrip® Technology in Illinois, the following requirements apply:

Cotton:

- DO NOT apply after June 12 or 1<sup>st</sup> square, whichever comes first

Soybean:

- DO NOT apply after June 12 or V4, whichever comes first.

## 11.6 [Additional Indiana-specific restrictions as found on www.TaviumApplicationRequirements.com]

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

**These instructions for A21472 Plus VaporGrip® Technology are only applicable to the state of Indiana**

Primary Brand Name:

**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:

**Tavium® Plus VaporGrip® Technology**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Additional specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)**

**syngenta**

### Application Directions

In order to apply A21472 Plus VaporGrip® Technology in Indiana, the following requirements apply:

#### Cotton:

- DO NOT apply after June 12 or 1<sup>st</sup> square, whichever comes first

Soybean:

- DO NOT apply after June 12 or V4, whichever comes first.



## 11.7 [Additional South Dakota-specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)]

### RESTRICTED USE PESTICIDE

To be used by certified applicators only; NOT to be used by non-certified persons working under the supervision of a certified applicator, except that non-certified persons may transport containers.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**This labeling expires 12/20/2025. DO NOT use or distribute this product after 12/20/2025.**

**These instructions for A21472 Plus VaporGrip® Technology are only applicable to the state of South Dakota**

Primary Brand Name:

**A21472 Plus VaporGrip® Technology**

Alternate Brand Name:

**Tavium® Plus VaporGrip® Technology**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1623

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using A21472 Plus VaporGrip® Technology as permitted according to this labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Additional specific restrictions as found on [www.TaviumApplicationRequirements.com](http://www.TaviumApplicationRequirements.com)**

**syngenta**

### Application Directions

In order to apply A21472 Plus VaporGrip® Technology in South Dakota, the following requirements apply:

- DO NOT apply after June 20.

[End of Optional Text]

---

Tavium®, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Plus VaporGrip® Technology is a trademark of Bayer CropScience LLC.

©20XX Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.
--

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina, 27419-8300

A21472 Plus VaporGrip Technology 1623 MAS 0422 AMEND-D.DEC2022-CL - di - 2/10/2023  
000100-01623.20221215D.A21472\_PLUS\_VAPORGRIP\_TECHNOLOGY.AMEND.DEC2022-CL