



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 21, 2020

Ronald E. Hampton, Ph.D.
Regulatory Product Manager
Syngenta Professional Solutions
P.O. Box 18300
Greensboro, NC 27419-8300

Subject: Registration Review Label Mitigation for Indoxacarb
Product Name: Provaunt WDG
EPA Registration Number: 100-1636
Application Date: May 17, 2019
Decision Number: 565397

Dear Dr. Hampton:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Indoxacarb Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2
EPA Reg. No. 100-1636
Decision No. 564963

If you have any questions about this letter, please contact Jaclyn Pyne by phone at 703-347-0445, or via email at pyne.jaclyn@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

(Master label)

INDOXACARB **GROUP** **22A** **INSECTICIDE**

PROVAUNT® WDG

Insecticide

A water-dispersible granular insecticide for the control of lepidopterous larvae, including armyworms, cutworms, sod webworms, bagworms, fall webworms, gypsy moth caterpillars, tent caterpillars, tussock moth caterpillars, yellownecked caterpillars, and ants (excluding fire, harvester, pharaoh, and carpenter), and other listed pests infesting landscape and recreational turf grass, including golf courses and landscape ornamentals.

Active Ingredient:

Indoxacarb*

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4(trifluoromethoxy) phenyl]amino]carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate20.0%

Other Ingredients 80.0%

Total: 100.00%

Provaunt® WDG is a water-dispersible granule (WG) that contains 0.2 lb indoxacarb per lb product.

*Indoxacarb belongs to the oxadiazine chemical class.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-1636

EPA Est. XXXX

[Batch Code: _____] (For nonrefillables only.)

Net Contents




TABLE OF CONTENTS

1.0 FIRST AID

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

2.2 Personal Protective Equipment (PPE)

2.2.1 User Safety Requirements

2.2.2 User Safety Recommendations

2.3 Environmental Hazards

DIRECTIONS FOR USE

3.0 PRODUCT INFORMATION

3.1 Insect Resistance Management

3.1.1 Integrated Pest Management (IPM) Programs

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

4.2 Application Equipment

4.3 Mixing Directions

4.3.1 Provaunt WDG Alone

4.3.2 Tank-Mix Precautions

4.3.3 Tank-Mix Compatability

4.3.4 Provaunt WDG In Tank Mixtures

5.0 RESTRICTIONS AND PRECAUTIONS

5.1 Use Restrictions

5.2 Spray Drift Management

6.0 TURF AND ORNAMENTAL USE DIRECTIONS

6.1 Application Dilution Chart

6.2 Recreational Turfgrass and Landscape Areas

6.3 Ornamental Plants within Exterior and Interior Landscapes

7.0 STORAGE AND DISPOSAL

8.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p style="text-align: center;">HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p>	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

2.2.1 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 User Safety Recommendations

Users Should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this

product will be reduced by avoiding applications when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Do not apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters).

This product is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in or adjacent to the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. Syngenta will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Syngenta. User assumes all risks associated with such non-specified use.

3.0 PRODUCT INFORMATION

Provaunt WDG is a water-dispersible granular insecticide that may be applied as a foliar or broadcast spray to control lepidopterous larvae, such as armyworms, cutworms, sod webworms, bagworms, fall webworms, gypsy moth caterpillars, tent caterpillars, tussock moth caterpillars, yellownecked caterpillars, and ants (excluding fire, harvester, pharaoh, and carpenter), and other listed pests on landscape and recreational turf grass, including golf courses and landscape ornamentals such as trees, shrubs, foliage plants, flowers, and non-bearing fruit and nut trees (i.e., trees that will not produce a raw agricultural commodity during the season of application).

Provaunt WDG may only be used on plants or turf grasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, interior plantscapes, ornamental gardens, parks, playgrounds, schools, golf courses (such as tee box areas, roughs, fairways, greens, and collars), athletic fields and other landscaped areas.

3.1 Insect Resistance Management

INDOXACARB	GROUP	22A	INSECTICIDE
------------	-------	-----	-------------

For resistance management, Provaunt WDG contains a Group 22A insecticide. Any insect population may contain individuals naturally resistant to Provaunt WDG and other Group 22A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same treated areas. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Provaunt WDG or other Group 22A insecticides with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classifications are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local Syngenta representative, retailer, or extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Syngenta at 1-866-SYNGENT(A) (866-796-4368).

Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

3.1.1 Integrated Pest Management (IPM) Programs

Provaunt WDG is recommended for IPM programs on turf and landscape ornamentals as it does not directly impact natural arthropod predator and parasitoid populations, including ladybird beetles, lacewings, minute pirate bugs and predatory mites. The feeding behavior of predatory beneficial arthropods will aid in extending natural control of other insect and mite pests and will reduce the possibility of secondary pest outbreaks. Provaunt WDG will reduce the target pest species that may serve as a food source for beneficial arthropods. If Provaunt WDG is tank-mixed with an insect control product that negatively impacts beneficial arthropods, the full benefit of Provaunt WDG to the IPM program will not be realized.

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

Provaunt WDG must be diluted with water before application.

Provaunt WDG applications should target listed pests. Use sufficient water to obtain thorough, uniform coverage. Apply repeat applications after 7 days, if necessary.

4.2 Application Equipment

Application equipment must be clean and free of previous pesticide deposits before mixing Provaunt WDG.

Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all application equipment to reduce the risk of forming hardened deposits that might become difficult to remove. Drain application equipment. Thoroughly rinse application equipment and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources or desirable vegetation. Dispose of waste rinsewater in accordance with local regulations.

4.3 Mixing Directions

4.3.1 Provaunt WDG Alone

1. Use clean, well maintained application equipment.
2. Fill sprayer tank 1/4 to 1/2 full of water.
3. Add Provaunt WDG directly to the sprayer tank. If tank mixing Provaunt WDG with other products, see the Tank-Mixtures section below.
4. Mix thoroughly to fully disperse the insecticide and continue agitation to keep the insecticide in suspension. Use mechanical or hydraulic agitation. Do not use air agitation.
5. The mixture should not be stored in the spray tank overnight.

4.3.2 Tank-Mix Precautions

Provaunt WDG may be tank-mixed with other pesticides. When tank-mixing Provaunt WDG with other pesticides, observe all precautions and limitations on each separate product label. No label application rates may be exceeded. Provaunt WDG cannot be mixed with any product containing a label prohibition against such mixing.

4.3.3 Tank-Mix Compatability

The physical compatibility of Provaunt WDG may vary with different sources of pesticide products and local cultural practices. Any tank-mixture that has not been previously tested should be prepared on a small scale (pint or quart jar) using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.

4.3.4 Provaunt WDG In Tank Mixtures

Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after the addition of each product.

1. Water-soluble bags
2. Provaunt WDG and other water-dispersible granules
3. Wettable powders
4. Water-based suspension concentrates
5. Water-soluble concentrates
6. Oil-based suspension
7. Emulsifiable concentrates
8. Adjuvants, surfactants, oils
9. Soluble fertilizers
10. Drift retardants

5.0 RESTRICTIONS AND PRECAUTIONS

5.1 Use Restrictions

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production.
- **DO NOT** formulate this product into other end-use products without written permission of Syngenta Crop Protection, LLC
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product in commercial nurseries and greenhouses or on sod farms.

- **DO NOT** apply more than 36 ounces (equivalent to 0.45 lb of active ingredient) of product per acre per season in broadcast applications.
- Wait a minimum of 7 days to retreat.
- Keep people and pets away from treated areas until treatment has dried.

5.2 Spray Drift Management

SPRAY DRIFT

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or plant canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size — Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT — Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift

Boom-less Ground Applications

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

- Take precautions to minimize spray drift.

6.0 TURF AND ORNAMENTAL USE DIRECTIONS

6.1 Application Dilution Chart

Application Volume (gallons per 1,000 sq ft)	Application Rate			Amount of Provaunt WDG needed for these volumes of finished spray (oz)			
	(oz per acre)	(oz per 1,000 sq ft)	(lb AI per acre)	1 gallon	5 gallons	10 gallons	100 gallons
1	3	0.07	0.038	0.07	0.34	0.69	6.9
1	6	0.14	0.075	0.14	0.69	1.38	13.8
1	9	0.21	0.113	0.21	1.03	2.07	20.7
1	12	0.28	0.150	0.28	1.38	2.75	27.5
1	15	0.34	0.188	0.34	1.72	3.44	34.4
1	18	0.41	0.225	0.41	2.07	4.13	41.3

2	3	0.07	0.038	0.03	0.17	0.34	3.4
2	6	0.14	0.075	0.07	0.34	0.69	6.9
2	9	0.21	0.113	0.10	0.52	1.03	10.3
2	12	0.28	0.150	0.14	0.69	1.38	13.8
2	15	0.34	0.188	0.17	0.86	1.72	17.2
2	18	0.41	0.225	0.21	1.03	2.07	20.7
3	3	0.07	0.038	0.02	0.11	0.23	2.3
3	6	0.14	0.075	0.05	0.23	0.46	4.6
3	9	0.21	0.113	0.07	0.34	0.69	6.9
3	12	0.28	0.150	0.09	0.46	0.92	9.2
3	15	0.34	0.188	0.11	0.57	1.15	11.5
3	18	0.41	0.225	0.14	0.69	1.38	13.8
4	3	0.07	0.038	0.02	0.09	0.17	1.7
4	6	0.14	0.075	0.03	0.17	0.34	3.4
4	9	0.21	0.113	0.05	0.26	0.52	5.2
4	12	0.28	0.150	0.07	0.34	0.69	6.9
4	15	0.34	0.188	0.09	0.43	0.86	8.6
4	18	0.41	0.225	0.10	0.52	1.03	10.3
5	3	0.07	0.038	0.01	0.07	0.14	1.4
5	6	0.14	0.075	0.03	0.14	0.28	2.8
5	9	0.21	0.113	0.04	0.21	0.41	4.1
5	12	0.28	0.150	0.06	0.28	0.55	5.5
5	15	0.34	0.188	0.07	0.34	0.69	6.9
5	18	0.41	0.225	0.08	0.41	0.83	8.3

• To convert from ounces to grams, multiply by 28.35 (1 ounce = 28.35 grams).

DO NOT use household utensils to measure Provaunt WDG

6.2 Recreational Turfgrass and Landscape Areas

Target Pest	Amount of Product per acre (oz)	Amount of Product per 1,000 sq ft (oz)	Amount of AI per acre (lb)
Caterpillars (including armyworms, cutworms and sod webworms)	3 to 6	0.07 to 0.14	0.038 to 0.075
European Crane Fly Larvae	9 to 18	0.21 to 0.41	0.113 to 0.225
Grasshoppers Mole Crickets* Annual Bluegrass Weevil Larvae*	18	0.41	0.225
Ants (excluding fire, harvester, pharaoh, and carpenter)	9 to 18	0.21 to 0.41	0.113 to 0.225

*Not approved for use in California

NOTE: For mole cricket control, make 1-2 applications on a 2 to 4 week interval when nymphs or adults are actively foraging. For best results, apply in morning when turf is wet or pre-wet with irrigation. Water in immediately after application with 1/8” of irrigation.

To ensure optimum control of lepidopterous larvae, delay watering (irrigation) and mowing for 24 hours after application unless instructed otherwise. If the grass area is being maintained at a mowing height of greater than 1 inch, the higher application rate (6 ounces per acre) may be required during periods of high pest pressure.

For maximum residual effectiveness, apply Provaunt WDG at up to 18 ounces per acre (0.412 ounces per 1,000 square feet = 0.225 lb AI/A).

Annual Bluegrass Weevil Control Program Targeting Overwintered Adults and First Generation Larvae*			
Target Pest	Amount of Product per acre (oz)	Amount of Product per 1,000 sq ft (oz)	Amount of AI per acre (lb)
Annual Bluegrass Weevil Adults (Preventative)	18	0.41	0.225
Approximately three weeks after the Preventative treatment (above) apply the treatment (below):			
Annual Bluegrass Weevil Larvae	18	0.41	0.225

* Not approved for use in California

6.3 Ornamental Plants within Exterior and Interior Landscapes

Target Pest	Amount of Product per 100 Gallons (oz)	Amount of AI per 100 Gallons (lb)
Leaf-feeding caterpillars , such as bagworms gypsy moth caterpillars tent caterpillars (including eastern tent caterpillar) tussock moth caterpillars (including white-marked tussock moth) webworms (including fall webworm) yellownecked caterpillars	1.88 to 3.75	0.023 to 0.046
European pine sawfly	3.75 to 7.5	0.046 to 0.092
Potato leafhopper	7.5	0.092

Apply the desired application rate of Provaunt WDG as a full-coverage foliar spray. Repeat treatment as directed to achieve best control using higher application rates within the listed rate range as pest pressure and foliage area increases. Repeat applications must be limited to no more than once per 7 days.

Certain plant species or cultivars may be sensitive to the final spray solution. If local experience is not available, then a small number of plants should be treated and observed for phytotoxicity for at least one week before making application to the entire planting.

7.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

Pesticide Disposal

Do not contaminate water, food or feed by storage and disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

Container Handling

Non-refillable Plastic and Metal Containers (less than or equal to 50 pounds): Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. For Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. For Metal Containers, offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Non-refillable Plastic and Metal Containers (greater than 50 pounds):

Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. For Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Non-refillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (size or shape too large to be tipped, rolled or turned upside down):

Non-refillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. For Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Non-refillable Paper or Plastic Bags, or Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC):

Non-refillable container. Do not reuse or refill this container. Completely empty paper bag, plastic bag, or fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper bag, plastic bag, or fiber sack in a sanitary landfill, or by incineration, or by other procedures approved by state and local ordinances.

Fiber Drums With Liners:

Non-refillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of liner in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

8.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Provaunt®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

©20XX Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

Provaunt WDG 1636 MAS 0518 AMEND 0519-CL – di – 5-14-19
000100-01636.20190516.Provaunt WDG.AMEND.MAY2019-CL