



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

100-1648

Date of Issuance:

8/30/19

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Saltro

Name and Address of Registrant (include ZIP Code):

Adora Clark  
 Federal Team Lead, Fungicides  
 Syngenta Crop Protection, LLC  
 PO Box 18300  
 Greensboro, NC 27419

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

 FOR

Cynthia Giles-Parker, Chief  
 Fungicide Branch, Registration Division (7505P)

Date:

8/30/19

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 100-1648.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 5/11/2018
- Alternate CSF 1 dated 5/11/2018

If you have any questions, please contact Lindsay Roe by phone at 703-347-0506, or via email at [roe.lindsay@epa.gov](mailto:roe.lindsay@epa.gov).

Enclosure


[Master Label]

**PYDIFLUMETOFEN** | **GROUP 7** | **FUNGICIDE**

**Saltro®**

**Fungicide**

An ADEPIDYN™ brand fungicide



Active Ingredient:	
Pydiflumetofen* .....	41.7%
Other Ingredients:	58.3%
Total:	100.0%

\*CAS No. 1228284-64-7

Saltro® is a flowable concentrate for seed treatment containing 4.17 lb pydiflumetofen per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## **CAUTION**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-

EPA Est.

\_\_\_\_\_  
Net Contents

[Batch Code: \_\_\_\_\_ ] (*For nonrefillables only.*)

## **TABLE OF CONTENTS**

### **1.0 FIRST AID**

### **2.0 PRECAUTIONARY STATEMENTS**

#### **2.1 Hazards to Humans and Domestic Animals**

#### **2.2 Personal Protective Equipment (PPE)**

##### 2.2.1 User Safety Requirements

##### 2.2.2 Engineering Controls

##### 2.2.3 User Safety Recommendations

#### **2.3 Environmental Hazards**

##### 2.3.1 Groundwater Advisory

##### 2.3.2 Surface Water Advisory

#### **2.4 Physical or Chemical Hazards**

## **DIRECTIONS FOR USE**

### **3.0 PRODUCT INFORMATION**

#### **3.1 Resistance Management**

### **4.0 APPLICATION DIRECTIONS**

#### **4.1 Mixing Directions**

##### 4.1.1 Tank-Mix Precautions

##### 4.1.2 Tank Mixtures

### **5.0 ROTATIONAL CROP RESTRICTIONS**

### **6.0 SEED CONTAINER LABEL REQUIREMENTS**

### **7.0 Seed Treatment Directions**

#### **7.1 Rapeseed (including Canola) – Crop Subgroup 20A**

#### **7.2 Soybean**

### **8.0 STORAGE AND DISPOSAL**

### **9.0 CONDITIONS OF SALE AND LIMITATION OF LIABILITY**

## 1.0 FIRST AID

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p><b>HOT LINE NUMBER</b>            For 24-Hour Medical Emergency Assistance (Human or Animal)            Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)            Call  <b>1-800-888-8372</b></p>	

## 2.0 PRECAUTIONARY STATEMENTS

### 2.1 Hazards to Humans and Domestic Animals

#### **CAUTION**

Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### 2.2 Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

#### 2.2.1 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### 2.2.2 Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### 2.2.3 User Safety Recommendations

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## 2.3 Environmental Hazards

Pydiflumetofen is toxic to fish, aquatic invertebrates, and oysters and shrimp. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### 2.3.1 Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

### 2.3.2 Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of pydiflumetofen from runoff water and sediment. Runoff of this product will be

reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

## 2.4 Physical or Chemical Hazards

Do not mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY; POOR DISEASE CONTROL; AND/OR ILLEGAL RESIDUES.**

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. Exception:** If the seed is treated with the product and the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

## 3.0 PRODUCT INFORMATION

Salstro is a seed treatment fungicide which provides protection against listed soil-borne and seed-borne diseases of rapeseed (including canola) crop subgroup 20A, and soybean. Salstro is effective against seed and seedling blight or damping-off caused by listed seed- and soil-borne pathogens, including *Fusarium virguliforme*, the causal organism of Sudden Death Syndrome (SDS) in soybean and seed- and soil-borne black leg (*Leptoshaeria maculens*) in canola. Salstro also provides activity against listed plant parasitic nematodes. Where rate ranges are shown, use the higher rate when disease pressure is expected to be severe.

Read all label directions for use. All applications must be made according to the use directions that follow.

### 3.1 Resistance Management

PYDIFLUMETOFEN	GROUP	7	FUNGICIDE
----------------	-------	---	-----------

For resistance management, Salstro contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to Salstro and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Salstro or other Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Syngenta Crop Protection at 1-866-796-4368. You can also contact your university extension specialist to report resistance.



## 4.0 APPLICATION DIRECTIONS

Apply Saltro as a water-based slurry utilizing standard seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Saltro does not control diseases caused by *Pythium* spp., *Phytophthora* spp or *Rhizoctonia* spp. If these diseases are expected to be a problem, apply Saltro with other seed treatment products that contain mefenoxam and/or fludioxonil as active ingredients.

This product must be used only by commercial seed treatment facilities. Do not use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

### 4.1 Mixing Directions

Thoroughly mix the specified amount of Saltro into the required amount of water and other seed treatment products for better application performance. Follow the manufacturer's application instructions for the seed treatment equipment being used. It's important to homogenize products prior to use and to maintain constant agitation of the slurry during the treatment.

This product does not contain dye and treated seed is not covered by an appropriate tolerance, tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR 153.155, therefore, all seed treated commercially with this product must be colored with an EPA-approved dye or colorant of a suitable color to prevent accidental use as food for man or feed for animals.

Allow seed to dry before bagging.

#### 4.1.1 Tank-Mix Precautions

If seed treatment products containing mefenoxam and/or fludioxonil are not registered on that crop, apply Saltro as a tank mix with other labeled seed treatment products that have activity against these diseases.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### 4.1.2 Tank Mixtures

**Rapeseed (including canola) Crop Subgroup 20A:** For additional control of certain seed- and soil-borne pathogens, Saltro may be combined with a seed treatment product containing, difenoconazole, mefenoxam, and fludioxonil. On canola only, Saltro may be combined with a seed treatment product containing thiamethoxam for insect control.

**Soybean:** Additional protection from common seed-borne and soil-borne pathogens may be achieved by adding seed treatment products containing mefenoxam, azoxystrobin, fludioxonil and/or thiabendazole. Saltro may also be mixed with CruiserMaxx® Vibrance® (EPA Reg. No. 100-1508) for protection from additional early-season diseases and insect pests. Alternatively, Saltro may also be mixed with Clariva® Elite Beans (EPA Reg. No. 100-1620) for soybean cyst nematode (SCN) protection and from additional early-season diseases and insect pests.

## 5.0 ROTATIONAL CROP RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from seed treated with Saltro, the field may be replanted according to the following schedule:

Crop, Crop Group, or Crop Subgroup	Plant-Back Interval
Blueberry and Bushberry, Crop Subgroup 13-07B Brassica Head and Stem Vegetable, Crop Group 5-16 Bulb Vegetables, Crop Group 3-07 Canola (Rapeseed Crop Subgroup 20A) Cereals (barley, oats, wheat, triticale, rye) Citrus Fruit, Crop Group 10-10 Corn Corn, sweet Cottonseed, Crop Subgroup 20C Cucurbit Vegetables, Crop Group 9 Fruiting Vegetables, Crop Group 8-10 Leaf Petiole Vegetables, Crop Subgroup 22B Leafy Greens, Crop Subgroup 4-16A Leaves of Root and Tuber Vegetables, Crop Group 2 Legume Vegetables, Succulent or Dried, Crop Group 6, Except Cowpea Forage and Cowpea Hay Mustard Greens and Brassica Leafy Greens, Subgroup 4-16B Peanut Peppers Pome Fruit, Crop Group 11-10 Potato Quinoa Root Vegetables, Crop Subgroup 1A Soybean, Except Forage, Hay, and Silage Sorghum Stone Fruit Crop Group 12-12 Strawberry, and Low Growing Berry, Crop Subgroup 13-07G, Except Cranberry Sunflower, Crop Subgroup 20B Tomatoes Tree Nuts, Crop Group 14-12 Tuberous and Corm Vegetables, Crop Subgroup 1C	0 days
All other crops Intended for Food and Feed	365 days

## 6.0 SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with pydiflumetofen fungicide.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with Salstro fungicide:

- Store away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwater.
- Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- Treated seed must be planted into the soil at a depth of at least 1 inch.
- Do not feed or harvest soybean hay, forage, and silage.
- Dispose of seed packaging in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice.
- For Rapeseed (including canola) Crop Subgroup 20A **DO NOT** exceed 0.291 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.
- For soybeans, **DO NOT** exceed 0.357 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.
- The maximum number of plantings of seed treated with Pydiflumetofen to the same crop area is 2 per calendar year.
- In the event of crop failure or harvest of a crop grown from seed treated with Salstro, crops may be replanted according to the following schedule:

Crop, Crop Group, or Crop Subgroup	Plant-Back Interval
Blueberry and Bushberry, Crop Subgroup 13-07B Brassica Head and Stem Vegetable, Crop Group 5-16 Bulb Vegetables, Crop Group 3-07 Canola (Rapeseed Crop Subgroup 20A) Cereals (barley, oats, wheat, triticale, rye) Citrus Fruit, Crop Group 10-10 Corn Corn, sweet Cottonseed, Crop Subgroup 20C	0 days

<b>Crop, Crop Group, or Crop Subgroup</b>	<b>Plant-Back Interval</b>
Cucurbit Vegetables, Crop Group 9 Fruiting Vegetables, Crop Group 8-10 Leaf Petiole Vegetables, Crop Subgroup 22B Leafy Greens, Crop Subgroup 4-16A Leaves of Root and Tuber Vegetables, Crop Group 2 Legume Vegetables, Succulent or Dried, Crop Group 6, Except Cowpea Forage and Cowpea Hay Mustard Greens and Brassica Leafy Greens, Subgroup 4-16B Peanut Peppers Pome Fruit, Crop Group 11-10 Potato Quinoa Root Vegetables, Crop Subgroup 1A Soybean, Except Forage, Hay, and Silage Sorghum Stone Fruit Crop Group 12-12 Strawberry, and Low Growing Berry, Crop Subgroup 13-07G, Except Cranberry Sunflower, Crop Subgroup 20B Tomatoes Tree Nuts, Crop Group 14-12 Tuberous and Corm Vegetables, Crop Subgroup 1C	
All other crops Intended for Food and Feed	365 days

## 7.0 Seed Treatment Directions

### 7.1 Rapeseed (including Canola) – Crop Subgroup 20A

Crops (Including cultivars, varieties, and/or hybrids of these)			
Borage	Gold of pleasure	Mustard seed	
Canola	Hare's ear mustard	Oil radish	
Crambe	Lesquerella	Poppy seed	
Cuphea	Lunaria	Rapeseed	
Echium	Meadowfoam	Sesame	
Flax seed	Milkweed	Sweet rocket	
Target Disease/Pest	Use Rate		Notes
	fl oz/100 lb of seed	grams ai/100 kg of seed	
Control of seed- and air-borne blackleg caused by <i>Leptosphaeria maculans</i>	1.23 (80 ml/100 kg seed)	40	<p>For additional fungicidal control on canola only, Salstro may be mixed with Vibrance® Flexi (EPA Reg. No. 100-1588).</p> <p>For additional disease and insect control on canola only, Salstro may be mixed with Helix® Vibrance® (EPA Reg. No. 100-1528).</p> <p>For additional insect control on Rapeseed (including Canola) – Crop Subgroup 20A, Salstro may be mixed with Fortenza® (EPA Reg. No. 100-1420).</p> <p>For additional insect control on Canola only, Salstro may be mixed with Cruiser® 5FS (EPA Reg. No. 100-941).</p> <p>Follow all appropriate treating and use directions, precautions, etc. as specified on the respective product labels.</p>
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Refer to Section 3.1.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Plantings:</b> 2</li> <li><b>Sections 4.0, 4.1, 5.0, and 6.0</b> contain additional product use restrictions.</li> <li><b>DO NOT</b> exceed 0.291 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.</li> </ul>			

## 7.2 Soybean

Crops (Including cultivars, varieties, and/or hybrids of these)		
Soybean		
Target Disease/Pest	Use Rate*	
	fl oz formulated product	mg ai/seed
Sudden Death Syndrome ( <i>Fusarium virguliforme</i> )	1.52 fl oz/ 100 lbs soybean	0.075
Early-season Septoria Brown Spot ( <i>Septoria glycines</i> )	0.714 fl oz/ 140,000 soybean seeds	
Plant parasitic nematodes Soybean Cyst ( <i>Heterodera glycines</i> ) Root Knot ( <i>Meloidogyne incognita</i> ) Reniform ( <i>Rotylenchulus reniformis</i> ) Lesion ( <i>Pratylenchus</i> spp.) Lance ( <i>Hoplolaimus</i> spp.)		
*assuming 3,000 soybean seeds/pound		
<b>Resistance Management:</b>		
<ul style="list-style-type: none"> <li>Refer to Section 3.1.</li> </ul>		
<b>USE RESTRICTIONS</b>		
<ul style="list-style-type: none"> <li><b>Maximum Annual Plantings: 2</b></li> <li><b>Do not feed or harvest soybean hay, forage, and silage.</b></li> <li><b>Sections 4.0, 4.1, 5.0, and 6.0</b> contain additional product use restrictions.</li> <li><b>DO NOT</b> exceed 0.357 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.</li> </ul>		

## 8.0 STORAGE AND DISPOSAL

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### Pesticide Storage

Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

### **Pesticide Disposal**

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### **Container Handling [less than or equal to 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume – and recap. Shake for 30 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [greater than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [greater than 5 gallons]**

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume



application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

## 9.0 CONDITIONS OF SALE AND LIMITATION OF LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

---



ADEPIDYN™, Clariva®, Cruiser®, CruiserMaxx®, Fortenza®, Helix®, Vibrance®, the ALLIANCE FRAME the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Viton® is a registered trademark of E.I. DuPont de Nemours & Company, Inc.

©20XX Syngenta

For non-emergency (e.g. current product information), call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

Salstro RAUI MAS 0718 NEW-VER C – ep – 8/28/19  
000100-0RAUI.20180716B.SALTRO\_NEW\_0718.pdf