



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1662

Date of Issuance:

4/1/20

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Pitore Liquid Treatment

Name and Address of Registrant (include ZIP Code):

Ronald Hampton
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 100-1662."

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

Date:

4/1/20

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 12/05/2019
- Alternate CSF 1 dated 12/05/2019
- Alternate CSF 2 dated 12/05/2019

If you have any questions, please contact Kevin Ulrich by phone at 703-347-0464, or via email at ulrich.kevin@epa.gov.

Enclosure: Stamped Label

Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
CHLORANTRANILIPROLE	GROUP	28	INSECTICIDE

PITORE® LIQUID TREATMENT

Insecticide

For control of localized infestations of listed termites and wood-destroying beetles and borers.

Do not use this product as the sole source of control for active, structural infestations by subterranean termites. The purpose of this application is to kill workers or winged reproductive forms of termites that are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, or a soil or foundation treatment. For active, structural infestations by subterranean termites, this product can only be used to supplement a termiticide that is registered as a stand-alone termite treatment. This product will not eliminate termite infestations or provide protection against future infestations.

Active Ingredients:

Thiamethoxam ¹	17.5%
Chlorantraniliprole ²	8.8%
<hr/>	
Other Ingredients:	73.7%
Total:	100.0%

¹CAS No. 153719-23-4
²CAS No. 500008-45-7

Pitore Liquid Treatment is a suspension concentrate that contains 1.67 lb thiamethoxam and 0.835 lb chlorantraniliprole active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

[Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)]

See additional Precautionary Statements and Directions for Use inside booklet.

EPA Reg. No. 100-XXXX
EPA Est. XXXX

Net Contents

[Batch Code: _____] (For nonrefillables only.)




TABLE OF CONTENTS

1.0 FIRST AID

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

- 2.1 Hazards to Humans and Domestic Animals
- 2.2 Personal Protective Equipment (PPE)
- 2.3 User Safety Requirements
- 2.4 User Safety Recommendations
- 2.5 Environmental Hazards
 - 2.5.1 Groundwater Advisory
 - 2.5.2 Surface Water Advisory
 - 2.5.3 Pollinator Precautions
- 2.6 Physical or Chemical Hazards

DIRECTIONS FOR USE

3.0 PRODUCT INFORMATION

- 3.1 Resistance Management

4.0 APPLICATION DIRECTIONS

- 4.1 Mixing Directions
 - 4.1.1 Dilution Preparation
 - 4.1.2 Foam Preparation

5.0 RESTRICTIONS

- 5.1 Use Restrictions

6.0 USE DIRECTIONS

- 6.1 Application Areas
 - 6.1.1 Surface Treatment
 - 6.1.2 Wood Injection

7.0 SPECIFIC USE DIRECTIONS FOR REMEDIAL CONTROL OF LISTED PESTS

- 7.1 Drywood Termites
- 7.2 Limited Area Treatment of Subterranean Termites in and Around Existing Structures
- 7.3 Wood-Destroying Beetles

8.0 STORAGE AND DISPOSAL

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID

HOTLINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call
1-800-888-8372

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate; butyl rubber ≥ 14 mils; nitrile rubber ≥ 14 mils; neoprene rubber ≥ 14 mils; polyvinyl chloride (PVC) ≥ 314 mils; or Viton® ≥ 14 mils
- Shoes and socks

2.3 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.4 User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.5 Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply where/when conditions could favor runoff. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues on blooming plants or weeds. Do not apply this product or allow it to drift to blooming plants or weeds while bees are foraging in or adjacent to the treatment area.

2.5.1 Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

2.5.2 Surface Water Advisory

This product may impact surface water quality due to spray drift and runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, such as ponds, streams, and springs, will reduce the potential loading of thiamethoxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours. (See manual at the following Internet address:

<http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>)

2.5.3 Pollinator Precautions

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to:

www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

2.6 Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

3.0 PRODUCT INFORMATION

Apply Pitore Liquid Treatment as a remedial treatment to control localized infestations of:

- Drywood Termites: *Cryptotermes*, *Incisitermes* and *Marginitermes* species
- Subterranean Termites: *Coptotermes*, *Heterotermes* and *Reticulitermes* species
- Wood-destroying Beetles: True powderpost beetles (*Lyctidae* species), death-watch beetles and ambrosia or bark beetles

Apply Pitore Liquid Treatment as crack-and-crevice, structural void, wood injection, spot, and/or general surface treatment.

DO NOT use Pitore Liquid Treatment as a preventive or stand-alone termite treatment.



DO NOT apply Pitore Liquid Treatment while bees are foraging.

DO NOT apply to plants that are fruiting. Only apply after flower petals have fallen off.

3.1 Resistance Management

THIAMETHOXAM	GROUP	4A	INSECTICIDE
CHLORANTRANILIPROLE	GROUP	28	INSECTICIDE

For resistance management, please note that Pitore Liquid Treatment contains both a Group 4A and Group 28 insecticide. Any insect population may contain individuals naturally resistant to Pitore Liquid Treatment and other Group 4A or Group 28 insecticides. The resistant individuals may dominate the insect population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists. Chlorantraniliprole belongs to the diamide class of chemistry.

Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

For additional information on Insecticide Resistance Management:

- Contact Syngenta at 1-866-SYNGENT(A) (1-866-796-4368)
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org>

Syngenta encourages responsible product stewardship to ensure effective long-term control of the insect pests on this label.

4.0 APPLICATION DIRECTIONS

4.1 Mixing Directions

Always use clean, well maintained application equipment. For best results, clean application tanks, hoses, and nozzles prior to mixing, especially if the tank has been used to apply a repellent insecticide or termiticide.

4.1.1 Dilution Preparation

Pitore Liquid Treatment must be mixed in water and applied as a dilute finished mixture using directions contained in the mixing table below.

Table 1. Amount of Pitore Liquid Treatment to Add to Spray Tank to Prepare Diluted Spray Solution

Dilution Concentration (% Active Ingredients)	Fluid Ounces (ml) of Pitore Liquid Treatment to Mix Listed Tank Volumes			
	1 Gallon	3 Gallons	5 Gallons	10 Gallons
0.15%	0.68 fl oz (20 ml)	2.0 fl oz (60 ml)	3.4 fl oz (100 ml)	6.8 fl oz (200 ml)

For small hand-pump sprayers (less than or equal to 5 gallons), mix thoroughly by shaking sprayer before and during treatment.

For large sprayers (greater than 5 gallons), mix Pitore Liquid Treatment in the following manner:

1. Fill applicator tank $\frac{1}{4}$ to $\frac{1}{2}$ full with water.
2. Begin agitation. For most systems, this means starting the pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose.
3. Before pouring into tank, shake the container of Pitore Liquid Treatment well.
4. Add required amount of Pitore Liquid Treatment according to **Table 1**.
5. Let pump run and allow recirculation through the hose.
6. Add the remaining amount of water.

Recirculate to complete mixing. The mixture should not be stored in the tank overnight. If this cannot be avoided, agitate the mixture thoroughly before using.

4.1.2 Foam Preparation

Foam applications to wall voids in stud walls should utilize dry foam only (e.g., 25:1 expansion ratio). Apply foam only to wall voids where termites or termite damage are detected or suspected.

Prepare a 0.15% dilution of Pitore Liquid Treatment and mix with the manufacturer's specified volume of foaming agent as follows:

- For wall galleries and spot applications, use an expansion ratio of 25:1
- For subsurface applications, concrete block, etc., use an expansion ratio in the range of 5:1 to 15:1

Use a foaming agent that is non-repellent to the target termite species. If sufficient foam volume cannot be applied to achieve the recommended rate of Pitore Liquid Treatment, apply an additional 0.15% dilution of Pitore Liquid Treatment as a liquid to assure proper treatment volumes and concentration in the treatment zone.

Table 2: Mixing Table for Pitore Liquid Treatment Foam

Pitore Liquid Treatment Use Dilution	Finished Mixture (Gallons)	Foam Expansion Ratio*	Finished Foam (Gallons)
0.15	1.0	25:1	25
	1.66	15:1	
	2.5	10:1	
	5.0	5:1	

* Add the manufacturer's specified quantity of foaming agent to the Pitore Liquid Treatment dilution.

5.0 RESTRICTIONS

5.1 Use Restrictions

- **DO NOT** apply this product in a way that will contact any other person. Keep people or pets away from treated area until dry.
- **DO NOT** apply as a broadcast spray in living areas. In living areas, use only as a void application, spot treatment, or crack-and-crevice application where occupants will not be exposed to direct contact.
- **DO NOT** treat soil that is frozen or water-saturated soil that will not accept the insecticide.
- **DO NOT** apply into electrical switches or receptacles or other uncoated wiring where electrical shock hazards exist.
- **DO NOT** contaminate public or private water supplies.

6.0 USE DIRECTIONS

6.1 Application Areas

6.1.1 Surface Treatment

Apply a 0.15% dilution of Pitore Liquid Treatment as a coarse spray at low pressure (<25 psi). Apply sufficient volume to cover the surface to the point of wetness, but avoid applying to the point of runoff. Alternatively, apply a 0.15% dilution of Pitore Liquid Treatment as a foam. Retreat as directed to maintain control. Allow treated surfaces to dry before contacting them.

For inaccessible surfaces, drill and treat the interior of structural voids using wood injections.

6.1.2 Wood Injection

For injection, use a pointed injection tip to apply a 0.15% dilution of Pitore Liquid Treatment as a liquid or foam to structural voids and insect galleries in damaged wood. Multiple injection points at varying depths may be necessary for best control. Spacing of the holes will depend on the distribution of insect activity and galleries. Injection holes may be clustered in areas with insect activity as indicated by damage, live insects or other detection methods. For control of drywood termites, refer to **Section 7.1** of this label.

Care should be taken to avoid electrical wiring, plumbing, etc. when drilling and injecting. Avoid drilling completely through wood.

7.0 SPECIFIC USE DIRECTIONS FOR REMEDIAL CONTROL OF LISTED PESTS

7.1 Drywood Termites

To control localized infestations of drywood termites, including species of *Incisitermes*, *Cryptotermes* and *Marginitermes*, apply a 0.15% dilution of Pitore Liquid Treatment as described below and in **Section 6.1.2** (Wood Injection). For foam-mixing instructions, refer to **Table 2** in **Section 4.1.2** (Foaming Instructions).

Locate galleries by using visual signs (e.g., fresh fecal pellets or blistered wood), the presence of live pests, mechanical sounding techniques (tapping on the wood surface and listening for changes in sound to indicate changes in wood density), listening devices, motion detection devices or other technologies that help pinpoint drywood termite activity.

To apply using the wood injection method, drill small diameter holes for the injection tip to intersect galleries with infested wood. Drywood termite emergence holes, or “pellet kick-out holes”, connect directly to galleries and are indicators of potential sites to drill. Apply up to 50 ml (1.7 fl oz) of Pitore Liquid Treatment at the 0.15% dilution at each injection hole. Holes must be plugged with an appropriate inert material after treatment.

Reapply if drywood termite activity within treatment areas is detected following treatment. For best results, inject a 0.15% dilution of Pitore Liquid Treatment into new injection holes drilled between previous injection sites.

7.2 Limited Area Treatment of Subterranean Termites in and Around Existing Structures

To control localized infestations of subterranean termites, apply a 0.15% dilution of Pitore Liquid Treatment as a coarse spray at low pressure (<25 psi) to the soil or the structure adjacent to the infestation. Applications should be made in sufficient volumes to coat the target surface, but less than the volume that creates runoff. Retreat as directed to maintain control. Allow treated surfaces to dry before contacting them.

This application is only intended for remedial control and is not intended as a preventive treatment. Use an approved termite control method for structural protection.

7.3 Wood-Destroying Beetles

For control of true powderpost beetles (*Lyctidae* species), death-watch beetles and ambrosia or bark beetles, apply Pitore Liquid Treatment to insect galleries, structural voids, spaces between wood elements of a structure, joints between wood and foundation, and/or as a surface treatment to exposed wooden surfaces in non-living space areas, such as crawlspaces, basements, and attics.

Apply a 0.15% dilution of Pitore Liquid Treatment as a coarse spray at low pressure (<25 psi) onto the surface in a volume to coat the target surface, but less than the volume that creates runoff. Alternatively, apply a 0.15% dilution of Pitore Liquid Treatment as a foam. Retreat as directed to maintain control. Allow treated surfaces to dry before contacting them.

8.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Pitore[®], the ALLIANCE FRAME
the Syngenta Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Viton[®] is a registered trademark of E.I. DuPont de Nemours & Company, Inc.

©20XX Syngenta

For non-emergency (e.g. current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

PITORE LIQUID TREATMENT XXXX NEW PRODUCT-E JUL2019-CL – di – 3/31/2020

000100-0XXXX.20190723E.PITORE-LIQUID-TREATMENT.NEW.JUL2019-CL