



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1765

Date of Issuance:

10/8/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A16148 ST

Name and Address of Registrant (include ZIP Code):

Patsy Laird
Regulatory Product Manager, Seedcare
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Kevin R. Ulrich
for

Kevin Ulrich (for)
Nathan Mellor, Chief
Fungicide Herbicide Branch, Registration Division (7505T)

Date:

10/8/25

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 100-1765."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 04/14/2025
- Alternate 1 CSF dated 04/14/2025

If you have any questions, please contact Hector Escobar at 202-566-1371 or at escobar.hector@epa.gov.

Enclosure: Stamped Label

SEDAXANE	GROUP	7	FUNGICIDE
----------	-------	---	-----------

A16148 ST
[Alternate Brand Name: Hyder]
Fungicide

A seed treatment product for protection against listed diseases on bulb onion crop subgroup 3-07A and Cucurbit vegetables crop group 9.

Active Ingredient:

Sedaxane*43.7%

Other Ingredients:56.3%

Total:100.0%

*CAS No. 874967-67-6

A16148 ST is a flowable concentrate for seed treatment containing 4.3 lb sedaxane per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-XXXX

EPA Est.

Net Contents




TABLE OF CONTENTS

1.0 FIRST AID

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

2.2 Personal Protective Equipment (PPE)

2.2.1 User Safety Requirements

2.2.2 Engineering Controls

2.2.3 User Safety Recommendations

2.3 Environmental Hazards

3.0 PRODUCT INFORMATION

3.1 Resistance Management

4.0 APPLICATION DIRECTIONS

4.1 Tank Mixtures

5.0 ROTATIONAL CROPS RESTRICTIONS

6.0 USE RESTRICTIONS AND PRECAUTIONS

6.1 Use Restrictions

7.0 SEED CONTAINER LABEL REQUIREMENTS

8.0 SEED TREATMENT DIRECTIONS

8.1 Bulb Onion Crop Subgroup 3-07A [– Not Registered for Use by California]

8.2 Cucurbit Vegetables Crop Group 9 [– Not Registered for Use by California]

9.0 STORAGE AND DISPOSAL

10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p align="center">SYNGENTA HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372</p>	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, causes moderate eye irritation, harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves including barrier laminate, butyl rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton™ ≥14 mils
- Shoes plus socks

2.2.1 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

2.2.3 User Safety Recommendations

Users Safety Recommendations

User should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

This product is toxic to fish. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notifications to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves including barrier laminate, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

3.0 PRODUCT INFORMATION

A16148 ST is a seed treatment fungicide which provides protection against listed soil-borne and seed-borne diseases of crop plants. A16148 ST is effective against seed and seedling blight or damping-off caused by listed seed- and soil-borne pathogens, including *Rhizoctonia* spp. Where rate ranges are shown, use the higher rate when disease pressure is expected to be severe.

3.1 Resistance Management

For resistance management, please note that A16148 ST contains a Group 7 fungicide, sedaxane. Any fungal population may contain individuals naturally resistant to A16148 ST and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

Sedaxane belongs to the carboxamide class of chemistry and is a succinate dehydrogenase inhibitor (SDHI). SDHI's inhibit fungal metabolism by binding to the succinate dehydrogenase enzyme, thereby disrupting cellular respiration and energy generation.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled.
- Adopt an integrated disease management program for fungicide use that includes scouting; uses historical information related to pesticide use and crop rotation; and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance, contact Syngenta Crop Protection at 1-866-796-4368. You can also contact your university extension specialist to report resistance.

4.0 APPLICATION DIRECTIONS

Apply A16148 ST as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of A16148 ST into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155(c). Allow seed to dry before bagging. Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).

A16148 ST does not control diseases caused by *Pythium* or *Phytophthora* spp. If these diseases are expected to be a problem, apply A16148 ST with other seed treatment products that contain mefenoxam as an active ingredient. If a mefenoxam-containing seed treatment product is not registered on that crop, apply A16148 ST as a tank mix with another labeled seed treatment product that has activity against these diseases.

4.1 Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. This product must not be mixed with any product which prohibits such mixing.

Bulb Onion Crop Subgroup 3-07A: For additional control of certain seed- and soil-borne pathogens, A16148 ST may be combined with seed treatment products containing mefenoxam, fludioxonil, picarbutrazox, and/or azoxystrobin. For insect control, A16148 ST may be tank mixed with a seed treatment product containing thiamethoxam, spinosad, or cyromazine.

Cucurbit Vegetables Crop Group 9: For additional control of certain seed- and soil-borne pathogens, A16148 ST may be combined with seed treatment products containing mefenoxam, fludioxonil, picarbutrazox, and/or azoxystrobin. For insect control, A16148 ST may be tank mixed with a seed treatment product containing thiamethoxam.

5.0 ROTATIONAL CROPS RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from seed treated with A16148 ST, crops may be replanted according to the following schedule:

ROTATIONAL CROP PLANT BACK INTERVALS				
Crop Planted	Immediate Plant Back	Minimum 30-Day Plant Back	Minimum 60-Day Plant Back	Minimum 12-Month Plant Back
Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet	Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Potato Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet	All other crops		
Potato	Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet		All other crops except potato	Potato

6.0 USE RESTRICTIONS AND PRECAUTIONS

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift.

6.1 Use Restrictions

- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feed and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, including in row ends.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. **DO NOT** contaminate water bodies when disposing of planting equipment washwater.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** apply more than 0.023 lb ai/A (10.4 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Bulb Onion Crop Subgroup 3-07A.
- **DO NOT** apply more than 0.0024 lb ai/A (1.1 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Cucurbit Vegetables Crop Group 9.

7.0 SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with sedaxane fungicide.
- **DO NOT** use for feed, food, or oil purposes.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with A16148 ST fungicide:

- The contents of this bag are for planting purposes only.
- **DO NOT** use for fuel or ethanol production purposes.
- Store away from feed and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, including in row ends.
- **DO NOT** plant treated seed by broadcasting to the soil surface. Ensure that all planted seeds are thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local

requirements. **DO NOT** contaminate water bodies when disposing of planting equipment washwater.

- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** apply more than 0.023 lb ai/A (10.4 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Bulb Onion Crop Subgroup 3-07A.
- **DO NOT** apply more than 0.0024 lb ai/A (1.1 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Cucurbit Vegetables Crop Group 9.
- In the event of crop failure or harvest of a crop grown from seed treated with A16148 ST, crops may be replanted according to the following schedule:

ROTATIONAL CROP PLANT BACK INTERVALS				
Crop Planted	Immediate Plant Back	Minimum 30-Day Plant Back	Minimum 60-Day Plant Back	Minimum 12-Month Plant Back
Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet	Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Potato Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet	All other crops		
Potato	Bulb Onion Crop Subgroup 3-07A Cereal Grains Crop Group 15 Corn (field, pop, sweet) Cotton Cucurbit Vegetables Crop Group 9 Legume Vegetables (Succulent or Dried) Crop Group 6 Peanut Rapeseed (including canola) Crop Subgroup 20A Sorghum Soybean Sugar Beet		All other crops except potato	Potato

8.0 SEED TREATMENT DIRECTIONS

8.1 Bulb Onion Crop Subgroup 3-07A [– Not Registered for Use by California]

Crop	Disease Controlled	Use Rate	
		fl oz/100 lb of seed	grams ai/100 kg of seed
Bulb Onion Crop Subgroup 3-07A Daylily, bulb Fritillaria, bulb Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Lily, bulb Onion, bulb Onion, Chinese, bulb Onion, pearl Onion, potato, bulb Shallot, bulb Cultivars, varieties, and/or hybrids of these	Seed Decay, Seedling Blight and Damping-off caused by <i>Rhizoctonia solani</i>	0.15 – 0.30	5 – 10
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.1. 			
USE RESTRICTIONS			
<ul style="list-style-type: none"> DO NOT apply more than 0.011 lb ai/A (5 g ai/A) per application of sedaxane-containing products as a seed treatment on Bulb Onion Crop Subgroup 3-07A. DO NOT apply more than 0.023 lb ai/A (10.4 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Bulb Onion Crop Subgroup 3-07A. A16148 ST-treated seed may be planted on the same acres up to 2 times per year. 			

8.2 Cucurbit Vegetables Crop Group 9 [– Not Registered for Use by California]

Crop	Disease Controlled	Use Rate	
		fl oz/100 lb of seed	grams ai/100 kg of seed
Cucurbit Vegetables Crop Group 9 Chayote (fruit) Chinese waxgourd (Chinese preserving melon) Citron melon Cucumber Gherkin Gourd, edible (includes hyotan, cucuzza, hechia, and Chinese okra) <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, and Chinese cucumber) Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Clause melon, and snake melon) Pumpkin Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini) Squash, winter (includes butternut squash, calabaza, Hubbard squash, acorn squash, and spaghetti squash) Watermelon	Seed Decay, Seedling Blight and Damping-off caused by <i>Rhizoctonia solani</i>	0.15 – 0.30	5 – 10
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.1. 			
USE RESTRICTIONS			
<ul style="list-style-type: none"> DO NOT apply more than 0.0012 lb ai/A (0.5 g ai/A) per application of sedaxane-containing products as a seed treatment on Cucurbit Vegetables Crop Group 9. DO NOT apply more than 0.0024 lb ai/A (1.1 g ai/A) per calendar year of sedaxane-containing products as a seed treatment on Cucurbit Vegetables Crop Group 9. A16148 ST-treated seed may be planted on the same acres up to 2 times per year. 			

9.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [(less than or equal to 5 gallons)]

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Handling [(greater than 5 gallons)]

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Handling [(greater than 5 gallons)]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC
©20XX Syngenta

<p>For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-866-796-4368.</p>
--

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

A16148 ST XXXX NEW-B 0425-CL – jeb – 09/18/2025
000100-0XXXX.20250429B.A161480466 ST.NEW-0425-CL.pdf