

INSECT DUST NO. DIRECTIONS

Follow directions carefully. Timing and method of application, weather and crop conditions, mix with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of material not in strict accordance with directions on herewith.

Sprinkle Rawleigh Insect Dust from a sifter-top (or better still use Rawleigh Insect Duster). Be sure to cover entire plant thoroughly and evenly—especially under leaves. Plants and fruits are not harmed by Rawleigh Insect Dust, nor does it injure the

Dust as soon as the insects appear. Select a time when the air is calm and preferably when foliage is damp from rain or dew.

For killing and controlling many sucking and chewing insects. For complete list see other side of package. Works by paralyzing insects upon contact when eaten. Make sure dust comes in actual contact with the insect.

Ready to use. This dust is ready to use as it comes from package. Repeat as necessary to control worms or insects which hatch out after the dusting. The number of applications depends on the number of insects and whether or not they are migrating from other fields. Dust again as soon as they reappear.

Will Not Damage Tender Foliage

POTENT—EFFICIENT
RELIABLE

*An Economical Contact and
Stomach Insecticide*

CAUTION

ready to use. This dust is ready
 as it comes from package.
 is necessary to control
 or insects which hatch out
 the first dusting. The number
 depends upon the
 of insects and whether or
 they are migrating from other
 . Dust again as soon as they re-
 ar.

**Economical Contact and
 Stomach Insecticide**

not damage tender foliage

**EFFICIENT
 D**

NOV - 2 1953

**Rotenone . . .
 Other Ether Extr
 a minimum of
 Sulfur . . .**

**Total . . .
 (Including:
 other inert**

Peas  Pea Aphid	Pea Aphid, 1/8 inch, pea green. One of the largest plant lice. Massed together in tops of stalks or underside. Pea Weevil, 1/8 to 1/5 inch. Dull colored with black and white markings.	Dust while dew is on. Dust when in blossom and as pods form to help control flies that deposit eggs. Dosage: 80 lbs. per acre.	yellow with black lines. Young that do damage are flesh colored, soft bodied. Potato Aphid, less than 1/8 inch, pink or green, long delicate wings and tail extending beyond body.	young bugs are seen, or eggs found on underside of leaves. Control same as all plant lice by dusting plants with Rawleigh Insect Dust. Dosage: 134 lbs. per acre.	Tobacco Flea Beetle, 1/16 to 1/12 inch. Jumps like flea when disturbed.	Water-melons Cucumbers Cantaloupes 	Striped Cucumber Beetle, 1/5 inch long, yellow above, black head and lengthwise stripes. Spotted Cucumber Beetle, 1/4 inch long, greenish-yellow. Head and legs black. 12 spots on back. Flea Beetle, see cabbage. Melon Lice. Melon Worm, white to light green, 3/4 to 1 inch.	Apply Rawleigh Insect Dust while dew is on, reaching underside of leaves as much as practical. Repeat as necessary. Dosage: 67 lbs. per acre.
Cabbage Cauliflower Broccoli Brussels Sprouts Lettuce Turnips Kohlrabi	Cabbage Worm, velvety green caterpillar, 1/2 to 1 1/2 inches. Cabbage Looper, pale green worm becoming striped, doubles up as it travels. Diamond-back, light green hairy caterpillar, 2/5 inch. Flea Beetles, small, dark colored bugs that jump like fleas. Feed on underside of leaves of radishes, turnips, eggplant, cabbage, tomatoes, potatoes.	Apply Rawleigh Insect Dust as soon as insects or eggs of the insect are found. Repeat in a week if insects continue numerous. Dosage: 67 lbs. per acre.	Tomato  Tomato Horn Worm  Moth Tomato Horn Worm	Young Tomato Horn Worm, fleshy, juicy green worms that devour leaves. Rear projection like a horn. Flea Beetles, eat small round holes in leaves. Jump like fleas when disturbed.	Sift & dusting of Rawleigh Insect Dust over the worms and the leaves where worms or beetles are seen. Dosage: 134 lbs. per acre.	Flowers Vines Shrubs	Aphis, or plant lice. Slugs Sowbugs Mites Ants	Thorough dusting of foliage & stems with Rawleigh Insect Dust gives good control of these parasites.

MANUFACTURED BY
THE W. T. RAWLEIGH COMPANY
 FREEPORT, ILLINOIS 61032

PETS

Cats
Dogs

Fleas

Dust into hair & into bedding
 Rawleigh Insect Dust