

DIRECTIONS

Place Rawleigh RAT and MOUSE KILLER in locations frequented by rats and mice and preferably where they are known to feed.

Bait should be protected from humans, livestock, poultry, or pets. This can be done by providing bait boxes which are accessible only to rats and mice.

FOR RATS

Bait should be used in amounts from 1/2 pound to 1 pound for each bait station and maintained for at least 10 days.


FOR MICE

The amount should be reduced to 1 tablespoonful, but more bait placements used.

Where continuous sources of infestation are present, such as from nearby dumps or fields, permanent feed stations should be established and kept refilled as needed. Be sure that feeding is not interrupted by lack of material.

Bait placements should be at 8- to 10-foot intervals and maintained at least 15 days. Average killing times are from 6 to 10 days, but some feeding may continue up to 4th week.

USDA REG. NO. 108-26



CAUTION

Pival, the active ingredient of this product, is an anti-coagulant chemical, therefore baits must be protected from, or placed where humans, livestock, poultry or pets will not accidentally eat them.

In case of eating, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIAN

Administration of blood transfusions combined with intravenous injections and oral doses of Vitamin K₁ is indicated.

R
RAWLEIGH

RAT AND MOUSE KILLER

ACTIVE INGREDIENT

2-Pivalyl-1, 3-Indandione* .025%

INERT INGREDIENTS 99.975%

Total 100.000%

*U.S. Patent No. Re. 23,080



NET WT. 32 OZ. (2 LBS.)

RAWLEIGH CO. FREEPORT, ILL. 61022