

[108-27]

DIRECTIONS

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of milk, milk equipment, food, dishes, utensils, and water. Avoid contamination of feed and feed products. Remove animals and poultry before treating. Contamination of fish-bearing waters will harm fish.

NOTE TO PHYSICIAN: Malathion upon repeated, prolonged, careless use may cause cholinesterase inhibition. Atropine is antidotal.

Directions for Use

HOGS: For the control of sarcoptic mange on hogs, mix 1 pint to 11 gallons of water. Apply as a complete coverage spray to all animals in the herd, bedding and walls. Use extreme care to thoroughly cover all body surfaces of the animal, including the inside of the ear. After spraying, the swine should be kept out of the sun and wind for a few hours. One thorough spraying will usually control sarcoptic mange of swine.

For the control of lice on swine, mix 1 pint to 11 gallons of water. Apply a complete coverage spray to animals, pens and litter. One treatment is usually sufficient.

BEEF CATTLE: For the control of lice on beef cattle, mix 1 pint to 11 gallons of water. Apply a complete coverage spray. One treatment is usually sufficient.

For the control of ticks and horn flies, mix 1 pint to 11 gallons of water. Apply a complete coverage spray. Repeat at two-week intervals if needed.

DO NOT APPLY TO LACTATING DAIRY ANIMALS OR NON LACTATING DAIRY ANIMALS WITHIN TWO WEEKS OF BREASTING.

DO NOT TREAT CALVES UNDER ONE MONTH OF AGE.

USDA REG. NO. 108-27



50%
MALATHION
Emulsifiable Concentrate

ACTIVE INGREDIENTS	
Malathion*	50%
Petroleum Hydrocarbons	39%
INERT INGREDIENTS	
	11%
Total 100%	

*O, O-dimethyl dithiophosphate
of diethyl mercaptosuccinate
ONE PINT LIQUID

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions
Manufactured by
THE W. T. RAWLEIGH COMPANY
Freeport, Illinois

DIRECTIONS CONTINUED

PEAS: Use ½ pint per 30 gallons of water for control of Pea Aphid, making one or more applications as needed.

BEANS: Use ½ pint per 30 gallons of water for control of Mexican Bean Beetle, Spider Mites, making two or more applications as needed.

CABBAGE: Use ½ pint to 1 pint per 50 gallons of water for control of Aphids, Imported Cabbage Worm, Cabbage Looper, making one or more applications as needed.

POTATOES: Use ½ pint per 50 gallons of water for control of Aphids, making one or more applications as needed.

NOTE: Make no application on peas, potatoes, beans, 3 days before harvest. Make no application on cabbage 7 days before harvest.

ORNAMENTALS: For the control of Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle adults, Four-spotted Leaf Bug, Tarnished Plant Bug, Rose Leaf Hopper and Potato Leaf Hopper, use 1 to 2 teaspoonfuls per gallon of water, making one or more applications as needed.

For control of: Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers, use ½ pint per 20 gallons of water, making one or more applications as needed.

For control of: Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes, use 1 pint to 50 gallons of water. Make one or more applications as needed, after crawlers have settled on the foliage.

For control of: Pine Needle Scale, use 1 pint per 25 gallons of water. Make one or more applications as needed, after crawlers have settled on the foliage.

NOTE: Do not use on Boston, Maiden Hair, Pteris Ferns as well as species of Crassula. Do not apply on Petunias.

[108-27]

