


# RAWLEIGH® Vapona® Insecticide Cattle & Barn Spray



DDVP for fast control of face flies, horn flies, stable flies, house flies, mosquitoes, gnats, on dairy animals, beef cattle, horses; around refuse areas, kennels

#### ACTIVE INGREDIENTS

*2,2-Dichlorovinyl Dimethyl Phosphate-0.930%	*Vapona® Insecticide — A Registered Trademark
*Related Compounds ..... 0.070%	**Equivalent to 0.0135% (butylcarbityl)
Pyrethrins ..... 0.002%	(6-propylpiperonyl) ether and 0.0034%
**Technical Piperonyl Butoxide ..... 0.017%	of related compounds
Petroleum Distillate ..... 98.981%	
Total ..... 100.000%	

FOR COMMERCIAL FARM USE. TO BE APPLIED BY TRAINED SERVICE PERSONS ONLY.

**FOR DAIRY CATTLE, BEEF ANIMALS:** To protect dairy and beef animals from face flies, horn flies, house flies, stable flies, mosquitoes and gnats, spray the animal daily using 1 to 2 ounces of this product per head. Cover the animal with spray with particular attention to back, flanks and legs.

Do not soak the skin or wet the hide.

If face flies are present, spray head and neck lightly.

Experience indicates that two applications of 1 ounce of the product morning and night give outstanding control of the above pests. This product may be applied as a mist spray, with a hand or automatic sprayer. Do not apply in excess of 2 fluid ounces per animal per day.

When a barn fogger is used, fog 1 to 1½ fluid ounces (½ pint = 8 fluid ounces or 1 cup) of this solution for each 10 running feet of barn length in average width barn. Let fog settle before milking. Do not fog this solution over animals that have had direct application of this solution within eight hours.

**SUPPLEMENTAL DIRECTIONS:** By diluting one part Rawleigh Vapona Insecticide—Cattle and Barn Spray with one part of fuel oil, diesel oil or kerosene, and applying as a coarse spray (sprinkling can), an insecticide contact and vaporizing spray is obtained to control flies, gnats, mosquitoes, roaches, ants, fleas, scabbugs and brown dog ticks.

In milk sheds, stables, livestock barns, loafing sheds, pig pens and outdoor areas, treat at the rate of one quart per 1000 square feet. Treat around doorways, feed storage rooms, alleyways, and stalls at the same rate. Apply this dilute solution as a coarse mist spray to contaminate milk or milk-handling equipment.

In poultry houses, apply to manure, window sills, outside of penned enclosures and on floor of feed storage rooms at the rate of one quart of dilute solution per 1000 square feet.

**DOG KENNELS:** In treating dog kennels, apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks, apply 2 quarts of this dilute solution per 1000 square feet of area.

For the control of fly maggots in maggot-breeding areas, such as garbage dumps and manure piles, apply one to two quarts of this dilute solution per 100 square feet of area. For the control of house flies, mosquitoes, gnats around picnic grounds, loading docks, outdoor latrines, parking areas, refuse areas, around service stations, open air drive-ins, ice cream stands, garbage collection and disposal areas, use at the rate of 1 quart of dilute solution per 1000 square feet.

**WARNING:** Do not get in eyes or on skin. Avoid breathing spray mist. Do not contaminate water, food or feedstuffs. May be fatal if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. If swallowed, induce vomiting immediately, and get medical attention. Aspirin is advised.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

Do not reuse this empty container. Destroy by perforating or burying in a safe place.

**WARNING — KEEP OUT OF REACH OF CHILDREN**  
THE W. T. RAWLEIGH COMPANY FREEPORT, ILLINOIS 61032  
KEEP CONTAINER TIGHTLY CLOSED WHEN NOT IN USE  
EPA REG. NO. 108-37-AA EPA EST. 108-IL-1

**NET CONTENTS 5 GALLONS**

63