



ACCEPTED
MAY 1 1984
108-37

RAWLEIGH® Cattle & Barn Spray

DDVP for fast control of face flies, horn flies, stable flies, house flies, mosquitoes, gnats, on dairy animals
and cattle, horses, around refuse areas, kennels

ACTIVE INGREDIENTS

| | | |
|--------------------------------------|----------|--|
| 2,2-Dichlorovinyl Dimethyl Phosphate | 93.0% | |
| Related Compounds | 0.070% | *Equivalent to 0.0135% (butylcarbityl) |
| Pyrethrins | 0.002% | (6-propylpiperonyl) ether and 0.0034% |
| *Technical Pyrethrins Butoxide | 0.017% | of related compounds |
| Petroleum Distillate | 98.981% | |
| Total | 100.000% | |

FOR COMMERCIAL FARM USE TO BE APPLIED BY TRAINED SERVICE PERSONS ONLY.

FOR DAIRY CATTLE, BEEF ANIMALS: This spray is intended for use on dairy cattle, beef animals, horses, and other animals in the home and barn. It is effective against face flies, horn flies, stable flies, house flies, mosquitoes, and gnats. It is also effective against ticks and lice. It is safe for use on animals and humans.

Directions: Spray the animals with the spray. Do not spray the face of the animal. Do not spray the eyes, ears, or mouth of the animal. Do not spray the animal's food or water. Do not spray the animal's bedding. Do not spray the animal's stall or stallion.

Supplemental Directions: This spray is also effective against ticks and lice. It is safe for use on animals and humans.

For Kennels: This spray is intended for use in kennels and other animal housing areas. It is effective against ticks and lice. It is safe for use on animals and humans.

Directions: Spray the kennel with the spray. Do not spray the animals. Do not spray the food or water. Do not spray the bedding. Do not spray the stallion.

Supplemental Directions: This spray is also effective against ticks and lice. It is safe for use on animals and humans.

THE W. T. RAWLEIGH COMPANY FREEPORT, ILLINOIS 61032 U.S.A.
EPA Reg. No. 108-37 AA EPA Est. 334 IL 1

NET CONTENTS 5 GALLONS

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

ACCEPTED

1984

108-37

RAWLEIGH

DIRECTIONS
IMPORTANT Child-Resistant Cap
TO OPEN Press down and turn

FOR DAIRY CATTLE, BEEF ANIMALS

To protect dairy and beef animals from face flies, horn flies, house flies, stable flies, mosquitoes and gnats, spray the animal, always using 1 to 2 ounces of this product per head. Cover the animal with spray with particular attention to back, flanks and legs.

Do not soak the skin or wet the hide

If face flies are present, spray head and neck lightly.

Expecting to use this spray on 100 head of dairy cattle of the above pests, this product may be diluted to a mist spray with a hand or electric sprayer. Do not spray in excess of 2 to 4 ounces per head per day.

When spraying, use a nozzle that will spray in a fan pattern. Do not spray directly into the eyes or ears. Do not spray directly into the mouth. Do not spray directly into the face. Let top settle before milking. Do not spray this solution on animals that are pregnant or lactating.

SUPPLEMENTAL DIRECTIONS

For control of eye gnats, spray the animal with Cattle and Barn Spray with long nozzle. Do not spray directly into the eyes. Use as a dust spray for control of eye gnats. Do not spray directly into the eyes. Do not spray directly into the mouth. Do not spray directly into the face.

For control of house flies, spray the animal with Cattle and Barn Spray with long nozzle. Do not spray directly into the eyes. Use as a dust spray for control of house flies. Do not spray directly into the eyes. Do not spray directly into the mouth. Do not spray directly into the face.

CATTLE AND BARN SPRAY

DDVP
FOR FAST CONTROL OF
• FACE FLIES • HOUSE FLIES
• STABLE FLIES • HORN FLIES
• MOSQUITOES • GNATS

ON DAIRY ANIMALS
• BEEF CATTLE • HORSES
• AROUND REFUSE AREAS
• OUTDOOR LIVING AREAS
• KENNELS

ACTIVE INGREDIENTS
DDVP 1.0%
P-PP-1 0.5%
P-PP-2 0.5%
P-PP-3 0.5%
P-PP-4 0.5%
P-PP-5 0.5%
P-PP-6 0.5%
P-PP-7 0.5%
P-PP-8 0.5%
P-PP-9 0.5%
P-PP-10 0.5%

KEEP CONTAINER TIGHTLY CLOSED WHEN NOT IN USE

In poultry houses, apply to manure outside of perched enclosures and on storage rooms at the rate of one quart of dilute solution per 1000 square feet.

DOG KENNELS.

In treating dog kennels, apply to outside dog house sides and eaves. For control of head lice, apply 2 quarts of this dilute solution per 1000 square feet.

For the control of fly maggots in indoor areas such as garbage dumps and manure piles, dilute this solution per 1000 square feet. For control of house flies, mosquitoes and gnats in areas such as porches, patios, refectories, dining rooms, outdoor, atrium, parking refectories, playgrounds, restrooms, open air, swimming stands, garbage collection and disposal at the rate of 1 quart of dilute solution per 1000 square feet.

WARNING: Do not get in eyes or on skin. If in eyes, flush with copious amounts of water. If on skin, wash with soap and water. Do not get on face. Wash thoroughly with water after spraying and before eating or drinking. Do not get on clothing. Do not get on food.

This product is toxic to fish and wildlife. Birds to which it is applied should be kept out of any water. Apply to areas as specified on this label. Do not use on waterfowl. Do not use on aquatic life.

EPA Reg. No. 108-37 AA EPA Est. 37A

THE W. T. RAWLEIGH Company

WARNING:
See this panel for additional precautions
KEEP OUT OF REACH OF CHILDREN

NET CONT. 1 GALLON