

ACCEPTED
 MAR 5 1971
 FEDERAL INSECT
 AND MAMMAL REPELLENT ACT



Contains Stabilized DDVP


units, apply to dropping area. Repeat as necessary.

Garbage Cans: Apply bait around and inside garbage can. Repeat as necessary.

For Reducing Populations of Resistant and Non-Resistant Houseflies In and Around Outdoor Living Areas, Picnic Grounds, Backyard Areas, Patios, Loading Docks, Outdoor Latrines and Parking Areas.

For Refuse Areas Around Service Stations, Open-Air Drive-Ins, Ice Cream Stands, Garbage Collection and Disposal Areas.

Use dry bait at 1/2 lb. per 1,000 sq. ft. of area. Apply on ground or areas where flies congregate. For open-air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.



dry

CAUTION: Not for Use Inside Homes Harmful If Swallowed, Inhaled or Absorbed Through the Skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not re-use container. Destroy when empty.

W. T. RAWLEIGH CO., FREEPORT, ILL. 61032

FOR FAST FLY CONTROL AROUND

- MILK SHEDS
- DAIRY BARNs
- HORSE BARNs
- STOCK BARNs
- PIG PENS
- KENNELs
- PATIOS
- OUTDOOR LIVING AREAs



VAPONA^(R) INSECTICIDE
 Dry Fly Bait

Use and Directions for Application

Dairy Barns, Milk Sheds, Stables, Live-stock Barns, Loafing Sheds, Pig Pens and Outdoor Areas: Scatter dry bait at the rate of 1/4 lb. per 1,000 sq. ft. on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly limed floors place treatment on feed sacks or on paper. Repeat procedure as necessary for practical fly control.

Cage Poultry and Turkeys on Wire: Scatter dry bait at the rate of 1/4 lb. per 100 sq. ft. on droppings beneath cages or perches. Repeat at weekly intervals.

Poultry Houses: Scatter dry bait at the rate of 1/4 lb. per 100 sq. ft. where flies congregate but keep out of reach of birds. Apply to manure, on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary.

Dog Kennels: Scatter dry bait at the rate of 1/4 lb. per 1,000 sq. ft. outside dog runways, on window sills, ledges and other surfaces out of reach of animals. For caged

ACTIVE INGREDIENTS:* BY WT.
 2,2-dichlorovinyl dimethyl phosphate - - - - 0.46%
 Related compounds - - - 0.04%
 INERT INGREDIENTS - - 99.50%
 Total - - - - 100.00%
 Keep container tightly closed when not in use
 *Equivalent to 0.5% of Vapona (R) Insecticide. A Registered Trademark.

Net Wt. 26 Oz. (1 Lb. 10 Oz.)



vapona^(R) insecticide

Contains Stabilized DDVP

General Use

For Residential Use: Use in Restrooms, Bathrooms, Kitchens, Hallways, Living Areas, Porch, Garages, Stairways, Patios, Loading Dock, and for Entrances and Parking Areas. For Retail Areas: Around Service Stations, Air Drive-In, Ice Cream Stand, and for Collection and Disposal Areas.

FOR FAST FLY CONTROL AROUND

- MILK SHEED
- DAIRY BARN
- HORSE BARN
- STOCK BARN
- PIG PENS
- KENNELS
- PATIOS
- OUTDOOR LIVING AREA



VAPONA^(R) INSECTICIDE

Dry Fly Bar

Use in Restrooms, Bathrooms, Kitchens, Hallways, Living Areas, Porch, Garages, Stairways, Patios, Loading Dock, and for Entrances and Parking Areas. For Retail Areas: Around Service Stations, Air Drive-In, Ice Cream Stand, and for Collection and Disposal Areas.

dry

For more information, call 1-800-368-7272