


PM 31

116-16

10/2



If swallowed, drink promptly a large quantity of egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.  
Store above 32° F (0° C)  
Do not reuse empty container.  
Open dumping is prohibited.

**Pesticide disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

This product fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 550 ppm of hardness calculated as CaCO<sub>3</sub> when tested by the AOAC Germicidal and Detergent Sanitizer—Official Method.

SOLVAY Animal Health, Inc.  
Mendota Heights, MN 55120 USA  
Manufactured at Charles City, IA 50616 USA  
1-800-894-9403

LG1895 PMS

ACCEPTED  
FEB 7 1995  
116-16



List No. 1895

# Germex

20% Quaternary Ammonium Compound

For agricultural use only

Disinfectant, sanitizer and deodorizer

Authorized by USDA for use in federally inspected meat, poultry and egg processing plants



Active ingredients:  
n-Alkyl (C14, 60%, C16, 30%, C12, 5%, C18, 5%)  
dimethylbenzylammonium chloride .. 20.0%  
Inert ingredients: Water ... 77.5%  
Isopropanol ... 2.5%

Keep out of reach of children

**Danger:**

See side panel for additional precautionary statements

EPA Registration No. 116-16  
EPA Est. 116-IA-1

**1 Gallon (128 fl oz/3.785 liters)**

SOLVAY Animal Health, Inc.

**BEST AVAILABLE COPY**

2 of 2



**Direction for Use:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Purpose	Dilution	Directions
<b>General disinfectant</b>		
Ordinary surface	1/2 cup (120 ml) per 15 U.S. gal. (57 liters) water or 1 tablespoon (15 ml) per 2 gal. (7.6 liters) (400 ppm)	For use in poultry and livestock buildings and equipment. Thoroughly clean all surfaces before spraying or brushing with Germex. If soap is used, rinse thoroughly before disinfecting. <b>Special directions: In poultry houses and livestock buildings:</b> 1. Remove all poultry or animals, all feedstuffs, trucks, coops, crates, etc., from the premises. 2. Empty all troughs, automatic feeders, and other feeding and watering equipment. 3. Remove droppings or manure from floors, walls and surfaces of facilities normally occupied or traversed by poultry or animals. 4. Remove all litter. 5. Saturate all surfaces, including equipment, with Germex solution. 6. Immerse in the solution all halters, ropes and other types of equipment used in handling animals. 7. Ventilate buildings, coops and other closed spaces. 8. Do not rehouse poultry or livestock, and do not reuse equipment, until the Germex solution has dried or set or been absorbed. 9. Rinse <u>thoroughly</u> with clean, drinkable water, all treated feed troughs, racks, waterers, mangers and other equipment before reuse.
Porous surface or area difficult to clean	3/4 cup (180 ml) per 10 U.S. gal. (38 liters) water (800 ppm)	

Purpose	Dilution	Directions
Farrowing swine	1 tablespoon (15 ml) per 4 U.S. gal. (15.2 liters) water (200 ppm)	Before farrowing, apply the lukewarm solution over all areas of udder and flanks, rinse thoroughly.
Dipping hatching eggs	1/2 cup (120 ml) per 30 U.S. gal. (114 liters) water or 1 tablespoon (15 ml) per 4 gal. (15.2 liters) (200 ppm)	To reduce surface contamination, dip eggs in solution at 110° F (43° C) to 120° F (49° C) before placing them in incubator.
Sanitizing meat- and poultry- processing utensils and equipment	1 cup (240 ml) per 60 U.S. gal. (228 liters) water or 1 tablespoon (15 ml) per 4 gal. (15.2 liters) (200 ppm)	Pre-cleanse utensils and equipment. Then, immerse utensils for at least 2 minutes or for contact time specified by governing sanitary code. Place sanitized utensils on a rack or drainboard to dry without rinse.

**Hazards to humans and domestic animals**

**Danger:** Keep out of reach of children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling product. Harmful or fatal if swallowed. Avoid contamination of food.

**Statement of Practical Treatment:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

4:59:48 AM

ACCEPTED

FEB 7 1995

176-16

**BEST AVAILABLE COPY**