

## FARM and HOME DISINFECTANT

## **ACTIVE INGREDIENTS:**

Coal Tar Phenols (Tar Acids)	14.0°°
Neutral Coal Tar Oils	61.6%
Anhydrous Soap	14.4°°
INERT INGREDIENTS	10.0°°
Phenol Cofficient A.O A.C Method	

Salmonella typhosa 6 minimum

**DANGER:** Keep out of reach of children. See back of panel for additional cautions and antidote statement.

division of rhodia inc

## Sec. 10. 1834

Best results will be obtained if warm water is used. Mix until solution becomes milky.

HOG HOUSES SHEEP BARNS DOG KENNIELS CATTLE BARNS HORSE STABLES YARDS and PENS

Remove all litter, bedding and dirt. Mix 1 pint of Hess & Clark Farm & Home Disinfectant with 4 gallons of water (1 gallon to 48 gallons of water). Spray or sprinkle all surfaces until thor oughly wet. This practice will aid in the control of lice fleas, ticks and infectious disease organisms.

MATAGER Do Not Use in Dairy Barns.

## AND MARKE OF POURRY HOUSES.

Remove all poultry and feed from premises, trucks coops or crates. Empty all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. For general disinfecting, mix 1 pint of Farm & Home Disinfectant with 6 gallons of water (1 gallon to 48 gallons water) and spray or sprinkle all surfaces thoroughly.

After treatment, ventilate buildings, coops and other closed spaces. Do not house poultry until treatment has been absorbed, set or dried.

FEEDERS WATERERS INCUBATORS BOOTS
EQUIPMENT AND UTENSILS

Remove all dirt and manure. Mix 1 pint Farm & Home Disinfectant with 4 gallons of water (1 gallon to 32 gallons of water). Thoroughly scrub equipment. All treated feed racks, mangers, troughs, equipment, automatic feeders, fountains and waterers must be thoroughly rinsed with potable water prior to reuse.

May be fatal if swallowed or absorbed through the skin. Call a physician immediately if taken internally. Do not get in eyes, on skin, on mucous membranes, or on clothing. May produce severe burns. For eyes, flush thoroughly with plenty water and immediately get medical attention. Wash thoroughly after handling and keep container closed when not in use. DO NOT BREATHE SPRAY MIST.

Internal, induce vomiting by running finger down throat. External, wash with soap and water.

Rinse empty container thoroughly with water and discard it.