

[134-57]

RESIDUAL FLY SPRAY 2EC can be applied by a knapsack or power sprayer.

RESIDUAL FLY SPRAY 2EC is a clear liquid. When added to water, it will disperse readily and will remain physically stable with a minimum amount of agitation.

**PREPARATION OF A 1% RESIDUAL FLY SPRAY 2EC**

**AMOUNT OF SPRAY DESIRED    AMOUNT OF RESIDUAL FLY SPRAY 2EC NEEDED**

1 PINT	4 TEASPOONS
2 QUARTS	1/4 CUP
6 QUARTS	1/2 PINT
3 GALLONS	1 PINT
6 GALLONS	1 QUART
12 1/2 GALLONS	2 QUARTS
25 GALLONS	1 GALLON

**DIRECTIONS FOR USE:** For control of house flies. When mixing, spray tank should be at least 1/2 full before adding Hess & Clark Residual Fly Spray 2EC. Stir to maintain a good suspension.

**RESIDUAL WALL SPRAYS:** For control of house flies, including resistant strains in dairy barns, calf barns, hog pens, poultry houses and other farm buildings. Apply a 1% spray. Thoroughly wet all fly resting areas to the point of runoff. Apply spray to ceilings, walls and floors. One gallon of spray will cover 500 to 1000 square feet of surface. Repeat application as necessary.

Remove all animals from area to be treated prior to spraying.

**SPOT SPRAYS:** For localized housefly control. Apply a 1% spray to areas where flies congregate. Repeat as necessary.

**MAGGOT SPRAYS:** For control of housefly maggots. Apply a 1% spray to suitable breeding areas such as poultry droppings, garbage dumps, and manure piles. Repeat applications as additional manure or garbage is added.

**GENERAL OUTDOOR USE:** For control of house flies around homes and recreation areas, garbage cans, animal quarters, food processing plants, garbage dumps, loading docks and warehouses. Apply a 1% spray to exposed areas. Repeat application as necessary.



EPA REG. NO. 134-57  
NET CONTENTS 1 GAL.

# RESIDUAL FLY SPRAY



**CONTROLS FLIES UP TO 8 WEEKS  
AND SHOULD BE APPLIED  
AS A 1% SPRAY.**

**KEEP OUT OF THE REACH  
OF CHILDREN. SEE SIDE PANEL FOR  
ADDITIONAL WARNINGS.**

Keep away from domestic animals. This product is toxic to wildlife. Birds or other wildlife feeding on treated areas may be killed.

**ACTIVE INGREDIENT:**


Dimethoate [O,O-Dimethyl  
S-(N-Methylcarbamoylmethyl)  
Phosphorodithioate] 23.0%

**INERT INGREDIENTS:** 76.6%  
**TOTAL** 100.0%

Manufactured by

division of rhodia inc.





EPA REG. NO. 134-57  
NET CONTENTS 1 GAL.

# RESIDUAL FLY SPRAY



*...residual ... fly ...*

**CONTROLS FLIES UP TO 8 WEEKS  
AND SHOULD BE APPLIED  
AS A 1% SPRAY.**

**ACTIVE INGREDIENT:**

Dimethoate [O,O-Dimethyl  
S-(N-Methylcarbamoylmethyl)  
Phosphorodithioate 23.4%  
(Equivalent to 2 lbs per gallon)

**INERT INGREDIENTS:** 76.6%  
**TOTAL** 100.0%

Manufactured by

division of rhodia inc.

**KEEP OUT OF THE REACH  
OF CHILDREN. SEE SIDE PANEL FOR  
ADDITIONAL WARNINGS.**

Keep away from domestic animals.  
This product is toxic to wildlife. Birds  
or other wildlife feeding on treated  
areas may be killed.

This product may cause irritation of eyes, nose,  
throat and skin. Harmful or fatal if inhaled, swal-  
lowed or absorbed through skin. Avoid breathing  
vapor or spray mist. Use with adequate ventilation.  
Vapor harmful, causes eye irritation. Avoid contact  
with eyes and skin. Wash thoroughly after handling.  
In case of contact with eyes, flush with plenty of  
water for at least 15 minutes; get medical attention.  
This product may cause  
cholinesterase inhibition. Antidote is Atropine.

Do not store below freezing temperatures.

Do not apply when weather conditions favor drift  
from area treated. Do not contaminate water by  
cleaning of equipment, or disposal of waste. Keep  
out of streams, lakes or ponds.

Food utensils such as teaspoons, tablespoons or  
measuring cups should not be used for food pur-  
poses after use with pesticides. Only apply this  
product with a knapsack or power sprayer.

Apply this product only as specified on this label.  
Do not reuse empty container. Destroy empty con-  
tainer by crushing or perforating and burying in  
a safe place.