

Vapor-Phase Surface Disinfectant containing 2-(Hydroxymethyl)-2-Nitro-1, 3-Propanediol Which Slowly Decomposes Yielding Formaldehyde

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE — KEEP UPRIGHT WHILE MOVING AND IN STORAGE. Container equipped with a pressure relief closure. KEEP FROM FREEZING. Store in original container only, in safe manner. Do not store under conditions which might adversely affect the container or its ability to function properly. Reduce stacking height where local conditions can affect package strength. Open container in a well ventilated area, avoid breathing vapor. Keep container tightly closed when not in use.

SPILL CONTROL — Soak up spill in a dry absorbant such as clay. Sweep it up, put into disposal containers and dispose as directed below. Wear protective clothing and equipment consistent with good pesticide handling procedures while exposed to spilled material.

PESTICIDE DISPOSAL — "Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance."

CONTAINER DISPOSAL — Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

DC&R[®] Disinfectant

ACCEPTED

OCT 15 1986

Active Ingredients
 2-(Hydroxymethyl)-2-Nitro-1, 3-Propanediol 19.20%
 Formaldehyde 2.28%
 Alkyl (C₁₂-67%, C₁₄-25%, C₁₆-7%, C₈, C₁₀, C₁₃-1%)
 dimethyl benzyl ammonium chloride 3.08%
 Inert Ingredients 75.44%
 Total 100.00%

EPA Reg. No. 134-65

"Only a hint of formaldehyde"

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED — Induce vomiting (if conscious). Call a physician.
 IF IN EYES — Flush immediately with water.
 IF ON SKIN — Wash thoroughly with soap and water.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes, on skin, or on clothing.
 Harmful if swallowed. Do not use on animals.

KEEP FROM FREEZING

Hess & Clark, Inc. Net Contents: 5 gal.
 ASHLAND, OHIO 44805

EPA Reg. No. 134-65
 EPA Est. No. 134-OH-1
 U.S. Pat. 4,008,332

DIRECTIONS FOR USE (Continued)

To be used in the disinfection of farm buildings and equipment which are used for poultry and livestock production. Especially suited for equipment which may harbor or spread many disease germs within the farm operation or to other farms. Dilute DC&R Disinfectant one ounce per gallon of water. Use as follows:

EQUIPMENT

Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Then thoroughly wet all exposed parts of the cleaned equipment used in the poultry or livestock operation.

All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

Immerse cleaned halters, ropes and other types of equipment used in handling and restraining animals; forks, all shovels and scrapers used for removing litter and manure and allow DC&R Disinfectant to saturate all surfaces for 10 minutes. Do not spray onto feed to be consumed by animals.

When disinfecting trucks used for hauling poultry or livestock, it is desirable to rinse the truck cab a few minutes after the spraying is completed. Do not rinse other parts of the truck. The release of formaldehyde is so slight that little or no odor can be detected. However, allow reasonable ventilation for animals transported following disinfection.

BUILDINGS

Evacuate all animals from buildings. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Wet all surfaces by allowing DC&R Disinfectant to saturate all surfaces for 10 minutes. Equipment inside the building (cages for laying hens, for example) should be first cleaned with soap or detergent, rinsed with water and then saturated as above with this product.

Ventilate buildings, cars, boats, coops and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

