# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

APR 2,5 200

SEPA Office of Pesticide Programs

Hess & Clark, Inc. 110 Hopkins Drive Randolph, WI 53956

Attention: Jennifer J. Seifert

Subject: DC&R Disinfectant EPA Registration No. 134-65 Notification Dated March 27, 2007

This will acknowledge receipt of your notification for an alternate warranty statement, submitted under the provisions of FIFRA Section 3(c)(9). Based on a review of the submitted material, the following comments apply.

The Notification is in compliance with PR Notice 98-10 and is acceptable. This information has been added to your files.

If you have any questions concerning this letter, please contact Martha Terry at (703) 308-6217.

Sincerely

Marshall Swindell Product Manager (33)

Regulatory Management Branch 1 Antimerobials Division (7510P)

≎EPA	Environmental I	red States Protection A ton, DC 20460	gency	X	Registr Amend Other		OPP Identifier Number
	Α	pplication fo	r Pesticid	e - Sectio	n I		
1. Company/Product Number	r		!	oduct Manage	Г	3. Pr	oposed Classification
134-65		<del></del>	Marshall	Swindell			None Restricted
4. Company/Product (Name) DC&R Disinfectant			PM#				
5. Name and Address of App Hess & Clark, Inc. 110 Hopkins Drive Randolph, WI 53956	plicant (Include ZIP Code  t is a new address	,	(b)(i), my to: EPA Re	product is s	imilar or iden	itical in co	FIFRA Section 3(c)(3) mposition and labeling
		S	ection - II	t Name			
: 1			1 1			<u> </u>	<u> </u>
Amendment - Explain			<u> </u>	final printed lal Agency letter d	•	se to	
Resubmission in resp	onse to Agency letter da	ted	[ ] •	'Me Too" Appl	ication.		
X Notification - Explain	below.			Other - Explain	below.		
Explanation: Use addition	nel negate) if necessary	(For section 1 and	Section II )				
Pursuant to PR-Notice-98-10, in regard to the subject product, we are hereby notifying the Agency of an alternate warranty statement.							
		Se	ction - III				<u> </u>
1. Material This Product Will	Be Packaged In:						
Child-Resistant Packaging	Unit Packaging	Wat	ter Soluble Pad	kaging	2. Type of	Centainer	
Yes	Yes	<u> </u>	Yes		1	_ Metal Plastic	
No	No		J No			Glass	
* Certification must be submitted		No. per If "' container Bec	Yes Kage wgt	No. per container		Paper Other (S	pecify)
3. Location of Net Contents (	Information 4.	Size(s) Retail Con	tainer 1	5. (	ocation of La	bul Direction	ns .
Label C	gatainer			F		-	
6. Manner in Which Lebel is	Affixed to Product	Lithograph Paper glued Stenciled		Other _			
	<u></u>		ction - IV			<del></del>	
1. Contact Point /Complete	items directly below for		<del></del>	ontacted, if ne	cessary, to pi	ocess this	aprécation.)
Name Jennifer J. Seifert		Title	ager, Regula			r	No. (include Area Code)
	nents I have made on thi y knowlingliy false or mis aw.						6. Date Application Received (Stamped)
2. Signature 3. Title							
Jennilo I	Put)	Mar	nager, R	egulator	/ Affairs		
4 Typed Name	/	5. Date	1				
Jennifer J. Seifert		ن ا	3/27/07				

375

Manufacturing 110 Hookins Drive Randolph, WI 53956 920/326-5141 Fax: 920/326-5135 inform@neogen.com www.neogen.com

8601 1998 6568

Fed Ex:

Neogen Corporation 944 Nandino Boulevard Lexington, KY 40511 800/477-8201 859/255-1221 Fax: 859/255-5532 inform@neogen.com www.neogen.com

# HESS&CLARK, INC.

28 March 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

RE:

DC&R® Disinfectant

EPA Reg. No. 134-65

Gentlemen/Ladies:

Pursuant to PR-Notice 98-10, in regard to the subject product, we are hereby notifying the Agency of an alternate warranty statement for the following product: DC&R® Disinfectant.

Following is my statement as required by PR-Notice 98-10:

This notification is consistent with the provisions of PR-Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with terms of PR-Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Attached please find the completed application form marked for Notification, as well as a "before" and "after" label, with the additional warranty statement clearly marked.

If you should have any questions regarding this matter, please contact me at 920-326-2461.

Sincerely,

Hess & Clark, Inc.

Jennifer J. Seifer

Manager, Regulatory Affairs

Enclosure(s): 3



# Disinfectant

# Bactericidal • Fungicidal • Virucidal\*

For Control of Canine Parvovirus and Feline Panleukopenia in Veterinary Clinics and Kennels

ACTIVE INGREDIENTS: 2-(Hydroxymethyl)-2-Nitro-1, 3-Propanediol	19.20%
Alkyl (C12-67%, C14-25%, C16-7%, C8, C10, C18-1%)	)
dimethyl benzyl ammonium chloride	3.08%
Formaldehyde	
INERT INGREDIENTS:	<u>75.44%</u>
TOTAL:	100.00%
EPA Reg. No 134-65	EPA EST NO 61282-WI-1

# KEEP OUT OF REACH OF CHILDREN DANGER

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-800-498-5743

### FIRST AID

A CONTROL THE CONTROL TO STREET

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.  Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye Contact poison control center or doctor for treatment advice.
If on skin or on clothing	Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably month-to-month, if possible. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison centrel center or doctor immediately for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For a Medical Emergency involving this product, call 1-800-498-5743

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage or skin burns. Hamful if absorbed through skin or inhaled. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Avoid breathing vapor or spray mist. Prolonged or repeated skin contact may cause allergic reaction in some individuals. PERSONAL PROTECTIVE EQUIPMENT (FFE).

Pesticide handlers must wear Long-sleeved shirt and long pants, chemical resistant gloves (such as any waterproof material, rubber, etc.), wear protective eyewear (goggles, face shield, or safety glasses), shoes plus socks. In addition, when engaged in pouring this product, a respirator with either an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter

#### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash separately from other laundry. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the totlet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Worsh the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.



#### ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuanes, oceans or public waters unless in accordance with the requirements of a National Polihitant Discharge Elimination Systems (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer treatment plant authority. For guidance confact your local State Water Board or Regional Office of EPA.

GENERAL: DC&R Disinfectant is a non-flammable and non-corrosive formulation. DC&R Disinfectant can be used in the following areas. Poultry and Turkey Farms, Hatcheries, Swine Operations, Dairy Farms, Cattle Farms, Veal Farms, Equine Farms, Emm and Ostrich Farms, Zoos, Vetermary Clinics, Kennels, Pet Shops, Federally Inspected Meat and Poultry Establishments except where meat and food products are handled, Asumal Life Sciences Laboratories and in Farm Foot Baths. DC&R Disinfectant can be used to disinfect pens, hutches, feeders, waterers, cages and livestock and poultry hauling velucles and equipment.

DC&R has proven effective as a disinfectant by the AOAC use dilution test modified for spray products against the following organisms. All dilutions 1 128 tested at 400 ppm water hardness (as calcium carbonate) and 5% calf serum

Mycoplasma pneumoma Salmonella cholenaessus Salmonella typhimizimi Salmonella enteritidis Pasteurella multocida Clostindium perfungens Eschenchia coli Campylobacter jojum Klabsiella pneumonae Enterobacter aerogenes Proteus vulgaris Streptococcus pyogenes Bordetella bronchiseptica Actinobacillus suis

\*DC&R has proven virucidal efficacy for use on hard, nonporous, inanimate environmental surfaces. All dilutions 1:128 tested with 400 ppm water hardness (as calcium carbonate) and 5% boying scrum.

Pseudorabies Neweastle disease Avian laryngotrachettis
Transmissible gastroenteritis Avian influenza Swine influenza virus
Canine purvovirus\*\* Feline panleukopenus\*\* Infectious bronclutus virus
Porcine Respiratory and Reproductive Syndrome (PRRS) Infectious Bursal Disease Virus

DC&R has proven effective sungecidal activity by the AO.AC use dilution test modified for spray products. All dilutions 1 128 tested with up water and 5% call serium

Candida albicans Trychophyton mentagrophytes

#### DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

To be used for disinfecting hard, non-porous surfaces of farm buildings and equipment which are used for poultry and livestock production and veterinary clinics, kennels and other non-livestock buildings. Especially suited for equipment which may harbor or spread many germs or microorganisms within the farm operation, veterinary facility or to other farms. Apply DC&R Disinfection with a mechanical spray device. Spray all surfaces to be disinfected until thoroughly wet Dilute DC&R Disinfectant one ounce per gallon of water. Use as follows:

Poultry and Livestock Buildings: Evacuate all arumals from buildings. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chures and other factifities and fixtures occupied or traversed by animals Empty all troughs, tacks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and mise with water. Spray all surfaces and allow DC&R Disinfectant to saturate all surfaces for 10 minutes Equipment inside the building (cages for laying heris, for example) should be first cleaned with soap or detergent, rinsed with water and then saturated as above with this product.

Ventilate buildings, coops and other closed spaces. Do not house poultry, investock or employ equipment until treatment has been absorbed, set or dired, then thoroughly scrub all treated feed issubs, troughs, automatic feeders, fountiaris and waterers with soap or detergent and runse with potable water before reuse.

Equipment: Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all equipment surfaces (cages for laying hens, for example) with soap or detergent and rinse with water. Then thoroughly spray or soak all exposed parts of the cleaned equipment used in the poulity, investock, or veterinary operation.

All treated feed racks, food bowls, mangers, troughs, automatic feeders, fountains and waterers must then be thoroughly combined again with delargants and missed with possible water prior to reuse.

Immerse cleaned halters, ropes and other types of equipment used in handling and restraining animals, forks, and all shovels and accepts used for removing litter and manure or feces and allow DC&R Disinfectant to saturate all surfaces for 10 minutes. Do not spray onto feed to be consumed by animals.

FOGGING: DC&R Disinfectant can be used in fogging (wet mist) operations as an adjunct either proceding or following regular cleaning and disinfecting procedures. DC&R at 1 to 2% is appropriate A 1% to 2% meeting is achieved by mixing 1-1/4 to 2-1/2 ounces of DC&R Disinfectant to 1 gallon (128 ounces) of water. Fog (wet mist) until the area is moist using automatic wet fogger according to manufacturer's use directions.

Reentry Restrictions: Entry by any person-other than a correctly equipped handler is PROHIBITED in the entire enclosed building/structure from the start of application until aeration reduces the air concentration level of formaldehyde in the working area to less than 0.75 ppm. The air concentration of formaldehyde must be measured before reentry is permitted (OSHA issued a final rule for the PL of formaldehyde as 0.75 ppm, May 27, 1992, Federal Register, Vol. 57, p. 22290). Any handler who enters the treated area during the entry-restricted period must wear a respirator with either an organic vapor removing cartridge with a prefilter approved for pesticidos (MSHA/MOSH approval number prefix TC-14G), or a NIOSH approval respirator with an organic vapor (OV) cartridge or canister with any N. R. P or HE prefilter.

\*\*FOR CANINE PARVOVIRUS & FELI Parvovirus and Fehne Panleukopenia when and Feline Panleukopenia thoroughly clean i of water. Apply with a cloth, mop, sponge, minutes Except for the change in dose, th cleaning and second water inse directions lis ALL OTHER NON-LIVESTOCK BULL Pet Shops: Thoroughly clean all surfaces disinfected and allow to saturate all surfaces gallon of water, unless the objective is to cot directions specific to that use. Ventilate build

## STORAGE AND DIS

Do not contaminate water, food or feed by st STORAGE: KEEP UPRIGHT WHILE M closure KEEP FROM FREEZING. Store is which might adversely affect the container conditions can affect package strength. Open highly closed when not in use.

SPILL CONTROL: Soak up spill in a dr. dispose as directed below. Wear protective of while exposed to spilled material.

PESTICIDE DISPOSAL: "Pesticida waste is a violation of Federal Law If these was State Pesticide or Environmental Control. Office for guidance"

CONTAINER DISPOSAL: Triple ruse dispose of in a sanitary landfill, or momen out of smoke

#### CONDITIONS OF SALI


NOTICE: Read the entire Directions for Use using this product. If the terms are not accepta

Manufacturer and Seller warrant that this two purposes stated in the Directions for Use, sub under normal use conditions. This warranty abnormal condutions or under conditions not r and User assume the risk of any such use. MANUFACTURER AND SELLER MAK PARTICULAR PURPOSE NOR ANY OTH in no event shall Manufacturer or Seiler be a handling of mis product. THE EXCLLSIVE OF MANUFACTURER AND SELLER (INCLUDING CLAIMS BASED ON BE LIABILITY OR OTHERWISE) RESULTE-RETURN OF THE PURCHASE PRICE OF SELLER, THE REPLACEMENT OF THE P Manufacturer and Seller offer this product, Limitations of Warranty and Liability, which representative of Manufacturer

Manufactured by Hess & Clark, Inc. 110 Hopkins Drive Randolph, WI 53956



<sup>\*\*</sup>See specific instructions for Canine parvovirus and Feline panleukopenia





## Virucidal\*

Vetermary Chrice and Kennels

(\$4,6)4*!***!!*!   * * *	19,209
*******************	3.08%
	2.28%
	75.44%
**************	100.00%

EPA EST NO 61282-WI-1

# CHILDREN

DDUCT, CALL 1-800-498-5743

th water for 15-20 minutes at 5 minutes, then continue rinsing or treatment advice imediately with plenty of water for doctor for treatment advice thing, call 911 or an ambulance, (th-to-mouth, if possible Call a divice.

Do not induce venting unless

when Do not give anything by

e the product container or label with you

ill 1-800-498-5743

#### ENTS

ill if absorbed through skin or inhaled Do s, face shield, or safety glasses). Avoid illergic reaction in some individuals

resistant gloves (such as any waterproof ty glasses), shoes plus socks. In addition, vapor removing cartridge with a prefilter or a canister approved for pesticides 3 respirator with an organic vapor (OV)

I no such instructions are available for tlaundry. Users should wash hands before d remove clething immediately if pesticite move PPE immediately after handling this la, wash thoroughly and change into clean

#### FNVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuanes, oceans or public waters unless in accordance with the requirements of a National Pollutant Discharge Elimination Systems (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer treatment plant authority. For guidance contact your local State Water Board or Regional Office of EPA.

GENERAL: DC&R Disinfectant is a non-flammable and non-corresive formulation. DC&R Disinfectant can be used in the following areas. Poultry and Turkey Farms. Hatcheries, Swine Operations, Dairy Farms, Cattle Farms, Veal Farms, Equine Farms, Emu and Ostrich Farms, Zoos, Veterinary Clinics, Kennels, Pet Shops, Federally Inspected Meat and Poultry Establishments except where meat and food products are handled, Animal Life Sciences Laboratories and in Farm Foot Baths. DC&R Disinfectant can be used to disinfect pens, butches, feeders, waterers, cages and beestock and poultry hauling vehicles and equipment.

DC&R has proven effective as a disinfectant by the A O.A.C. use dilution test modified for spray products against the following organisms. All dilutions 1, 128 tested at 400 ppm water hardness (as calcium carbonate) and 5% calf serum

Mycoplasma pneumonia	Salmonella choleraesius	Salmonella typhimuriur
Salmonella enteritidis	Pasteurella multocida	Clostridium perfringens
Eschenchia coli	Campylobacter jejuni	Klebsiella pneumoniae
Enterobacter aerogenes	Proteus vulgaris	Streptococcus pyogenes
Bordetella bronchiseptica	Actnobacillus suis	

\*DC&R has proven virue;dal efficacy for use on hard, nonporous, inanimate environmental surfaces. All dilutions I 128 tested with 400 ppm water hardness (as calcium carbonate) and 5% bovine serum.

	Pseudorabies	Newcastle disease	Avian laryngotracheitis	
Transmissible gastroenteritis Canine parvovirus**		Avian influenza	Swine influenza virus Infectious bronchitis virus	
		Feline panleukopenia**		
	Porcine Respiratory and Reproductive	Infectious Bursal Disease Virus		

<sup>\*\*</sup>See specific instructions for Canine parvovirus and Feline parleukopenia.

DC&R has proven effective fungioidal activity by the AOAC, use dilution test medified for spray products. All dilutions 1:128 tested with tap water and 5% calf scrum

Candida albicaris Trychophyton mentagrophytes

#### DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

To be used for disinfecting hard, non-porous surfaces of farm buildings and equipment which are used for poulitry and livestock production and veterinary clinics, kennels and other non-livestock buildings. Especially suited for equipment, which may harbor or spread many germs or microorganisms within the farm operation, veterinary facility or to other farms. Apply DC&R Disinfectant with a mechanical spray device. Spray all surfaces to be disinfected until thoroughly wet Dilute DC&R Disinfectant one ounce per gallon of water. Use as follows:

Poultry and Livestock Buildings: Evacuate all animals from buildings. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, clutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Throughly, team all surfaces with scop or detergent and rinse with water. Spray all surfaces and allow DC&R Disinfectant to saturate all surfaces for 10 minutes Equipment inside the building (cages for laying herss, for example) should be first cleaned with scap or detergent, rinsed with water and then saturated as above with this product.

Ventilate buildings, coops and other closed spaces Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried, then thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

Equipment: Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all equipment surfaces (cages for laying heas, for example) with soap or detergent and raise with water. Then thoroughly spray or soak all exceeded acts of the cleaned equipment used in the roughly, invative, or exterious operation.

seak all exposed parts of the cleaned equipment used in the poultry, livestock, or veterinary operation.

All treated feed racks, food bowls, mangers, troughs, automatic feeders, fountains and waterers must then be

thoroughly scrubbed again with detergents and nased with potable water prior to reuse.

Immerse cleaned halters, ropes and other types of equipment used in handling and restraining enimals, forks, and all shovels and scrapers used for removing litter and manure or feces and allow DC&R Disinfectant to saturate all surfaces for 10 minutes. Do not spray onto feed to be consumed by animals.

FOGGING: DC&R Disinfectant can be used in fogging (wet mist) operations as an adjunct either preceding or following regular cleaning and disinfecting procedures. DC&R at 1 to 2% is appropriate. A 1% to 2% maxture is achieved by mixing 1-1/4 to 2-1/2 ounces of DC&R Disinfectant to 1 gallon (128 ounces) of water Fog (wet mist) until the area is most using automatic wet fogger according to manufacturer's use directions.

Reentry Restrictions: Entry by any person-other than a correctly equipped handler is PROHIBITED in the entire enclosed building structure from the start of application until aeration reduces the air concentration level of formaldehyde in the working area to less than 0.75 ppm. The air concentration of formaldehyde must be measured before recentry is permitted. (OSHA issued a final rule for the PL of formaldehyde as 0.75 ppm, May 27, 1992, Federal Register, Vol. 57, p. 22290). Any handler who enters the treated area during the entry-restricted period must wear a respirator with either an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approval respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter

\*\*FOR CANINE PARVOVIRUS & FELINE PANL FUKOPENIA: DC&R Disinfectant is effective against Canine Parvovirus and Feline Pardeukopenia when used at a three curiese per gallon dilution. To centrel Canine Parvovirus and Feline Parleukopenia theroughly clean all surfaces and mix three (3) ounces of DC&R Distinfectant in one gallen of water. Apply with a cloth, mop, sponge, as a spray or by soaking. Allow to air dry or area may be mixed after 45 minutes. Except for the change in dose, the identical animal removal, pre-cleaning, water mixe, treatment, second cleaning and second water rinse directions lated in the building and entitiment sections above, must be followed.

ALL OTHER NON-LIVESTOCK BUILDINGS AND FACILITIES including Veterlaary Clinics, Kennels, and Pet Shops: Thoroughly clean all surfaces with soap or detergent and time with water. Spray all surfaces to be disinfected and allow to saturate all surfaces for 10 mmutes. The use rate is one (1) ounce of DC&R Disinfectant per gallon of water, unless the objective is to control Canne Parvovius or Peline Panleukopenia, then you must follow the directions specific to that use Vernilate braidings and other thosed spaces.

# STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

STORAGE: KEEP UPRIGHT WHILE MOVING AND IN STORAGE Container equipped with a pressure relief closure. KEEP FROM PREEZING Store in original container only, in safe manner Do not store under conditions which might adversely affect the container or its ability to function properly. Reduce stacking height where local conditions can affect package strength. Open container in a well-ventilated area, avoid breathing vapor. Keep container tightly closed when not in use

SPILL CONTROL: Soak up spill in a dry absorbent such as clay Sweep it up, put into disposal containers and dispose as directed below. Wear protective elections and equipment consistent with good pesticide handling procedures while exposed to spilled material.

PESTICIDE DISPOSAL: "Pesticide wastes are toxic Improper disposal of excess pesticide, spray mixture, or misate is a violation of Federal Law. If these wastes carniet be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazzirdous Waste representatives at the nearest FPA Regional Office for guidance."

CONTAINER DISPOSAL: Imple runse (or equivalent) Then offer for recycling or reconditioning, or junctum and dispose of in a sentiary landfill, or incineration, or, if allowed by State and Local authorities, by burning if burned, stay out of enote.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product if the terms are not acceptable, report the product at once, unopened, and the purchase prior will be refunded.

Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use

MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FIRNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY FXCEPT AS STATED ABOVE In no event shall Manufacture or Seller be liable for any incidental, consequented or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE HABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAMIS, LOSSES, INJURIES OR DAMAGES (INCILIDING CLAMIS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISD, RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Nide and Limitations of Westernty and Liability, which may not be modified except by written agreement signed by a drift authorized expressionable of Manufacturer.

Manufactured by: Hess & Clark, Inc. 110 Hopkins Drive Randolph, WI 53956

