

DE-PESTER®

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

SOIL INSECTS ATTACKING CORN, BEETS, BEANS, BRUSSELS SPROUTS AND CABBAGE

Unless otherwise specified, rates are for dilution with sufficient water to cover one acre.

INSECT	RATE OF APPLICATION	METHOD OF APPLICATION
Ants	1 1/2 pounds per 3 gallons of water (2 1/2 quarts)	Thoroughly wet soil or ornamental turf at first sign of infestation.
Wireworms	6 1/2 pounds	Work into soil to 4" depth at planting time.
White Grubs	12 1/2 pounds	Apply to soil and turf in Spring.
White Fringed Beetle Larva	6 1/2 pounds	Work into soil to 4" depth at planting time.
Asiatic Garden Beetle	25 pounds	Work into soil to 4" depth at planting time.
Cabbage Maggots	2 1/2 pounds	Work into soil to 4" depth at planting time.
Chinch Bugs	2 1/2 pounds	Apply to soil surface at first signs of infestation.
Onion Maggots	2 1/2 pounds	Apply when seeding.
Cutworms	2 1/2 pounds	Apply to soil surface around plants.
Corn Rootworm	2 1/2 pounds	Apply when seeding as row treatment.
Seed Corn Maggot	2 1/2 pounds	Apply when seeding.

Apply as pre-plant, planting or transplanting application.

RESIDUAL CONTROL

Prepare a spray by diluting 1 pound **Chlordane W-40** with 2 1/2 gallons of water. Apply this spray as a spot treatment to infested areas such as cracks, crevices, the undersides of carpets and infested areas for the control of **carpet beetles, cockroaches, scorpions, silverfish, spiders (black widow included)** in homes.

GARDEN AND TRUCK CROPS

INSECT	CROP	APPLICATION
Tomato Hornworm	Tomatoes	2 1/2 pounds in 100 gal. water per acre.
Onion Thrips	Onions	2 1/2 pounds in 100 gal. water per acre with 100 gal. water repeated if needed.
Cabbage Looper	Cabbage	2 1/2 pounds in 100 gal. water per acre.
Cabbage Worms	Cabbage	2 1/2 pounds in 100 gal. water per acre.
Imported Cabbage Worm	Cabbage	2 1/2 pounds in 100 gal. water per acre.
Marquim Cabbage Bug	Cabbage	2 1/2 pounds in 100 gal. water per acre.
Colorado Potato Beetle	Potatoes	2 1/2 pounds in 100 gal. water per acre.
Beet Webworm	Beets	2 1/2 pounds in sufficient water to cover 1/2 acre. Apply early before damage is severe.

*Do not apply to cabbage after edible parts begin to form. Do not apply to tomatoes after fruit begins to form. Do not apply to green or spring onions. No application should be made to beets after seeding stage if tops are to be used for food. Keep children and pets off treated lawns until material is watered into soil and grass is dry.



Chlordane W-40 contains 40% active ingredient.

CAUTION. Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid contamination of feed and foodstuffs. Avoid breathing spray mist. Keep out of reach of children. Do not apply to dairy animals, calves or in dairy barns. Do not use on feed or forage crops to be fed to dairy animals or those being finished for slaughter. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residue in excess of official tolerances.

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds.

NOTICE

For more information on the use of this product, contact your local agricultural agent or the manufacturer, T. H. Green, Inc., 1000 North 1st Street, St. Paul, Minnesota 55101.



CAUTION Keep Out of Reach of Children
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

CONTAINS 40% CHLORDANE

**WETTABLE POWDER FOR YARD,
GARDEN AND FARM USE**

ACTIVE INGREDIENT:

Technical Chlordane 40%

INERT INGREDIENTS 60%

100%

(Contains 24% Octachloro-4,7-methano tetrahydroindane and 16% related compounds)

USDA Reg. No. 148-47

2-69

