

ACCEPTED
 12/14/73
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND NAUGHTICIDE ACT
 FOR ECONOMIC PESTICIDE REGISTER-
 UNDER NO. 45-13 SUBJECT
 OF DETACHED COMMENTS.



HEAD



CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. In case of eye contact flush with plenty of water.

This product is toxic to fish. Keep out of any body of water.

Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purposes.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, moisture and wind movements are ideal for vaporization and drift of the vapor.

DED-WEED®

ME-6

4 LBS. 2,4-D PER GALLON

MIXING INSTRUCTIONS

Determine the amount of Ded-Weed ME-6 to be added in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed ME-6 recommended per acre. Add this amount of Ded-Weed ME-6 to one-half the water required for coverage. Mix thoroughly and add remaining water. Continue mixing until spray is uniform. Best results are obtained by treating weeds are growing vigorously.

FOR PERENNIAL WEEDS*

In Pasture, Rangelands and Non-Cultivated Land Such as Fence Rows, Right-of-Way, etc.

WEEDS	AMOUNT DED-WEED ME-6 PER ACRE	AMOUNT 2,4-D ACID PER ACRE	TIME OF APPLICATION
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 1/2 pints	3/4 lb.	Apply during early bud stage and again when rosettes are formed.
BINDWEED	1 1/2 pints	3/4 lb.	Apply bud to bloom stage.
WHITE TOP (HOARY CRESS)	1 to 1 1/2 pints	1/2 to 3/4 lb.	In spring when growing actively and foliage fully expanded.
RUSSIAN KNARWEED	1 1/2 pints	3/4 lb.	In spring before bloom.

for vaporization and drift of the vapor.

1 POUND, 12.5 POUNDS PER GALLON

RUSSIAN KNARWEED	1 1/2 pints	3/4 lb.	In spring be fore bloom	5 g
---------------------	-------------	---------	----------------------------	--------



**KILLS WEEDS IN CROP and NON-CROP AREAS,
ALONG FENCE ROWS and RIGHT-OF-WAYS**



**THOMPSON-HAYWARD
CHEMICAL COMPANY**

Kansas City, Kansas 66101

NET CONTENTS 1 GALLON





HEAD



MIXING INSTRUCTIONS

Determine the amount of Ded-Weed ME-6 to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed ME-6 recommended per acre. Add this amount of Ded-Weed ME-6 to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

FOR PERENNIAL WEEDS*

In Pasture, Rangelands and Non-Crop Land Such as Fence Rows, Right-of-Ways.

WEEDS	AMOUNT DED-WEED ME-6 PER ACRE	AMOUNT 2,4-D ACID PER ACRE	TIME OF APPLICATION	REMARKS
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 1/2 pints	3/4 lb.	Apply during early bud stage and again when rosettes are formed.	Use 1 1/2 pints Ded-Weed ME-6 per acre, each spraying.
BINDWEED	1 1/2 pints	3/4 lb.	Apply bud to bloom stage.	Reapply in late fall.
WHITE TOP (HOARY CRESS)	1 to 1 1/2 pints	1/2 to 3/4 lb.	In spring when growing actively and foliage fully expanded.	Retreat fall rosettes with 1 1/2 pints Ded-Weed ME-6 per acre.
RUSSIAN KNAWEED	1 1/2 pints	3/4 lb.	In spring before bloom.	Stops top growth and

**"USE CHEMICALS SAFELY.
FOLLOW LABEL DIRECTIONS!"**

WEED CONTROL IN CROPS

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

BROADLEAF WEEDS IN	AMOUNT DED-WEED ME-6 PER ACRE	AMOUNT 2,4-D ACID PER ACRE	TIME OF APPLICATION	REMARKS
SPRING WHEAT RYE, BARLEY	1/2 to 3/4 Pint	1/4 to 1/2 lb.	Apply full tiller to boot stage and after milk stage to harvest.	Do not treat in fall. Do not apply from boot to milk stage or in seedling stages. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
OATS	1/2 Pint	1/4 lb.		
FALL WHEAT	1/2 to 1 Pint	1/4 to 1/2 lb.		
CORN PRE-EMERGENCE (Field Corn)	1 to 1 1/2 Quarts	1 to 1 1/2 lbs.	Anytime after planting, preferably 2 to 5 days but before emergence.	Do not apply on sandy soil. Use recommended amounts in 5 to 20 gallons of water per acre.
CORN, SORGHUMS POST-EMERGENCE	3/8 to 1/2 Pint	3/16 to 1/4 lb.	When weeds are small and crop is 4 to 18 inches tall.	Use drop nozzles on crop more than 8 inches tall, to keep spray off foliage.
PASTURES RANGELAND	1 1/2 pints	3/4 lb.	When annual or perennial broadleaf weeds are growing actively fall or spring, perennials near bud stage.	Legumes present may be damaged. Do not graze dairy animals on treated areas within 7 days after application. Do not apply when grass is in boot to milk stage. Do not apply to seedling grasses, or after heading beans.

in boot to milk stage. Do not apply to seedling grasses, or after heading begins.

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED ME-6

- | | | |
|----------------------------|---------------------------|-----------------------------|
| <u>Beggar Ticks</u> | <u>Fanweed</u> | <u>Ragweed</u> |
| <u>Bindweed (Field)***</u> | <u>Lamb's Quarters</u> | <u>Russian Thistle***</u> |
| <u>Bull Thistle</u> | <u>Leafy Spurge***</u> | <u>Shepherds-Purse</u> |
| <u>Burdock</u> | <u>Marsh Elder</u> | <u>Saw Thistle (Annual)</u> |
| <u>Canada Thistle***</u> | <u>Mustards</u> | <u>Stinkweed</u> |
| <u>Carpet Weed</u> | <u>Pennywort</u> | <u>Sunflower</u> |
| <u>Chickweed</u> | <u>Pigweed</u> | <u>Wild Carrot</u> |
| <u>Cocklebur</u> | <u>Plantain</u> | <u>Wild Morning Glory</u> |
| <u>Curly Dock</u> | <u>Purslane (Pursley)</u> | |
| <u>Dandelion</u> | <u>Radish, Wild</u> | |

***REPEAT TREATMENTS NECESSARY

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

FOR DILUTION CALCULATIONS, 1 quart contains 1 lb. 2,4-D acid

Reduce Hazards!

FOLLOW THIS RINSE AND DRAIN PROCEDURE FOR PESTICIDE CONTAINERS



1
Empty container into spray tank. Then drain in vertical position for 30 seconds.



2
Add a measured amount of rinse water (or other diluent) so container is 1/4 to 1/5 full. For example, one quart in a one-gallon container



3
Rinse container thoroughly, pour into tank, and drain 30 sec. Repeat three times. Add enough fluid to bring tank up to level.




4
Crush pesticide container immediately. Sell as scrap for recycling or bury. Do not reuse.

EXAS

PLATE COMPANY, INC.

DATE	DESCRIPTION	SIZE	YELLOW	
6-2-64	THOMPSON CAN.	1 gal	RED	<input type="checkbox"/>
		marks	BLUE	<input type="checkbox"/>
		F style	BLACK	<input type="checkbox"/>
			VARNISH	<input checked="" type="checkbox"/>
			IMPRINT	<input type="checkbox"/>
			SPECIAL	<input type="checkbox"/>



OATS	1/2 Pint	1/4 lb		Stages Do not forage or graze treated grain fields with in 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
FALL WHEAT	1/2 to 1 Pint	1/4 to 1/2 lb		
CORN PRE-EMERGENCE (Field Corn)	1 to 1 1/2 Quarts	1 to 1 1/2 lbs	Anytime after planting, preferably 2 to 5 days but before emergence	Do not apply on sandy soil. Use recommended amounts in 5 to 20 gallons of water per acre.
CORN SORGHUMS POST-EMERGENCE	1/2 to 1 Pint	3/16 to 1/4 lb	When weeds are small and crop is 4 to 18 inches tall	Use drop nozzle on crop more than 8 inches tall to keep spray off foliage.
PASTURES RANGELAND	1 1/2 pints	3/4 lb	When annual or perennial broadleaf weeds are growing actively in the spring season in the treated stage.	Legumes present may be damaged. Do not graze dairy animals on treated areas within 7 days after application. Do not apply when grass is in boot to milk stage. Do not apply to seedling grasses, or after heading begins.

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED ME-6

- | | | |
|----------------------------|---------------------------|-----------------------------|
| <u>Beggar Ticks</u> | <u>Fanweed</u> | <u>Ragweed</u> |
| <u>Bindweed (Field)***</u> | <u>Lamb's Quarters</u> | <u>Russian Thistle***</u> |
| <u>Bull Thistle</u> | <u>Leafy Spurge***</u> | <u>Shepherds-Purse</u> |
| <u>Burdock</u> | <u>Marsh Elder</u> | <u>Saw Thistle (Annual)</u> |
| <u>Canada Thistle***</u> | <u>Mustards</u> | <u>Stinkweed</u> |
| <u>Carpet Weed</u> | <u>Pennywort</u> | <u>Sunflower</u> |
| <u>Chickweed</u> | <u>Pigweed</u> | <u>Wild Carrot</u> |
| <u>Cocklebur</u> | <u>Plantain</u> | <u>Wild Morning Glory</u> |
| <u>Curly Dock</u> | <u>Purslane (Pursley)</u> | |
| <u>Dandelion</u> | <u>Radish, Wild</u> | |

***REPEAT TREATMENTS NECESSARY

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

FOR DILUTION CALCULATIONS, 1 quart contains 1 lb. 2,4-D acid

Reduce Hazards!

FOLLOW THIS RINSE AND DRAIN PROCEDURE FOR PESTICIDE CONTAINERS



1
Empty container into spray tank. Then drain in vertical position for 30 seconds



2
Add a measured amount of rinse water (or other diluent) so container is 1/4 to 1/5 full. For example, one quart in a one-gallon container



3
Rinse container thoroughly, pour into tank, and drain 30 sec. Repeat three times. Add enough fluid to bring tank up to level



4
Crush pesticide container immediately. Sell as scrap for recycling or bury. Do not reuse

TEXAS
PLATE COMPANY, INC.

DATE JOB NO.	DESCRIPTION 1/2" x 1/4" x 1/8" A style	SIZE	YELLOW RED BLUE BLACK VARNISH IMPRINT SPECIAL
-----------------	-------------------------------------------	------	-----------------------------------------------------------------

