

DE-PESTER

## IMPORTANT

Since the effectiveness of RAT-TROL depends on small doses over a long time, baiting should be continued as long as evidence of feeding is observed. Except in extremely heavy infestations, this should be from 10 days to 15 days. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

## DIRECTIONS RATS

Put RAT-TROL BAIT in shallow containers (coffee can lids or empty sardine cans) and place in localities where rats are known to frequent. Each placement should amount to 1/4 to 1 pound of bait. Baits should be placed in locations not accessible to children, pets and domestic animals, or in tamper-proof bait boxes. Protection from rain or moisture also is desirable.

Baits should be examined regularly and replenished as long as feeding is evident. Baiting should continue for a minimum of 10 days.

Sour, moldy or otherwise decomposed bait should be replaced immediately.

## MICE

Follow the same directions as for rats except that the placements should be more numerous and smaller (1 tbs. of bait per placement) at 8 to 12 foot intervals where mice are known to be feeding and living. The period of baiting should also be longer to obtain good control. Baiting should continue for a minimum of 15 days.

**CAUTION:** RAT-TROL contains as its active ingredient an anti-coagulant chemical; therefore, baits prepared with RAT-TROL must be protected from or placed where humans, livestock, poultry or pets will not eat them. In case baits are eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

This material may not be effective under all conditions of use and particularly if directions are not followed. Since all effective rodenticides are more or less toxic, every precaution should be exercised in their use. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, or property.

**NOTE TO PHYSICIANS** When a human has been known to have ingested any RAT-TROL, blood transfusions, combined with intravenous injections and oral doses of Vitamin K, are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

## NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product

# RAT-TROL<sup>®</sup> BAIT

**CAUTION** Keep Out of Reach of Children  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

**CONTAINS DIPHACIN**

**KILLS MICE AND RATS**

ACCEPTED

FEB 5 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 148-200. SUBJECT  
TO ATTACHED COMMENTS.

### ACTIVE INGREDIENT

\*Diphacinone-2-diphenylacetyl 1,3-indandione.... 0.005%

INERT INGREDIENTS..... 99.995%

100.000%

\*U. S. Patent No. 2,672,483

**BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY**