

ture and wind movements are ideal for vaporization and drift of the vapor.

Do not re-use this container. Destroy by burying in a safe place. Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides, or other material to susceptible crops.

Do not store near fertilizers, seed, insecticides, fungicides or feed stuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover lawns.

Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.

MIXING INSTRUCTIONS

Determine the amount of Ded-Weed LV-69 to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed LV-69 recommended per acre. Add this amount of Ded-Weed LV-69 to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

FOR PERENNIAL WEEDS* IN PASTURE, RANGELANDS AND NON-CROP LANDS SUCH AS FENCE ROWS, DITCH BANKS, RIGHT-OF-WAYS

WEEDS	Amount Ded-Weed LV-69 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 quart	1 lb.	Apply during early bud stage and again when rosettes are formed.	Use 1 quart Ded-Weed LV-69 per acre, each spraying.
BINDWEED	3/4 to 1 quart	3/4 to 1 lb.	Apply bud to bloom stage.	Re-apply in late fall.
WHITE TOP (HOARY CRESS)	3/4 to 1 quart	3/4 to 1 lb.	In spring when growing actively and foliage fully expanded.	Re-treat fall rosettes with 1 to 2 quarts Ded-Weed LV-69 per acre.
RUSSIAN KNAPOWEED	1 quart	1 lb.	In spring before bloom.	Stops top growth and prevents seed production. Re-treat as necessary.

*Hard-to-kill weeds, such as Canadian Thistle, may not be controlled by one application but top growth and re-seeding usually will be prevented. Repeated treatments generally give good control.

For control of resistant species of brush and certain timberland applications, use other Thompson-Hayward products such as Ded-Weed LV-33 Brush Kil and Ded-Weed LV-6 Brush Kil.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ESTER

DED-WEED. LV-69

LOW VOLATILE ESTER

CONTAINS 2,4-D 4 LBS. PER GALLON

CAUTION Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

KILLS WEEDS IN PASTURES, CROP and NON-CROP AREAS, FENCE ROWS and RIGHT-OF-WAYS

ACTIVE INGREDIENTS:	
2,4-dichlorophenoxyacetic acid isooctyl ester.....	67.0%
(Equivalent to 2,4-dichlorophenoxyacetic acid—44.4%)	
INERT INGREDIENTS	33.0%
	100.0%

USDA Reg. No. 148-241

12-67



DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY
WEED CONTROL IN CROPS

BROADLEAF WEEDS IN	Amount Ded-Weed LV-69 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
SPRING WHEAT, RYE, BARLEY	3/4 to 3/4 pint	3/4 to 3/4 lb.	Apply full tiller to boot stage and after milk stage to harvest.	Do not treat in fall. Do not apply from boot to milk stage or in seedling stages.
OATS	3/4 Pint	3/4 lb.		
FALL WHEAT	3/4 to 1 pint	3/4 to 3/4 lb.		Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
FIELD CORN PRE-EMERGENCE	1 to 1 1/2 quarts	1 to 1 1/2 lbs.	Any time after planting, preferably 2 to 5 days, but before emergence.	Do not apply on sandy soil. Use recommended amounts in 5 to 20 gallons of water per acre.
CORN, SORGHUMS POST-EMERGENCE	3/4 to 3/4 pint	3/4 to 3/4 lb.	When weeds are small and crop is 4 to 18 inches tall.	Use drop nozzles on crop more than 8 inches tall to keep spray off foliage.
PASTURES, RANGELAND	3/4 to 1 quart	3/4 to 1 lb.	When annual or perennial broadleaf weeds are growing actively, fall or spring; perennials near bud stage.	Legumes present may be damaged. Do not graze dairy animals on treated areas within 7 days after application. Do not apply to seedling grasses.

FOR DILUTION CALCULATIONS—
1 quart contains 1 lb. 2,4-D acid

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED LV-69

Carpet Weed Lamb's Quarters Pigweed
Cocklebur Mustards Wild Morning Glory
If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. This product is toxic to fish. Keep out of any body of water. Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purposes.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.). Although this product has been formulated with low volatile esters in order to reduce the hazard of vapor damage, vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, mois-