

DE-PESTER

CAUTION!

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid prolonged breathing of spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate utensils, feed, water or foodstuffs. Keep out of the reach of children. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.


DO NOT SPRAY DIRECTLY ON SKIN OR ON ANIMALS. Remove birds and pets and cover fish bowls and aquaria before spraying room. An average-sized room would require 15-20 seconds of spraying.

CAUTION

Contents Flammable.
Do Not Puncture, Incinerate, or Store Above 120 F.
Do Not Use Near Open Flame or Electrical Spark.
Contains Petroleum Distillate
Keep Out of Reach of Children.

~~NOTICE: THIS PRODUCT DOES NOT Warrant any warranty as to hazards or results.~~

*Equivalent to 0.5% VAPONA® Insecticide



NOX-INSEX^R

KILLS


Roaches, Mosquitoes, Flies, Ants,
Silverfish and Flying Moths

CAUTION—CONTENTS UNDER PRESSURE
Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

ACTIVE INGREDIENTS	BY WEIGHT
2, 2-dichlorovinyl dimethyl phosphate	0.46%*
Related Compounds	0.04%*
Petroleum Hydrocarbons	14.50%
INERT INGREDIENTS	85.00%
TOTAL	100.00%

USDA Reg. No. 148-259



THOMPSON-HAYWARD CHEMICAL COMPANY
Kansas City, ~~MISSOURI~~ **KANSAS**

BEFORE USE READ CAUTION STATEMENT CAREFULLY

To control mosquitoes, flies, gnats, flying moths and similar insects close all doors and windows and direct mist upward and spray toward four corners of the room. Completely fill the room with mist, which can be accomplished in 15 to 20 seconds, for a room 10x12x3 (approximately 1000 cubic feet).

Silverfish, roaches, water bugs and ants may be controlled by directing spray into all places where insects seek shelter and treatment should include treatment of all cracks and crevices (for control of ants, also spray ant trails and places of entrance such as door frames and window sills). Contact as many insects as possible when spray is applied. Best results are obtained if treated area is left closed for at least 15 minutes after treatment. Repeat application as frequently as necessary for insect control. All cooking utensils, dishes, food containers and food stuff should be covered before applying spray.

HOW TO APPLY: Hold container upright directing nozzle toward area to be treated. Press button top of container.

ACCEPTED WITH COMMENTS

NOTICE: SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, CONCERNING THE USE OF THIS PRODUCT OTHER THAN INDICATED IN THE LABEL. SELLER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.