

MIXING INSTRUCTIONS

Determine the amount of Ded-Weed ME-9 to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed ME-9 recommended per acre. Add this amount of Ded-Weed ME-9 to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

FOR PERENNIAL WEEDS* IN PASTURE, RANGELANDS AND NON-CROP LAND SUCH AS FENCE ROWS, RIGHT-OF-WAYS

WEEDS	Amount Ded-Weed ME-9 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 1/3 pints	1 lb.	Apply during early bud stage and again when rosettes are formed.	Use 1 1/3 pints Ded-Weed ME-9 per acre, each spraying.
BINDWEED	1 to 1 1/3 pints	3/4 to 1 lb.	Apply bud to bloom stage.	Re-apply in late fall.
WHITE TOP (HOARY CRESS)	3/4 to 1 1/3 pints	1/2 to 1 lb.	In spring when growing actively and foliage fully expanded.	Re-treat fall rosettes with 1 1/3 to 2 2/3 pints Ded-Weed ME-9 per acre.
RUSSIAN KNAPWEED	1 1/3 pints	1 lb.	In spring before bloom.	Stops top growth and prevents seed production. Re-treat as necessary.

*Hard-to-kill weeds, such as Canada Thistle, may not be controlled by one application but top growth and reseeding usually will be prevented. Repeated treatments generally give good control.

CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. This product is toxic to fish. Keep out of any body of water. Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purposes.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, moisture and wind movements are ideal for vaporization and drift of the vapor.

Do not re-use this container. Destroy by burying in a safe place. Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides, or other material to susceptible crops.

Do not store near fertilizers, seed, insecticides, fungicides or foodstuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover lawns.

Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.



DED-WEED. ME-9

BUTYL ESTER

2,4-D 6 LBS. PER GALLON

CAUTION Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

KILLS WEEDS IN PASTURES, CROP and NON-CROP AREAS, FENCE ROWS and RIGHT-OF-WAYS

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid butyl ester..... 76.4%
(Equivalent to 2,4-dichlorophenoxyacetic acid—61.0%)

INERT INGREDIENTS 23.6%
100.0%

USDA Reg. No. 148-289



DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

CROP USE

BROADLEAF WEEDS IN	Amount Ded-Weed ME-9 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
SPRING WHEAT, RYE, BARLEY†	1/3 to 1/2 pint	1/4 to 1/3 lb.	Apply full tiller to boot stage and after milk stage to harvest	Do not treat in fall. Do not apply from boot to milk stage or in seedling stages.
OATS	1/3 pint	1/4 lb.		Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
FALL WHEAT†	1/3 to 2/3 pint	1/4 to 1/2 lb.		
CORN PRE-EMERGENCE (Field Corn)	1 1/3 to 2 pints	1 to 1 1/2 lbs.	Any time after planting, preferably 2 to 5 days, but before emergence.	Do not apply on sandy soil. Use recommended amounts in 5 to 20 gallons of water per acre.
CORN, SORGHUM POST-EMERGENCE†	1/4 to 1/3 pint	1/8 to 1/4 lb.	When weeds are small and crop is 4 to 12 inches tall.	Use drop nozzles on crop more than 8 inches tall to keep spray off foliage.
PASTURES, RANGELAND	1 to 1 1/3 pints	3/4 to 1 lb.	When annual or perennial broadleaf weeds are growing actively, fall or spring; perennials near bud stage.	Legumes present may be damaged. Do not graze dairy animals on treated areas within 7 days after application. Do not apply to seedling grasses.

†Always apply 2,4-D with enough water to give uniform coverage. For aerial applications, 2 1/2 to 4 gallons of spray solution per acre are usually recommended.

FOR DILUTION CALCULATIONS—
1 1/3 pints contain 1 lb. 2,4-D acid

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED ME-9

Carpet Weed Lamb's Quarters Pigweed
Cocklebur Mustards Wild Morning Glory

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

For control of resistant species of brush and certain timberland application use other Thompson-Hayward products such as Ded-Weed LV-33 Brush Kil and Ded-Weed LV-6 Brush Kil.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and/or handling is contrary to label instructions.