

MIXING INSTRUCTIONS

Determine the amount of Ded-Weed BE-6D to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed BE-6D recommended per acre. Add this amount of Ded-Weed BE-6D to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. This product is toxic to fish. Keep out of any body of water.

Avoid contamination of water supplies that may be used to irrigate or water susceptible crops or used for domestic purposes. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, moisture and wind movements are ideal for vaporization and drift of the vapor.

Do not re-use this container. Destroy by burying in a safe place.

Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides or other material to susceptible crops.



2,4-D 6 POUNDS PER GALLON BUTYL ESTER

ACTIVE INGREDIENT:		
2,4-dichlorophenoxyacetic acid butyl ester*	76.4%	
INERT INGREDIENTS:	23.6%	
	Total	100.0%

*Equivalent to 2,4-dichlorophenoxyacetic acid—61.0%

KILLS WEEDS IN PASTURES, CROP and NON-CROP AREAS, FENCE ROWS and RIGHT-OF-WAYS

CAUTION

Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACCEPTED
 148-289
 JUL 12 1977

NET CONTENTS GAL. (litres)



P.O. BOX 2383, KANSAS CITY, KANSAS 66110

Do n
sectr
caus-
gras-
Do n
flame
meth-
age c
erty :

FO
11

CO

Carp-
Cock
Lamb

If sto
may
agita
For c
and
other
as D
LV-4

NOTI
press
of th
the li
and/
such
label

Do not store near fertilizers, seed, insecticides, fungicides or foodstuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover. Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.

FOR DILUTION CALCULATIONS—

1 1/3 pints contain 1 lb. 2,4-D acid

PARTIAL LIST OF WEEDS CONTROLLED BY JED-WEED BE-60

Carpet Weed Mustards
Cocklebur Pigweed
Lamb's Quarters Wild Morning Glory

If stored below freezing temperatures it may be necessary to warm to 40° F and agitate thoroughly before using. For control of resistant species of brush and certain timberland application use other Thompson-Hayward products such as Ded-Weed LV-2D2T and Ded-Weed LV-4T.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**DIRECTIONS
BEFORE USE READ DIRECTIONS
STATEMENT OF CONTROL**

BROADLEAF WEEDS IN	Amount Ded-Weed BE-60 Per Acre	Amount 2,4-D Acid Per Acre	
SPRING WHEAT RYE, BARLEY:	1 3/4 to 1 1/2 pint	1/4 to 1/3 lb	Apply after m
OATS	1 3/4 pint	1/4 lb	
FALL WHEAT:	1 3/4 to 2 3/4 pint	1/4 to 1/2 lb	
CORN PRE-EMERGENCE (Field Corn)	1 1/3 to 2 pints	1 to 1 1/2 lbs	Any time 2 to 5 c
CORN, SORGHUM POST EMERGENCE:	1 1/4 to 1 3/4 pint	3/16 to 1/4 lb	When crop is 4
PASTURES, RANGELAND	1 to 1 1/3 pt	3/4 to 1 lb	When weeds are perennial

**Do not graze dairy animals on treated areas within 7 days heading begins. Do not apply when grass is in boot to milking stage.
†Always apply 2,4-D with enough water to give uniform solution per acre are usually recommended

FOR PERENNIAL WEEDS* II AND NON-CROP LANDS RIGHT-O

WEEDS	Amount Ded-Weed BE-60 Per Acre	Amount 2,4-D Acid Per Acre	
CANADA THISTLE AND PERENNIAL SON THISTLE	1 1/4 pint	1 lb	Apply during winter
BINDWEED	1 to 1 1/3 pint	3/4 to 1 lb	Apply bud
WHITE TOP (HOARY CRESS)	2 3/4 to 1 1/3 pt	1/2 to 1 lb	In spring and foliage
RUSSIAN KNAWEED	1 1/3 pint	1 lb	In spring or

*Hard-to-kill weeds such as Canada Thistle may not be controlled usually will be prevented. Repeated treatments generally give

148-289

3.4%
3.6%
1.0%

KS 1
S)

10
1

Do not store near fertilizers, seed, insecticides, fungicides or foodstuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover. Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.

FOR DILUTION CALCULATIONS—

1 1/3 pints contain 1 lb. 2,4-D acid

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED BE-6D

Carpet weed Mustards
Cocklebur Pigweed
Lamb's Quarters Wild Morning Glory

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using. For control of resistant species of brush and certain timberland application use other Thompson-Hayward products such as Ded-Weed LV-2D2T and Ded-Weed LV-4T.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

148 - 289

**DIRECTIONS
BEFORE USE READ DIRECTIONS, CAUTION AND STATEMENTS CAREFULLY
CROP USE**

BROADLEAF WEEDS IN	Amount Ded-Weed BE-6D Per Acre	Amount 2,4 D Acid Per Acre	TIME OF APPLICATION
SPRING WHEAT RYE, BARLEY:	1 3 to 1 2 pint	1 4 to 1 3 lb	Apply full rate to boot stage and after milk stage to harvest
OATS	1 3 pint	1 4 lb	
FALL WHEAT:	1 3 to 2 3 pint	1 4 to 1 2 lb	
CORN PRE-EMERGENCE (Field Corn)	1 1 3 to 2 pints	1 to 1 1 2 lbs	Any time after planting, preferably 2 to 5 days but before emergence
CORN, SORGHUM POST EMERGENCE:	1 4 to 1 3 pint	3 16 to 1 4 lb	When weeds are small and crop is 4 to 12 inches tall
PASTURES, RANGELAND	1 to 1 1 3 pt	3 4 to 1 lb	When annual or perennial broadleaf weeds are growing actively, fall or spring, perennials near bud stage

**Do not graze dairy animals on treated areas within 7 days after application. Do not head out begins. Do not apply when grass is in boot to milk stage

†Always apply 2,4-D with enough water to give uniform coverage. For aerial application solution per acre are usually recommended

FOR PERENNIAL WEEDS* IN PASTURE, RANGE AND NON-CROP LAND SUCH AS FENCE ROWS, RIGHT-OF-WAYS

WEEDS	Amount Ded-Weed BE-6D Per Acre	Amount 2,4 D Acid Per Acre	TIME OF APPLICATION
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 1 3 pt	1 lb	Apply to the early bud stage and again when rosettes are formed
BINDWEED	1 1 3 pt	1 4 to 1 3 lb	Apply bud to bloom stage
WHITE TOP (HOARY CRESS)	1 1 3 pt	1 2 to 1 lb	In spring when growing actively and foliage fully expanded
RUSSIAN Knapweed	1 1 3 pt	1 lb	Apply in spring before bloom

*Hard-to-kill weeds such as Canada Thistle may not be controlled by one application. Repeated treatments generally give good control