

DE-PESTER®

DIRECTIONS

FLY CONTROL For the reduction of flies in and around buildings which house domestic animals, around yards and homes, apply spray or bait prepared according to table below. Apply spray to thoroughly wet 1 gal. of diluted spray per 1000 square feet of painted surface, 2 gallons of diluted spray per 1000 square feet of an painted surface where flies alight or congregate, such as walls, ceilings, stanchions in dairy barns, fences, manure piles, and around garbage cans. Livestock should be removed before making applications to animal shelters. Do not use in dwellings as necessary.

STRAIGHT SPRAYS		BAIT SPRAYS		
Amount Spray Desired	Amount Malathion W25 to use	Amount Malathion W25 to use	Add Either	
			Sugar	Molasses* -or Corn Syrup
2½ gals.	1 lb.	1 lb.	1 cup	1 cup
12 gals.	5 lbs.	5 lbs.	2½ lbs.	1 qt.
100 gals.	40 lbs.	40 lbs.	20 lbs.	2 gals.

*Do not use Molasses treated with preservative.

PLANT AND ANIMAL USES ALL RATES OF APPLICATION ARE FOR USE IN 100 GALLONS OF WATER

SMALL VOLUME RATE 5 level teaspoons per gallon = 2½ lbs./100 gal. water

VEGETABLES Apply 100 gal. of spray per acre or 1 gal. per 435 square feet. To control spider mites and aphids on Tomatoes and Eggplants; aphids on Peppers, Head Lettuce and Brussels Sprouts, and leafhoppers on Potatoes, apply 2½ pounds. To control aphids, imported cabbageworm and cabbage looper on Cabbage, Broccoli, Kale, Turnips and Mustard, spray 2½ pounds. Control thrips on Onions with 4 pounds. Repeat application as needed for continued control. Do not spray Tomatoes, Broccoli, Eggplant, Peppers or Onions within 3 days of harvest nor Cabbage, Kale, Turnips, Mustard, Brussels Sprouts or Lettuce within 7 days.

FRUIT Spray with 2½ pounds for control of rosy aphids, forbes scale, and mealybug on Apples; mealybug on Pears, leafhoppers, black-headed fireworms, spittlebug nymphs and cranberry fruitworm on Cranberries; aphids and spider mites on Strawberries. 2-2½ pounds are required for control of European red mite, clover mite, Willamette mite, and two spotted mite on Apples; pear psylla and mites on Pears; European red mites and two spotted mite on Peaches; 2 pounds will control woolly apple aphids, green apple aphids, and bud moth on Apples, black cherry aphids on Cherries. Apply spray at rate of 100 gallons per acre on Cranberries and Strawberries.

Dilute 3 pounds for control of codling moth, plum curculio and red banded leafroller on Pears and Apples, also fruit tree leafroller on Pears; Oriental fruit moth, plum curculio on Peaches. Apply 1-1½ pounds for leafhopper control on Grapes. Use 1 pound for control of mealy plum aphids on Plums and Prunes.

*Make at least two applications 10-12 days apart in summer months.

Malathion W25 may injure Bosc Pears in the Northeast or some varieties of Sweet Cherries, especially in the Northwest.

On all applications repeat as required for continued control. Do not apply within 3 days of harvest on Apples, Pears, Cherries, Cranberries, Grapes, Plums, Prunes, Strawberries 7 days on Peaches. May cause injury to Ribier Grapes. Malathion may injure foliage on some varieties of apple.

ORNAMENTALS Make thorough full-coverage sprays with repeated applications as necessary. For scales, apply when crawlers have settled on foliage. Injury may occur on certain ferns, including Boston Maidenhair and Pteris as well as some species of crassula. Use 2 pounds in 100 gallons of water to control tent caterpillar.

Use 2½ pounds per 100 gallons of water to control aphids, spider mites, whitefly and mealybugs. Use 3 pounds per 100 gallons of water to control Fleischer scale. Use 4 pounds per 100 gallons of water to control lacebugs, bagworms, and pine needle scale. Use 6 pounds per 100 gallons of water to control Florida red scale, soft brown scale, Monterey pine scale, black scale crawlers and Juniper scale.

ANIMALS To control lice on Beef Cattle and Hogs use 16 pounds per 100 gallons of water. For control of horn flies on Beef Cattle use 16 to 24 pounds, for ticks on Beef Cattle use 16 to 32 pounds. Apply a complete coverage spray. Apply also to hog pens and litter. For lice, one treatment may be sufficient, repeat only if needed. For ticks and horn flies, repeat applications at two week intervals if needed. **DO NOT APPLY TO DAIRY ANIMALS. DO NOT TREAT CALVES UNDER ONE MONTH OF AGE.**

CAUTION Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin, wash thoroughly after using. Avoid contamination of foodstuffs, food containers, feed and watering troughs. Keep out of reach of children. Don't spray in room where edible products are handled or in milk processing rooms. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances. This product is toxic to fish. Keep out of any body of water.

NOTICE : Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and/or handling is contrary to label instructions.

MALATHION W-25

CAUTION Keep Out of Reach of Childrer
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

Contains 25% MALATHION

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

ACTIVE INGREDIENTS

Malathion	25.0%
INERT INGREDIENTS	75.0%
Total	100.0%

O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

U.S.D.A. Reg. No. 148-303

10-66



ACCEPTED WITH COMMENTS