

Contains METHYL PARATHION

Do not mix or use in combination with other organic phosphates. Not suitable for use in hand-operated or knapsack-type sprays. This material must not be used as an insecticide without dilution.

For Use by Trained Operators in Airplane and Power Sprayers Only

DESTROY CONTAINER WHEN EMPTY

DE-PESTER®

METHYL PARATHION

E-2

2 lbs. Per Gallon

DIRECTIONS FOR USE

BEFORE USE READ WARNING AND NOTICE STATEMENTS CAREFULLY

Methyl Parathion E-2 is an emulsifiable concentrate containing 2 pounds of methyl parathion to the gallon for use upon dilution as a spray in the control of various insects and mites infesting agricultural crops.

MIXING INSTRUCTIONS

Add the recommended amount of **Methyl Parathion E-2** to about one-half the volume of water to be used in spraying. Mix well, then add the remaining water and mix until spray mixture is uniform. Continue mixing while spraying.

COTTON INSECTS—BOLL WEEVIL: Use 1 to 2 pints per acre for each application. In areas where weevils are present regularly, pre-square treatment consisting of 2 applications at 7- to 10-day intervals, beginning when cotton first starts fruiting, is recommended. In midseason, before peak fruiting, begin treatment after 25% of the squares are punctured. Continue spraying at 8-day intervals until control is effective. Resume treatment if insects reappear.

COTTON APHID: Use 1 to 2 pints per acre for each application. Usually as this spray is applied to control other insects, aphid infestations are reduced. Repeat as needed for continued control.

OTHER COTTON INSECTS: Use 1 to 2 pints per acre for the control of cotton leafworm and spider mites. Repeat applications as required for continued control. Do not apply within 5 days of picking.

SMALL GRAIN (Wheat, Oats, Barley, Rye) INSECTS: Greenbugs and other aphids; Use 1 to 1½ pints per acre for each application. Treatment should be repeated when reinfestation occurs. Do not apply after heads begin to form.

ALFALFA AND VETCH: For control of the spotted alfalfa aphid apply 1 to 1½ pints per acre in sufficient water for good coverage. Make no application within 15 days of pasturing or harvest.

CAUTION: Misuse of quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

Do not mix or use in combination with other organic phosphates. Not suitable for use in hand-operated or knapsack-type sprays. This material must not be used as an insecticide without dilution.

Workers entering fields within 24 hours after application should wear protective clothing.

NOTICE

Some insects may be resistant to the use of this material. For more information on the proper use of this material, contact your local agricultural agent or the manufacturer.

WARNING

FIRST AID TREATMENT
CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING SYMPTOMS
SEE NOTE TO PHYSICIAN BELOW

INHALATION: Remove patient from contaminated atmosphere; keep him under continuous observation. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.
SKIN: Immediately remove contaminated clothing and scrub skin with soap and water for 15 minutes. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.
INTERNAL: If swallowed, make patient vomit by giving warm salty or soapy water. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PATIENT
NOTE FOR PHYSICIANS
SYMPTOMS OF POISONING

TREATMENT

DO NOT USE BOTTLE

DANGER Keep Out of Reach of Children

SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING STATEMENTS ON OTHER PANELS.



POISON



ATROPINE IS AN ANTIDOTE. Keep an emergency supply on hand.

ACTIVE INGREDIENTS:

Methyl parathion 2.00 lbs. per gallon

INERT INGREDIENTS:

READ WARNING, NOTICE, CAUTION AND FIRST AID TREATMENT BEFORE USE

U.S. Reg. No. 146-337

1-68



Contains METHYL PARATHION

Do not mix or use in combination with other organic phosphates. Not suitable for use in hand-operated knapsack-type sprays. This material must not be used as an insecticide without dilution.

For Use by Trained Operators in Airplane and Power Sprayers Only

DE-PESTER®

METHYL PARATHION E-2

2 lbs. Per Gallon

DIRECTIONS FOR USE

BEFORE USE READ WARNING AND NOTICE STATEMENTS CAREFULLY.

Methyl Parathion E-2 is an emulsifiable concentrate containing 2 lbs. per gallon of active ingredient. It is used for the control of various insect pests infesting cotton, alfalfa, and vetch.

MIXING INSTRUCTIONS

Add the recommended amount of **Methyl Parathion E-2** to about one-half the volume of water to be used in spraying. Mix well, then add the remaining water and mix until a uniform mixture is formed. Continue mixing while spraying.

COTTON INSECTS—BOLL WEEVIL: 1 to 2 pints per acre for each application in areas where weevils are present. Apply early, pre-bloom treatment consisting of 2 applications at 7 to 10 day intervals. Begin weevil control at first cotton bloom. Recommended for multiple applications. For control of weevils on cotton after the first bloom, apply 1 pint per acre. Control of weevils on cotton after the first bloom is best controlled by the use of a combination of insecticides.

COTTON APHID: 1 to 2 pints per acre for each application. Apply early, pre-bloom treatment consisting of 2 applications at 7 to 10 day intervals. Begin aphid control at first cotton bloom. Recommended for multiple applications. For control of aphids on cotton after the first bloom, apply 1 pint per acre.

OTHER COTTON INSECTS: 1 to 2 pints per acre for each application. Apply early, pre-bloom treatment consisting of 2 applications at 7 to 10 day intervals. Begin control of other cotton insects at first cotton bloom. Recommended for multiple applications. For control of other cotton insects on cotton after the first bloom, apply 1 pint per acre.

SMALL GRAIN (Wheat, Oats, Barley, Rye) INSECTS: 1 to 2 pints per acre for each application. Apply early, pre-bloom treatment consisting of 2 applications at 7 to 10 day intervals. Begin control of small grain insects at first grain bloom. Recommended for multiple applications. For control of small grain insects on small grain after the first bloom, apply 1 pint per acre.

ALFALFA AND VETCH: 1 to 2 pints per acre for each application. Apply early, pre-bloom treatment consisting of 2 applications at 7 to 10 day intervals. Begin control of alfalfa and vetch insects at first alfalfa and vetch bloom. Recommended for multiple applications. For control of alfalfa and vetch insects on alfalfa and vetch after the first bloom, apply 1 pint per acre.

CAUTION: Do not mix or use in combination with other organic phosphates. Not suitable for use in hand-operated or knapsack-type sprays. This material must not be used as an insecticide without dilution.

Workers entering fields within 24 hours after application should wear protective clothing.

NOTICE

DANGER Keep Out of Reach of Children

SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING STATEMENTS ON OTHER PANELS.

ACTIVE INGREDIENTS

INERT INGREDIENTS

READ WARNING, NOTICE, CAUTION AND FIRST AID TREATMENT BEFORE USE



FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING SYMPTOMS
SEE NOTE TO PHYSICIAN BELOW

INHALATION: Remove patient from contaminated area. Keep him under continuous observation. Do not wait for effect of atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.

SKIN: Immediately remove contaminated clothing and scrub skin with soap and water for 15 minutes. Do not wait for effect of atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.

INTERNAL: If swallowed, induce vomiting with 1 cup of water. Do not wait for effect of atropine (2 tablets) at once if symptoms include blurred vision or pinpoint pupils.