

148-359

UNDER THE FEDERAL INSECTICIDE ACT
 REGISTERED IN THE OFFICE OF THE
 FEDERAL REGISTER
 REGISTERED UNDER NO. 148-359



DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY.

CROP	INSECT	Amount to Use Per Acre (Pounds)	Intervals (Days Between Last Application and Harvest)
Strawberries	Aphids, Spider Mites	40	3
Beans**	Mexican Bean Beetle, Leafhopper	30-35	1
Cabbage	Aphids, Imported Cabbageworm, Cabbage Looper	30	7
Cucumber, Squash, Melons†	Aphids, Spider Mites, Leafhopper on Melon	30-35	1
Onions	Thrips	30-40	3
Peas	Pea Aphid	25	3
Potatoes	Aphids, Leafhopper	25-30	3
Tomatoes	Aphids, Spider Mites	35-40	1
Alfalfa-	Aphids, Potato Leafhopper, Spider Mites, Alfalfa Weevil, Larvae, Spittlebug	30	7
Ornamentals‡	Aphids, Spider Mites, Bagworms	Full Coverage Applications**	

**Make two or more applications as needed.
 †Do not apply Malathion to Cucurbits unless plants are dry.
 ‡Do not apply to Alfalfa in bloom. No application within 7 days of harvest or pasturing. Make repeat applications as needed.
 †Injury may occur on certain trees and on some species of crassula and canetti juniper. 30 lbs per acre on trees per 1,000 sq. ft. 40 lbs. per 1,450 sq. ft.

ANIMAL INSECT CONTROL

DAIRY AND BEEF CATTLE: For control of hornflies apply 3 tablespoons per animal. Apply dust on back and sides of animal and repeat application at 10-14 day intervals. Do not apply to dairy cattle more than five hours before milking or during milking. Do not use spray on dairy animals in Malathion. **BEEF CATTLE AND HOGS:** Control lice by spraying or dusting through covering of dust. Thoroughly dust hog pens. Repeat in 10 days thereafter as needed. Do not contaminate water troughs.

EGGS AND CATS: For control of ticks, apply sufficient dust to treat animals thoroughly. Use any garden type dust applicator or rotary type duster. Repeat application every 10-14 days. **ANIMAL QUARTERS:** For control of ticks, apply dust to walls, ceiling, and floor. Use rotary duster or puff duster. Apply 10-15 lbs per 1,000 sq. ft. of area.

POULTRY: Premise treatment for control of chicken red mite, northern fowl mite, poultry lice, apply one lb per 10 to 20 square feet of litter and floor space and to nests, roosts and adjacent areas. Apply with rotary hand duster, puff duster or other container. Repeat application every 10 days. Avoid contamination of feeds, troughs and water troughs.

CAUTION: Avoid contact with skin and clothing. Wash thoroughly after use. Avoid contamination of feeds and water troughs. Do not use within reach of children.

ENVIRONMENTAL CAUTIONS

Avoid contamination of water bodies. Do not use in areas where it may be injurious to fish, birds, bees, or other beneficial insects. Do not use in areas where it may be injurious to plants.

SEE CAUTION

NOTICE

This product is a restricted use pesticide. It is not to be used by anyone who is not registered with the EPA under the Federal Insecticide, Fungicide, and Rodenticide Act. For more information, contact the EPA at Washington, D.C. 20460.

MALATHION DUST No. 5

CAUTION Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

Contains 5% MALATHION

CONTROLS CROP, ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:

Malathion* 5.0%

INERT INGREDIENTS

..... 95.0%

100.0%

* (O,O-dimethyl phosphorothioic acid dimethylmercaptosuccinate)



**THOMPSON-HAYWARD
CHEMICAL COMPANY**

EPA EST. NO. 148-359