


[148-387] 1 of 2



PHOSDRIN*

E-2

2 Lbs. Per Gallon



DANGER Keep Out of Reach of Children

SEE ANTIDOTE STATEMENT AND OTHER REQUIRED
WARNING STATEMENTS ON SIDE PANELS



Do not mix or use in combination with other organic phosphates. Not suitable for use in hand-operated or knapsack-type sprayers. This material must not be used as an insecticide without dilution.

ACTIVE INGREDIENTS

INERT INGREDIENTS

pounds of actual Phosdrin) per acre. Apply with sufficient water to give uniform coverage. Begin applications when infestation appears and repeat as often as necessary to maintain control. Do not apply within 3 days of harvest.

CABBAGE, BROCCOLI AND LETTUCE: For control of Cabbage Looper, Imported Cabbage Worm and Aphids on cabbage, broccoli and lettuce, apply Phosdrin E-2 at the rate of one to two pints (1/4 to 1/2 pound of actual Phosdrin) per acre. Apply with sufficient water to give uniform coverage. Begin application when infestation appears and repeat as often as necessary to maintain control. Do not apply within 4 days of harvest of cabbage, within 3 days of harvest of broccoli or within 7 days of harvest of lettuce.

BEANS, TOMATOES, CLOVER, POTATOES, PEAS, ALFALFA: 1 quart per acre for control of lygus, aphid, spider mites, leaf hopper. For use on potatoes, do not exceed 1 pint per acre.

Do not use within 1 day of harvest.

STRAWBERRIES: 1 quart per acre for control of spider mite, aphid, lygus, salt marsh caterpillar. No application within one day of picking.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Birds feeding on treated areas may be killed.

Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PRECAUTIONS IN USING

Evidence of small wounds or absorption through the skin. Do not apply or allow drift to reach occupant or unprotected limbs. Operators should wear rubber gloves, water-tight protective clothing and effective mask. Frequent washing and drying of clothes to prevent residue from reaching and staining them. Keep out of treated fields for rest of day following treatment. Keep out of reach of children and domestic animals. Do not reuse containers by washing them with any solution; they destroy used containers to prevent reuse.

Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

Apply to any area not specified on the label. Birds feeding on treated areas may be killed.

FIRST AID TREATMENT

Atropine is the emergency antidote for Phosdrin poisoning. Atropine is obtainable only on a doctor's prescription. Consult your physician about obtaining an adequate supply of 1/100 grain Atropine tablets for emergency use.

2-PAM is also antidotal and may be administered in conjunction with atropine.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water) or soapy water. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. If material has been spilled on the skin, immediately remove the patient from the vicinity of the insecticide. Remove contaminated clothing and wash skin with soap and water. If the material has been accidentally splashed into the eyes, wash immediately with clear water for at least ten minutes. If warning symptoms appear, administer two 1/100 grain Atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (SEE WARNING SYMPTOMS BELOW.) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, reduced vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antidote of choice against the parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10 minute intervals in dose of 1 to 2 mg. grams until complete atropinization has occurred. Mouth care and intratracheal clear chest and postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause profound toxicity. In very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

DESTROY CONTAINERS WHEN EMPTY

Flush out thoroughly with water. Do not use where the water can be run off harmfully. Do not use in areas where holes or cracks exist that cannot be repaired for any purpose. Take to city dump or other suitable place where disposal attempt will be made to salvage for other use where possible.

NOTICE: Do not use this material in any area where it may be applied to the ground. Do not use this material in any area where it may be applied to the ground. Do not use this material in any area where it may be applied to the ground. Do not use this material in any area where it may be applied to the ground.