

ACCEPTED

7 1 6 1968

INSECTICIDE  
IDENTIFICATION  
SUBJECT  
COMMENTS

Net  
contents

Net  
contents

TOX - L SPRAY - DIP

**WARNING** Keep Out of Reach of Children  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

<b>ACTIVE INGREDIENTS:</b>	
Toxaphene*	45.00%
Gamma isomer of benzene hexachloride (from lindane).....	1.96%
Aromatic petroleum derivative solvent*.....	33.54%
Petroleum distillate.....	6.00%
<b>INERT INGREDIENTS</b>	13.50%
	<b>100.00%</b>

\*Technical chlorinated camphene (chlorine content 67-69%)

USDA Reg. No. 18-495

2-68

Manufactured for

Anchor Serum Company

St. Joseph, Missouri 64502

## BACKRUBBER USE

To aid in controlling Horn Flies, Mosquitoes, Gnats and Lice on Beef Cattle, mix thoroughly one pint of Stock Spray and Dip with 1½ gallons of diesel oil or No. 2 fuel oil. Apply mixture to rubbing device covering or pour into oil reservoir. Be sure covering is soaked.

Recharge units with mixture every 3 to 5 weeks as required to keep units working efficiently. Regular exposure of animals to treated backrubber for about 10 weeks is usually necessary for good results on lice. Caution—Do not mix with used motor or crankcase oil or greases. Do not allow calves under 6 months of age to use rubbing devices. Do not allow beef animals to use treated backrubber within 30 days of slaughter.

Do not treat sows within two weeks of farrowing, nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

**RESIDUAL SURFACE APPLICATION:** For control of Flies and Mosquitoes mix Stock Spray and Dip at the rate of 1 gallon to 40 gallons of water. Wet surfaces thoroughly, but not to the point of excessive runoff. Make application to outside walls of dairy barns, inside hog pens, on fences, garbage cans and similar places. Repeat applications as needed.

Do not use in poultry houses.

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Avoid contamination of feed and food stuffs.

**WARNING:** Do not spray or dip calves under 6 months of age. Do not store, pour or use near heat or open flame. Do not use in dairy barns or milk rooms, or on feed or forage to be fed to dairy animals. Do not dip animals within 60 days of slaughter or spray within 30 days of slaughter. Do not use on dairy cattle or dairy goats. Do not use on dogs or household pets. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

## DIRECTIONS

BEFORE USE READ DIRECTIONS, WARNING AND NOTICE STATEMENTS CAREFULLY

### MIXING INSTRUCTIONS

Add Stock Spray and Dip to required amount of water in accordance with dilution recommendations on this label. Agitate or stir the mixture thoroughly before each use.

**FOR SMALLER QUANTITIES**—One tablespoon of Stock Spray and Dip makes 1 gallon of spray or dip. **TO REPLENISH DIPPING VATS**—add Stock Spray and Dip, according to recommended dilutions—for all fresh water added to vat.

**CATTLE:** To kill Ticks, Horn Flies and Lice and to protect against reinfestation, mix Stock Spray and Dip at the rate of 1 gallon to 150 gallons of water. For best results thorough wetting of animals is necessary.

**SHEEP AND GOATS:** To control Lice and Sheep Ticks mix Stock Spray and Dip at the rate of 1 gallon to 500 gallons of water. One application is usually sufficient for control of these pests. For best results thorough wetting of animals is necessary. For sheep scab, mix Stock Spray and Dip at the rate of 1 gallon to 100 gallons of water and immerse sheep not less than ½ minute, ducking heads at least twice. When large numbers are dipped, the dip should be checked frequently by a vat-side test to assure the proper dipping strength. For control of Fleece Worms and Ticks dilute Stock Spray and Dip at the rate of 1 gallon to 150 gallons of water. Wet sheep thoroughly. Repeat application as needed.

**HOGS:** To control Lice dilute Stock Spray and Dip at rate of 1 gallon to 150 gallons of water. One application is usually sufficient to control this pest.

**CONTROL OF SARCOPTIC MANGE:** Beef cattle, hogs and sheep, mix Stock Spray and Dip at rate of 1 gallon to 100 gallons of water. Spray animals twice at a 14-day interval. For best results thorough wetting of animals is necessary.