

DE-PESTER

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

Apply DE-PESTER CHLORDANE 25G directly with lawn spreader or other commercial granular application equipment designed for lawn or turf use.

FOR LAWN INSECTS: To control ants (leaf-cutting ants, lawn ants and harvester ants) and armyworms, apply 1/5 pound of 25% Granular per 1,000 square feet (10 lbs. per acre). For boxelder bugs, chinch bugs, chiggers and ticks, apply 1/5 to 1/2 pound per 1,000 sq. ft. (10 to 15 lbs. per acre) of lawn or turf; for boxelder bugs make applications to base and trunks of boxelder trees, along foundations of buildings and other protected places.

To control Japanese beetle grubs, white grubs, wireworms and white tringe beetle, apply 1/2 pound of DE-PESTER CHLORDANE 25G per 1,000 sq. ft. (15 lbs. per acre) for a single season control. 4 to 5 years control can be obtained by using 1/2 to 1 pound per 1,000 sq. ft. (22 to 44 lbs. per acre). Wash the material into the turf following application.

FOR CRABGRASS CONTROL: Apply 6 pounds CHLORDANE 25G per 1,000 sq. ft. of lawn or turf to suppress crabgrass seedlings as germination or sprouting occurs. It is for use as an early season preventative treatment and is not effective for the control of crabgrass plants. Timing of application is important and should be applied before any germination which in many areas is during the period April 15 to May 1. This treatment will not burn or injure established lawns. Do not use on newly seeded lawns.

For Corn rootworm larva control on field corn, sweet corn and popcorn, apply 4 pounds DE-PESTER CHLORDANE G-25 per acre as a row treatment at planting time or 8 pounds DE-PESTER CHLORDANE G-25 per acre as a broadcast application at planting time.

For Potatoes: For the control of wireworms, apply 10 to 24 pounds of DE-PESTER CHLORDANE G-25 per acre at broadcast application and work into soil prior to or at planting time. For row application, use 2-1/2 to 3-1/3 pounds of DE-PESTER CHLORDANE G-25 per acre applied to the row at planting time.

For Sweet Potatoes to control flea beetle larva, white grubs and wireworms, apply 10 to 20 pounds of DE-PESTER CHLORDANE G-25 into the top few inches of soil. Apply only as pre-plant, planting or transplanting treatment.

CAUTION

Avoid inhalation. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Do not use on animals or pets. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying wastes, or by burning (Keep out of smoke).

This product is toxic to fish and wildlife.

Keep out of lakes, streams, and ponds.

Birds feeding on treated areas may be killed.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CHLORDANE G-25

CAUTION Keep Out of Reach of Children
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

CONTAINS 25% CHLORDANE

ACTIVE INGREDIENT:

Technical Chlordane† 25%

INERT INGREDIENTS:

..... 75%

..... 100%

†(Equivalent to 15% Octachloro-4, 7-methanotetrahydroindane and 10% of related compounds.)

USDA Reg. No. 148-602

2-69



APR 15 1964
FOR INFORMATION
ED UNDER NO
TO ATTACHED COMMENT 148-602